

EARTH RESOURCES

A CONTINUING BIBLIOGRAPHY WITH INDEXES

ISSUE 5

OCTOBER 1975

PREVIOUS EARTH RESOURCE BIBLIOGRAPHIES

Remote Sensing of Earth Resources	(NASA SP-7036(01))
Earth Resources	(NASA SP-7041(01))
Earth Resources	(NASA SP-7041(02))
Earth Resources	(NASA SP-7041(03))
Earth Resources	(NASA SP-7041(04))

This bibliography was prepared by the NASA Scientific and Technical Information Facility operated for the National Aeronautics and Space Administration by Informatics Information Systems Company.

EARTH RESOURCES

A Continuing Bibliography With Indexes Issue 5

A selection of annotated references to unclassified reports and journal articles that were introduced into the NASA scientific and technical information system and announced between January 1975 and March 1975 in

- Scientific and Technical Aerospace Reports (STAR)
- International Aerospace Abstracts (IAA)



This Supplement is available from the National Technical Information Service (NTIS), Springfield, Virginia 22161, for \$4.00. For copies mailed to addresses outside the United States, add \$2.50 per copy for handling and postage.

INTRODUCTION

The technical literature described in this continuing bibliography may be helpful to researchers in numerous disciplines such as agriculture and forestry, geography and cartography, geology and mining, oceanography and fishing, environmental control, and many others. Until recently it was impossible for anyone to examine more than a minute fraction of the earth's surface continuously. Now vast areas can be observed synoptically, and changes noted in both the earth's lands and waters, by sensing instrumentation on orbiting spacecraft or on aircraft.

This literature survey lists 601 reports, articles, and other documents announced between January and March 1975 in Scientific and Technical Aerospace Reports (STAR), and International Aerospace Abstracts (IAA).

The coverage includes documents related to the identification and evaluation by means of sensors in spacecraft and aircraft of vegetation, minerals, and other natural resources, and the techniques and potentialities of surveying and keeping up-to-date inventories of such riches. It encompasses studies of such natural phenomena as earthquakes, volcanoes, ocean currents, and magnetic fields; and such cultural phenomena as cities, transportation networks, and irrigation systems. Descriptions of the components and use of remote sensing and geophysical instrumentation, their subsystems, observational procedures, signature and analyses and interpretive techniques for gathering data are also included. All reports generated under NA SA's Earth Resources Survey Program for the time period covered in this bibliography will also be included. The bibliography does not contain citations to documents dealing mainly with satellites or satellite equipment used in navigation or communication systems, nor with instrumentation not used aboard aerospace vehicles.

The selected items are grouped in nine categories. These are listed in the Table of Contents with notes regarding the scope of each category. These categories were especially chosen for this publication, and differ from those found in STAR and IAA.

Each entry consists of a standard bibliographic citation accompanied by an abstract. The citations and abstracts are reproduced exactly as they appeared originally in STAR, or IAA, including the original accession numbers from the respective announcement journals. This procedure, which saves time and money, accounts for the variation in citation appearance.

Under each of the nine categories, the entries are presented in one of two groups that appear in the following order:

IAA entries identified by accession number series A75-10,000 in ascending accession number order.

STAR entries identified by accession number series N75-10,000 in ascending accession number order:

After the abstract section, there are five indexes:

subject, personal author, corporate source, contract number and report/accession number.

AVAILABILITY OF CITED PUBLICATIONS

IAA ENTRIES (A75-10000 Series)

All publications abstracted in this Section are available from the Technical Information Service. American Institute of Aeronautics and Astronautics, Inc. (AIAA), as follows: Paper copies are available at \$5.00 per document up to a maximum of 20 pages. The charge for each additional page is 25 cents. Microfiche⁽¹⁾ are available at the rate of \$1.50 per microfiche for documents identified by the "#" symbol following the accession number. A number of publications, because of their special characteristics, are available only for reference in the AIAA Technical Information Service Library. Minimum airmail postage to foreign countries is \$1.00. Please refer to the accession number, e.g. (A75-18063), when requesting publications.

STAR ENTRIES (N75-10000 Series)

One or more sources from which a document announced in STAR is available to the public is ordinarily given on the last line of the citation. The most commonly indicated sources and their acronyms or abbreviations are listed below. If the publication is available from a source other than those listed, the publisher and his address will be displayed on the availability line or in combination with the corporate source line.

Avail: NTIS. Sold by the National Technical Information Service to U.S. customers at the price shown in the citation following the letters HC (hard, paper, or facsimile copy). Customers outside the U.S. should add \$2.50 per copy for handling and postage charges to the price shown. (Prices shown in earlier STAR volumes, 1962-1974, have been superseded but may be calculated from the number of pages shown in the citation. The price schedule by page count was given in the last STAR issue of 1974 or may be obtained from NTIS.)

Microfiche⁽¹⁾ are available at a standard price of \$2.25 (plus \$1.50 for non-U.S. customers) regardless of age for those accessions followed by a "#" symbol. Accession numbers followed by a "+" sign are not available as microfiche because of size or reproducibility.

Initially distributed microfiche under the NTIS SRIM (Selected Research in Microfiche) is available at greatly reduced unit prices. For this service and for information concerning subscription to NASA printed reports, consult the NTIS Subscription Unit.

NOTE ON ORDERING DOCUMENTS: When ordering NASA publications (those followed by the "*" symbol), use the N accession number.

NASA patent applications (only the specifications are offered) should be ordered by the US-Patent-Appl-SN number.

Non-NASA publications (no asterisk) should be ordered by the AD, PB, or other *report* number shown on the last line of the citation not by the N accession number. It is also advisable to cite the title and other bibliographic identification.

Avail: SOD (or GPO). Sold by the Superintendent of Documents, U.S. Government Printing Office, in hard copy. The current price and order number are given following the availability line. (NTIS will fill microfiche requests, at the standard \$2.25 price, for those documents identified by a "#" symbol.)

⁽¹⁾ A microfiche is a transparent sheet of film. 105 by 148mm in size containing as many as 60 to 98 pages of information reduced to micro images (not to exceed 26:1 reduction).

- Avail: NASA Public Document Rooms. Documents so indicated may be examined at or purchased from the National Aeronautics and Space Administration, Public Documents Room (Room 126), 600 Independence Ave., S.W., Washington, D.C. 20546, or public document rooms located at each of the NASA research centers, the NASA Space Technology Laboratories, and the NASA Pasadena Office at the Jet Propulsion Laboratory.
- Avail: ERDA Depository Libraries. Organizations in U.S. cities and abroad that maintain collections of Energy Research and Development Administration reports, usually in microfiche form, are listed in *Nuclear Science Abstracts*. Services available from the ERDA and its depositories are described in a booklet, *Science Information Available from the Energy Research and Development Administration* (TID-4550), which may be obtained without charge from the ERDA Technical Information Center.
- Avail: Univ. Microfilms. Documents so indicated are dissertations selected from Dissertation Abstracts and are sold by University Microfilms as xerographic copy (HC) at \$10.00 each and microfilm at \$4.00 each regardless of the length of the manuscript. Handling and shipping charges are additional. All requests should cite the author and the Order Number as they appear in the citation.
- Avail: USGS. Originals of many reports from the U.S. Geological Survey, which may contain color illustrations, or otherwise may not have the quality of illustrations preserved in the microfiche or facsimile reproduction, may be examined by the public at the libraries of the USGS field offices whose addresses are listed in this Introduction. The libraries may be queried concerning the availability of specific documents and the possible utilization of local copying services, such as color reproduction.
- Avail: HMSO. Publications of Her Majesty's Stationery Office are sold in the U.S. by Pendragon House, Inc. (PHI), Redwood City, California. The U.S. price (including a service and mailing charge) is given, or a conversion table may be obtained from PHI.
- Avail: BLL (formerly NLL): British Library Lending Division, Boston Spa, Wetherby, Yorkshire, England. Photocopies available from this organization at the price shown. (If none is given, inquiry should be addressed to the BLL.)
- Avail: ZLDI. Sold by the Zentralstelle für Luftfahrtdokumentation und -Information, Munich, Federal Republic of Germany, at the price shown in deutschmarks (DM).
- Avail: Issuing Activity, or Corporate Author, or no indication of availability. Inquiries as to the availability of these documents should be addressed to the organization shown in the citation as the corporate author of the document.
- Avail: U.S. Patent Office. Sold by Commissioner of Patents, U.S. Patent Office, at the standard price of 50 cents each, postage free.
- Other availabilities: If the publication is available from a source other than the above, the publisher and his address will be displayed entirely on the availability line or in combination with the corporate author line.

ADDRESSES OF ORGANIZATIONS

American Institute of Aeronautics and Astronautics
Technical Information Service
750 Third Ave.
New York, N.Y. 10017

British Library Lending Division, Boston Spa, Wetherby, Yorkshire, England

Commissioner of Patents U.S. Patent Office Washington, D.C. 20231

Energy Research and Development Administration Technical Information Center P.O. Box 62 Oak Ridge, Tennessee 37830

ESA - Space Documentation Service ESRIN Via Galileo Galilei 00044 Frascati (Rome), Italy.

Her Majesty's Stationery Office P.O. Box 569, S.E. 1 London, England

NASA Scientific and Technical Information Facility P.O. Box 8757 B.W.I. Airport, Maryland 21240

National Aeronautics and Space Administration Scientific and Technical Information Office (KSI) Washington, D.C. 20546

National Technical Information Service Springfield, Virginia 22161

Pendragon House, Inc. 899 Broadway Avenue Redwood City, California 94063

Superintendent of Documents U.S. Government Printing Office Washington, D.C. 20402

University Microfilms A Xerox Company 300 North Zeeb Road Ann Arbor, Michigan 48106

University Microfilms, Ltd. Tylers Green London, England

U.S. Geological Survey 1033 General Services Administration Bldg. Washington, D.C. 20242

U.S. Geological Survey 601 E. Cedar Avenue Flagstaff, Arizona 86002

U.S. Geological Survey 345 Middlefield Road Menlo Park, California 94025

U.S. Geological Survey Bldg. 25, Denver Federal Center Denver, Colorado 80225


Zentralstelle für Luftfahrtdokumentation und -Information 8 München 86 Postfach 880 Federal Republic of Germany

TABLE OF CONTENTS


Subject Categories

Abstracts in the bibliography are grouped under the following categories:	page:
O1 AGRICULTURE AND FORESTRY Includes crop forecasts, crop signature analysis, soil identification, disease detection, harvest estimates, range resources, timber inventory, forest fire detection, and wildlife migration patterns.	1
02 ENVIRONMENTAL CHANGES AND CULTURAL RESOURCES Includes land use analysis, urban and metropolitan studies, environmental impact, air and water pollution, geographic information systems, and geographic analysis.	11
03 GEODESY AND CARTOGRAPHY Includes mapping and topography.	27
04 GEOLOGY AND MINERAL RESOURCES Includes mineral deposits, petroleum deposits, spectral properties of rocks, geological exploration, and lithology.	33
05 OCEANOGRAPHY AND MARINE RESOURCES Includes sea-surface temperature, ocean bottom surveying imagery, drift rates, sea ice and icebergs, sea state, fish location.	41
O6 HYDROLOGY AND WATER MANAGEMENT Includes snow cover and water runoff in rivers and glaciers, saline intrusion, drainage analysis, geomorphology of river basins, land uses, and estuarine studies.	47
07 DATA PROCESSING AND DISTRIBUTION SYSTEMS Includes film processing, computer technology, satellite and aircraft hardware, and imagery.	53
08 INSTRUMENTATION AND SENSORS Includes data acquisition and camera systems and remote sensors.	63
09 GENERAL Includes economic analysis.	75
SUBJECT INDEX PERSONAL AUTHOR INDEX CORPORATE SOURCE INDEX CONTRACT NUMBER INDEX REPORT/ACCESSION INDEX	B-1 C-1 D-1

TYPICAL CITATION AND ABSTRACT FROM STAR



TYPICAL CITATION AND ABSTRACT FROM IAA





EARTH RESOURCES

A Continuing Bibliography (Issue 5)

OCTOBER 1975

01 AGRICULTURE AND FORESTRY

Includes crop forecasts, crop signature analysis, soil identification, disease detection, harvest estimates, range resources, timber inventory, forest fire detection, and wildlife migration patterns.

A75-12033 # A comparison of Skylab and ERTS data for agricultural crop and natural vegetation interpretation. R. I. Welch, L. R. Pettinger, and C. E. Poulton (Earth Satellite Corp., Berkeley, Calif.). American Institute of Aeronautics and Astronautics and American Geophysical Union, Conference on Scientific Experiments of Skylab, Huntsville, Ala., Oct. 30-Nov. 1, 1974, AIAA Paper 74-1219, 12 p.

Images from the Skylab Earth Resources Experimental Package (EREP) system are compared with those obtained from the multispectral-scanner on the Earth Resources Technology Satellite (ERTS). The study is aimed at a comparative assessment of the usefulness of the EREP and ERTS remote sensing systems for identification and evaluation of-vegetation resources by testing the ability of photo interpreters to make the desired interpretations on selected types of EREP and ERTS imagery.

M.V.E.

A75-12872 Agricultural inventory techniques with orbital and high-altitude imagery. C. M. Hay (California, University, Berkeley, Calif.). (American Congress of Surveying and Mapping and American Society of Photogrammetry, National Fall Technical Convention, Columbus, Ohio, Oct. 11-14, 1972.) Photogrammetric Engineering, vol. 40, Nov. 1974, p. 1283-1293. NASA Order R-03-038-002.

An investigation was conducted regarding the feasibility of an employment of satellite imagery for land-use stratifications, taking into account present operational systems. It was found that satellite imagery of the quality of ERTS-1 can offer a feasible solution to the problem of out-of-date stratification. Other studies reported are concerned with the level of accuracy obtainable in a crop inventory for a specific crop using high-altitude aircraft photographs. G.R.

A75-15180 # Microwave emission characteristics of moist soils (Kharakteristiki izlucheniia vlazhnykh gruntov v SVCh diapazone). A. E. Popov, E. A. Sharkov, and V. S. Etkin (Akademiia Nauk SSSR, Institut Kosmicheskikh Issledovanii, Moscow, USSR). Meteorologiia i Gidrologiia, Oct. 1974, p. 49-57. 9 refs. In Russian.

The emissivity and polarization coefficient of moist-soil is calculated in the wavelength range from 0.8 to 226 cm for moisture up to 30 %. The calculations are based on experimental relationships

between the electrical parameters of the soil (attenuation factor and permittivity) and the moisture content. The effect of firmament illumination on the radio brightness temperature and the polarization coefficient of moist-soil emission is considered. It is shown that the optimal wavelength range for passive remote probing of moist soil is 20-75 cm, depending on the desired probing depth.

A.T.S.

A75-16241 Vegetation canopy reflectance. J. E. Colwell (Michigan, Environmental Research Institute, Ann Arbor, Mich.). Remote Sensing of Environment, vol. 3, no. 3, 1974, p. 175-183. 13 ref.

Possible cause-effect relationships in producing vegetation canopy reflectance are discussed. Hemispherical reflectance and even bidirectional reflectance measurements are shown to be inadequate to predict or understand vegetation canopy reflectance in many situations. Among the additional important parameters necessary for prediction and understanding of vegetation canopy reflectance are leaf hemispherical transmittance, leaf area and orientation, characteristics of other components of the vegetation canopy (stalks, trunks, limbs), soil reflectance, solar zenith angle, look angle, and azimuth angle. The effects of these parameters on vegetation canopy bidirectional spectral reflectance are described. (Author)

A75-16242 Active microwave measurement of soil water content. F. T. Ulaby, J. Cihlar, and R. K. Moore (University of Kansas Center for Research, Inc., Lawrence, Kan.). Remote Sensing of Environment, vol. 3, no. 3, 1974, p. 185-203. 22 refs. Contract No. NAS9-10261.

Measurements of radar backscatter from bare soil at 4.7, 5.9, and 7.1 GHz for incident angles of 0-70 deg have been analyzed to determine sensitivity to soil moisture. Because the effective depth of penetration of the radar signal is only about one skin depth, the observed signals were correlated with the moisture in a skin depth as characterized by the attenuation coefficient (reciprocal of skin depth). Since the attenuation coefficient is a monotonically increasing function of moisture density, it may also be used as a measure of moisture content over the distance involved, which varies with frequency and moisture content. The measurements show an approximately linear increase in scattering with attenuation coefficient of the soil at angles within 10 deg of vertical and all frequencies. At 4.7 GHz this increase continues relatively large out to 70 deg incidence, but by 7.1 GHz the sensitivity is much less even at 20 deg and practically gone at 50 deg. (Author)

A75-16682 Terrain measurement from holographic stereograms. J. T. McCrickerd (Northrop Corp., Hawthorne, Calif.). In: Coherent optics in mapping; Proceedings of the Tutorial Seminar and Technology Utilization Program, Rochester, N.Y., March 27-29, 1974. Palos Verdes Estates, Calif., Society of Photo-optical Instrumentation Engineers, 1974, p. 191-198. Grant No. DAAK02-73-C-0037.

01 AGRICULTURE AND FORESTRY

A study to determine the feasibility of measuring terrain from aerial stereo-photos displayed in the form of holographic stereograms. To compare left and right photos of a stereo-pair, a property of holograms is used, which is that two or more focused images can be superimposed on the same film, yet separately viewed under properly directed illumination. By laying one picture on the other, the distance is minimized between the two images of a feature, thereby reducing errors when simple measurement apparatus is used. A stereo-pair of a coordinate grid is also overlapped onto the stereogram so that all data can be differentially measured. For best results a set of grid points which are referred to points of known geographical coordinates is used.

A75-17104 # An examination of the extent of fire in the grassland and savanna of Africa along the southern side of the Sahara. W. Deshler (Maryland, University, College Park, Md.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 23-30.

A75-17106 # A method of specifying remotely sensed units for soil sample points. G. A. May, G. W. Petersen, F. Y. Borden, and D. N. Applegate (Pennsylvania State University, University Park, Pa.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 83-89. 6 refs.

Three sites were selected from Berks (shale-derived), Duffield (limestone-derived), and Penn (sitstone-derived) soils. A uniform training area was selected within each of these sites. The multispectral scanner data within each training area were analyzed by computer programs. Soil samples were collected within each training area. The angle and distance of each sampling point from the constructed base line was determined by a stadia rod. The angles and distances were input to a remotely sensed unit (RSU) identification program that outputs the scan line and element number (RSU) of each sampling point. Results from this program indicated that a high percentage of soil sampling locations were-within-previously designated training areas and that the duplication of soil samples within an RSU occurred at only one site.

A75-17113 # A clustering algorithm for unsupervised crop classification. L. Borriello (CSATA, Bari, Italy) and F. Capozza (Telespazio S.p.A., Rome, Italy). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 181-188, 5 refs.

In heuristic clustering algorithms, the initial value selected for cluster centers and thresholds has often a significant impact on the result which is going to be obtained, i.e., on the final position of the clusters and their number. Therefore, the initial values for the cluster centers must be known to a fairly good extent to ensure successful convergence of the algorithm to the true cluster centers. Adequate thresholds must further be selected in order to prevent scattered observations from affecting the convergence process. The algorithm described in this paper automatically determines the initial value for cluster centers and thresholds, by analyzing the histograms of the observations to be clustered. Thus, in this procedure the initial centers are close to the final ones and convergence is obtained in only a fraction of the iterations required by conventional heuristic clustering procedures. A table look-up procedure yields a further reduction in the computer time required to run this algorithm.

(Author)

A75-17122 * # Areal extent of snow in forested regions - A practical estimation technique using ERTS-1 data. W. C. Draeger and D. T. Lauer (California, University, Berkeley, Calif.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 333-339. 17 refs. Grant No. NGL-05-003-404.

A75-17143 # Evaluation of ERTS-1 and aircraft data for assessing internal drainage in irrigated agriculture. D. W. Ryland, F. A. Schmer, D. G. Moore (South Dakota State University, Brookings, S.D.), and W. A. Lidster (U.S. Bureau of Reclamation, Denver, Colo.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings... Volume 1.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 683-691. U.S. Bureau of Reclamation Contract No. 14-06-D-7358.

Research was conducted to investigate the application of remote sensing techniques for detecting areas with high water tables in irrigated agricultural lands. Aerial data were collected by the ERTS-1 satellite and aircraft over a portion of the Kansas/Bostwick Irrigation District in Republic County, Kansas. Areas having high water tables were visually located on the imagery and verified on the ground. Aircraft and ERTS-1 imagery were digitized and correlated with water table depths. Multiple regression, mode seeking, and K-class classification analyses were also applied to the data. Significant correlations were obtained between water table depths and the aircraft remotely sensed data both for the total area and for specific corn or fallow fields. For a corn field, 91% of the water table depths were correctly classified into depths less than 183 cm and depths greater than 183 cm. Significant correlations were obtained between ERTS-1 bands 5 and 7 and water table depth for both a corn and fallow field. (Author)

A75-17144 * # Use of visible, near-infrared, and thermal infrared remote sensing to study soil moisture. M. B. Blanchard (NASA, Ames Research Center, Moffett Field, Calif.), R. Greeley (Santa Clara, University, Santa Clara, Calif.), and R. Goettelman (LEE_Corp.,—Richmond; Va.).—In:—International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 693-700. 20 refs.

Two methods are used to estimate soil moisture remotely using the 0.4 to 14.0-micron wavelength region: (1) measurement of spectral reflectance, and (2) measurement of soil temperature. The reflectance method is based on observations which show that directional reflectance decreases as soil moisture increases for a given material. The soil temperature method is based on observations which show that differences between daytime and nighttime soil temperatures decrease as moisture content increases for a given material. In some circumstances, separate reflectance or temperature measurements yield ambiguous data, in which case these two methods may be combined to obtain a valid soil moisture determination. In this combined approach, reflectance is used to estimate low moisture levels; and thermal inertia (or thermal diffusivity) is used to estimate higher levels. The reflectance method appears promising for surface estimates of soil moisture, whereas the temperature method appears promising for estimates of near-subsurface (0 to 10 cm).

(Author)

A75-17145 * # Moisture detection from Skylab. J. R. Eagleman (Kansas, University, Lawrence, Kan.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 701-705. Contract No. NAS9-13273.

Five sets of data are described which relate the Skylab S193 radiometer and scatterometer data and S194 L-band radiometer temperature to soil moisture content of test sites in Texas and Kansas. Although the analysis of the S193 data are incomplete, analysis of the L-band radiometer data show that it is highly correlated with the soil moisture content of the surface layers of soil. Various correlations and predictions based on the relationship are described in this paper. (Author)

A75-17146 * # Remote detection of soil surface moisture. E. H. Stockhoff and R. T. Frost (GE Space Sciences Laboratory, Philadelphia, Pa.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 707-723. 8 refs. Contract No. NAS5-21689.

Polarimetric data concerning soil surface moisture were obtained during a series of flights over the Imperial Valley, California, and Phoenix, Arizona, during March 1972. A polarimeter was installed in NASA's Convair-990 aircraft, Galileo, above a window in the floor of the aft cargo compartment in such a manner that it could view from 42 deg ahead of, to 42 deg to the rear of the nadir. It had a 3 deg field of view and a 10 nm bandwidth centered at 641 nm. The moisture content of the solid surface for fields viewed by the polarimeter was measured by determining the angle through which the light had been scattered by the soil and by observing the degree of polarization of this light produced during its interaction with the soil. The polarimeter measures this degree of polarization in terms of Stokes parameters. Ground-truth samples of soil were obtained at several depths along the flight path. (Author)

A75-17158 * # Skylab S-192 ratio codes of soil, mineral, and rock spectra for ratio image selection and interpretation. R. K. Vincent and W. W. Pillars (Michigan, Environmental Research Institute, Ann Arbor, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 875-896. 10 refs. Contracts No. NAS9-13317; No. NAS9-13386.

A limited theoretical systems study is made of the Skylab S-192 12-channel multispectral scanner's capabilities for geological remote sensing. Laboratory spectra of rocks, minerals, soils, and some vegetation were used to rank the twelve channels and the best twelve of the 66 possible nonreciprocal spectral ratios according to their ability to discriminate among 17 classes of geological targets. Environmental and instrumental noise sources were ignored. The S-192 should be most useful for discriminating minerals deposited by hydrothermal alteration, weathering, and evaporation. Discrimination of igneous rocks will be difficult. Ratio images are recommended over automatic discrimination methods for those geological targets that can be enhanced by ratio imaging. Ratio codes were found for the twelve highest ranked ratios, for use in selecting the ratio images which will best enhance a target of interest. They can also be searched for false alarm candidates for a given target. A.T.S.

A75-17167 • # Preliminary results of fisheries investigation associated with Skylab-3. K. Savastano, E. Pastula, Jr., G. Woods (NOAA, National Marine Fisheries Service, Bay St. Louis, Miss.), and K. Faller (NASA, Johnson Space Center, Earth Resources Laboratory, Bay St. Louis, Miss.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1013-1042. 23 refs. NASA Project 240; NASA Order T-8217-B.

The purpose of the 15-month investigation now in the analysis phase is to establish the feasibility of utilizing remotely sensed data acquired from aircraft and satellite platforms to provide information concerning the distribution and abundance of oceanic gamefish. Data

from the test area, jointly acquired by private and professional fishermen and NASA and NOAA/NMFS elements, in the northeastern Gulf of Mexico has made possible the identification of significant environmental parameters for white marlin. Predictive models based on catch data and surface truth information have been developed and have demonstrated potential for reducing search significantly by identifying areas which have a high probability of being productive. Three of the parameters utilized by the model, chlorophylla, sea surface temperature and turbidity have been inferred from aircraft sensor data. (Author)

A75-17168 # Rapid stock assessment of pilchard populations by aircraft-borne remote sensors. D. L. Cram (Department of Industries, Sea Fisheries Branch, Cape Town, Republic of South Africa). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1043-1050. 5 refs.

A75-17169 # Determining range condition and forage production potential in California from ERTS-1 imagery. D. M. Carneggie and S. D. DeGloria (California, University, Berkeley, Calif.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1051-1059.

A75-17170 # Grass canopy bidirectional spectral reflectance. J. E. Colwell (Michigan, Environmental Research Institute, Ann Arbor, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1061-1085.

The green (550 nm), red (650 nm), and near IR (750 nm) bidirectional reflectance of grass canopies was studied both theoretically and empirically, in order to determine the feasibility of using remote sensing techniques to assess the standing biomass of grasslands. The investigation showed that the optimum spectral bands for remote determination of standing biomass of grasslands vary, depending on such things as the type of vegetation, the range of values of percent vegetation cover present, the soil reflectance, and the look angle and solar zenith angles. No single spectral band can be considered, a priori, to be effective in all situations. (Author)

A75-17171 # Forest defoliation assessment with satellite imagery. W. G. Rohde (Earth Satellite Corp., Washington, D.C.) and H. J. Moore (U.S. Department of Agriculture, Animal and Plant Health Inspection Service, Hyattsville, Md.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1089-1104. 7 refs.

A study of ERTS-1 multispectral scanner imagery (MSS) indicates that forest defoliation by insects such as the gypsy moth can be detected from satellite imagery and correlates well with aerial visual survey data. Two damage classes (heavy and moderate to light) and undefoliated areas can be detected by human interpreters from false-color composite imagery. The use of earth resources satellites in an integrated pest detection and control system will require more cloud-free images and more timely data acquisition and dissemination than were possible with ERTS-1.

A.T.S.

A75-17172 # Spectral reflectance studies on mineral deficiency in corn plants. H. A. Younes, A. G. Abdel-Samie (National Research Centre, Giza, Egypt), R. M. Abdel-Aal (Ministry of Agriculture, Soils and Water Research Institute, Giza, Egypt), and M. M. Khodair (National Institute for Standards, Giza, Egypt). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2.

Ann Arbor, Mich., Environmental Research

Institute of Michigan, 1974, p. 1105-1125. Research supported by the Oklahoma State University and NSF.

A sand-culture experiment was conducted to study the possibility of using spectrophotometric reflectance measurements on plant leaves and photography with different films to detect early symptoms of N, P, and K deficiency in corn plants. Experimental procedures, spectrophotometric measurements on single leaves and photography techniques are described and the results are presented in tabulated and illustrated forms. The data revealed that normal and deficient nutrient plants could be differentiated both photographically and photometrically. The differentiation depends on plant age, nutrient element, and the severity of nutrient deficiency. A detectable and noticeable difference in the near infrared spectrum between the nutrient deficient and normal leaves occurred, before any visual symptoms appeared. Photography proved that the best results can be obtained using false color infrared films with No. 25 filter. (Author)

A75-17173 # Agricultural resources investigations in Southern France and Northern Italy. In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1127-1159

A joint French-Italian study was made to evaluate the feasibility of using satellite and aircraft remote sensing to support agronomic research under European ecological conditions. Areas chosen for study included irregated rice fields and natural (beech forests) and artificial (poplar plantations and coniferous afforestations) forested areas. Eight test sites in France, Italy, and Madagascar were chosen to represent a wide range of climatic conditions. Results on two of the sites are treated. The use of ERTS multispectral scanner imagery makes it possible to evaluate total rice cultivation area and distinguish different varieties of rice. Moreover, it should be possible to distinguish rice varieties from other crops and to distinguish rice fields from natural bodies of water. Studies were also conducted on estimating rice yield, detecting diseased fields, and inventorying poplar plantations by remote sensing.

A75-17174 # Thermal behaviour of some rice fields affected by a yellows-type disease. C. De Carolis, G. Baldi (Ente Nazionale Risi, Mortara, Italy); S. Galli de Paratesi (Comitato Nazionale per l'Energia Nucleare, Centro per le Ricerche Comuni, Ispra, Italy), and G. M. Lechi (CNR, Milan, Italy). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1161-1170. 10 refs. Research sponsored by the Centro per le Ricerche Comuni.

Ground-based and airborne near-infrared and thermal-infrared surveys were made to study the radiation behavior of rice fields affected by 'giallume', a yellows-type disease. The near-infrared aerial surveys allowed exact measurement of the vegetative covering of water. The diseased rice areas were found to behave like thermally anomalous areas when compared to healthy areas. The black-body temperature of the diseased areas exceeds that of the healthy areas by several degrees K.

A.T.S.

A75-17175 * # First results from the crop identification technology assessment for remote sensing /CITARS/. F. G. Hall (NASA, Johnson Space Center, Houston, Tex.), M. E. Bauer (Purdue University, West Lafayette, Ind.), and W. A. Malila (Michigan, Environmental Research Institute, Ann Arbor, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1171-1191. 15 refs.

The CITARS task design, including objectives, analysis methodology and experimental procedures, is described and first results from this effort are presented. The extensive ground truth

data set acquired for the CITARS task is described and discussed in some detail. Results of the accuracy tests for the photo interpretative CITARS ground truth are given. Results of the assessment of the ERTS MSS data for cloud cover and electronic quality are presented. Some results of the geometric correction and registration of the time sequential CITARS ERTS data are given. Finally, the field boundary selection problem is addressed and the results of the use of new technology for boundary selection are presented. (Author)

A75-17176 * # ERTS-I data for classifying native plant communities - Central Colorado. R. S. Driscoll, R. E. Francis (U.S. Forest Service, Fort Collins, Colo.), J. A. Smith, and R. A. Mead (Colorado State University, Fort Collins, Colo.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1195-1211. 10 refs. NASA Order S-70251-AG.

A75-17185 # Color terrain map of Yellowstone National Park, computer-derived from ERTS MSS data. R. R. Root (National Park Service, Denver, Colo.), H. W. Smedes (U.S. Geological Survey, Denver, Colo.), N. E. G. Roller (Michigan, Environmental Research Institute, Ann Arbor, Mich.), and D. Despain (National Park Service, Yellowstone National Park, Wyo.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1369-1398. 11 refs. Research supported by the U.S. Geological Survey.

A75-17186 # Inventories of Delaware's coastal vegetation and land-use utilizing digital processing of ERTS-1 imagery. V. Klemas, D. Bartlett (Delaware, University, Newark, Del.), R. Rogers, and L. Reed (Bendix Corp., Aerospace Systems Div., Ann Arbor, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1399-1410. 6 refs.

A75-17187 # Texture analysis with Fourier series. H. Maurer (Zürich, University, Zurich, Switzerland): In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1411-1420. 14 refs. Translation.

Concerning a semi-automated or automated classification system for agricultural crop types from large-scale color aerial photographs, texture leads to excellent parameters for the discrimination. With a single Fourier series analysis textural parameters were computed from average density profiles through crop fields, measured with a ZEISS-Scanning-Microscope-Photometer. Then these parameters were tested with a discriminant analysis for their usefulness for the classification of agricultural crop types. The results are very promising. The combination of the measuring data from two different bands produces a perfect classification of all crop fields.

(Author)

A75-17189 # Investigation into the spectral signature of agricultural crops during their state of growth. T. A. de Boer (Instituut voor Biologisch en Scheikundig Onderzoek van Landbouwgewässen, Wageningon, Netherlands), N. J. J. Bunnik, H. W. J. van Kasteren, D. Uenk, W. Verhoef (Netherlands Interdepartmental Working Community for the Application of Remote Sensing Techniques, Delft, Netherlands), and G. P. de Loor (Physisch Laboratorium TNO, The Hague, Netherlands). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1441-1455. 15 refs. Research supported by the Physisch Laboratorium TNO.

The spectral signature from 400 to 2300 nm of 11 crops was determined at different phases of their growth. A fieldspectrometer, constructed according to a new principle and developed in the Netherlands, was used. To reduce the number of parameters that affect the directional reflectance of a canopy in its natural environment, the reflectance was measured only perpendicularly to the field surface while using an artificial light source. The purpose of this investigation was to determine, within the available atmospheric windows, the spectral bands in which the optimal differences between the signatures of different crops during their state of growth could be measured. (Author)

A75-17190 # Radar cross sections of vegetation canopies determined by monostatic and bistatic scatterometry. E. P. W. Attema, L..G. den Hollander (Technische Hogeschool Delft, Delft, Netherlands), T. A. de Boer, D. Uenk (Instituut voor Biologisch en Scheikundig Onderzoek van Landbouwgewassen, Wageningen, Netherlands), W. J. Eradus, G. P. de Loor (Physisch Laboratorium TNO, The Hague, Netherlands), H. van Kasteren, and J. van Kuilenburg (Netherlands Interdepartmental Working Community for the Application of Remote Sensing Techniques, Delft, Netherlands). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1457-1465. 10 refs.

A75-17216 * # Statistical separability of agricultural cover types in subsets of one to twelve spectral channels. R. Kumar and L. Silva (Purdue University, West Lafayette, Ind.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Acbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1891-1903, 18 refs. Research supported by the Instituto Nacional de Pesquisas Espaciais of Brazil; Grant No. NGL-15-005-112.

The purpose of this study was to determine the statistical separability of multispectral measurements from agricultural cover types: corn, soybeans, green forage (hay and pasture) and forest, in one to twelve spectral channels. Multispectral scanner data in twelve spectral channels in the wavelength range 0.4 to 11.7 microns, acquired for three flightlines were analysed by applying automatic pattern recognition techniques. The same analysis was performed for the data acquired a month later over the same three flightlines to investigate the effect of time on statistical separability of agricultural cover types. In the subsets of one to six spectral channels, the combination of wavelength regions (where V, N, M and T denote the visible, near infrared, middle infrared and thermal infrared wavelength regions, respectively): V, V M, V N M, V N M T, V V N M T, V V N M M T, respectively, were found to be the best choices for getting good overall statistical separability of the agricultural cover types for the data acquired. (Author)

A75-17973 * Microbial abundance and thermoluminescence of antarctic dry valley soils. J. D. Ingham, R. E. Cameron, and D. D. Lawson (California Institute of Technology, Jet Propulsion Laboratory, Pasadena, Calif.). Soil Science, vol. 117, no. 1, 1974, p. 46-57. 10 refs. NSF Grant No. C-585; Contract No. NAS7-100.

A75-18056 An intensive natural resources inventory aids land use planning and resources management in Puerto Rico. L. H. Liegel (U.S. Forest Service, Institute of Tropical Forestry, Rio Piedras, P.R.) and G. A. del Toro (Puerto Rico Department of Natural Resources, San Juan, P.R.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings.

Falls Church, Va., American Society of Photogrammetry, 1974, p. 57-65. 6 refs.

A75-18063 Geometronics-earth resource measurements and displays for the Forest Service. J. R. Swinnerton (U.S. Forest Service, Washington, D.C.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings. Falls Church, Va., American Society of Photogrammetry, 1974, p. 170-177.

More and better information is needed to support land use decisions on the National Forests. In response to this need, a new multilayer mapping system was recently adopted. The objective will be to incorporate special purpose mapping and graphic efforts into a coordinated system, tied to a common base. Concurrently, computer assisted systems are being implemented to streamline data handling. For example, after elevation data is properly formatted and stored, it can be analyzed with software routines which generate slope, aspect, contours, perspective, cross-sections or limits of viewed areas.

(Author)

A75-18066

The use of high altitude photography and ERTS-1 imagery for wildlife habitat inventory in Kansas. J. W. Merchant, Jr. (University of Kansas Center for Research, Inc., Lawrence, Kan.) and B. H. Waddell, Jr. (Kansas Forestry, Fish and Game Commission, Valley Falls, Kan.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings.

Falls Church, Va., American Society of Photogrammetry, 1974, p. 220-231, 8 refs.

A75-18068 Trends in the use of remote sensing for state resources and environmental management. F. J. Wobber (IBM Corp., Resources Management Dept., Gaithersburg, Md.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings. Falls Church, Va., American Society of Photogrammetry, 1974, p. 246-255

Remote sensing programs at the state level reflect many changes in remote sensing technology and particularly a trend towards practical problem-solving. Five state remote sensing case histories are described and analyzed for trends in remote sensing for state resources and environmental management. Trends including the growth of regulatory and legal applications, increased use of information systems and decision modeling, and modifications in user attitudes with respect to the scale of remote sensor records are noted. (Author)

A75-18073 Optical film density values from color IR photography for wetland soils mapping. S. T. Kim, A. J. Lewis, and P. E. Schilling (Louisiana State University, Baton Rouge, La.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings. Falls Church, Va., American Society of Photogrammetry, 1974, p. 323-331. 10 refs. Research supported by the Louisiana State University and Joint Legislative Committee on Environmental Quality of the State of Louisiana.

A75-18556 * Radar response to vegetation. F. T. Ulaby (University of Kansas Center for Research, Inc., Lawrence, Kan.). *IEEE Transactions on Antennas and Propagation*, vol. AP-23, Jan. 1975, p. 36-45. 14 refs. Contract No. NAS9-10261; Grant No. DAAK02-68-C-0089.

Active microwave measurements of vegetation backscatter were conducted to determine the utility of radar in mapping soil moisture through vegetation and mapping crop types. Using a truck-mounted boom, spectral response data were obtained for four crop types (corn, milo, soybeans, and alfalfa) over the 4-8 GHz frequency band, at incidence angles of 0 to 70 degrees in 10-degree steps, and for all four linear polarization combinations. Based on a total of 125 data sets covering a wide range of soil moisture, content, system design criteria are proposed for each of the aforementioned objectives.

Quantitative soil moisture determination was best achieved at the lower frequency end of the 4-8 GHz band using HH polarized waves in the 5- to 15-degree incidence angle range. A combination of low and high frequency measurements are suggested for classifying crop types. For crop discrimination, a dual-frequency dual-polarization (VV and cross) system operating at incidence angles above 40 degrees is suggested. (Author)

N75-10543*# Northern Prairie Wildlife Research Center, Jamestown, N. Dak.
UTILIZATION OF SKYLAB (EREP) SYSTEM FOR APPRAISING CHANGES IN CONTINENTAL MIGRATORY BIRD HABITAT Monthly Progress Report, Oct. 1974
David S. Gilmer, Principal Investigator Oct. 1974 4 p EREP (NASA Order T-4114-B)
(E74-10817; NASA-CR-140619) Avail: NTIS HC \$3.25 CSCL OBC

N75-10544*# Arizona Univ., Tucson. Office of Arid Lands

A STUDY TO EXPLORE THE USE OF ORBITAL REMOTE SENSING TO DETERMINE NATIVE ARID PLANT DISTRIBUTION Biannual Progress Report and Final Report

W. G. McGinnies, Principal Investigator, L. K. Lepley, E. F. Haase, J. S. Conn, H. B. Musick, and K. E. Foster Aug. 1974 38 p refs Original contains imagery. Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue, Sioux Falls, S. D. 57198 ERTS

(Contract NAS5-21812)

(E74-10818; NASA-CR-140620) Avail: NTIS HC \$3.75 CSCL 08F

The author has identified the following significant results. It is possible to determine, from ERTS imagery, native arid plant distribution. Using techniques of multispectral masking and extensive fieldwork, three native vegetation communities were defined and mapped in the Avra Valley study area. A map was made of the Yuma area with the aid of ground truth correlations between areas of desert pavement visible on ERTS images and unique vegetation types. With the exception of the Yuma soil-vegetation correlation phenomena, only very gross differentiations of desert vegetation communities can be made from ERTS data. Vegetation_communities_with_obvious-vegetation_density differences such as saguaro-paloverde, creosote bush, and riparian vegetation can be separated on the Avra Valley imagery while more similar communities such as creosote bush and saltbush could not be differentiated. It is suggested that large differences in vegetation density are needed before the signatures of two different vegetation types can be differentiated on ERTS imagery. This is due to the relatively insignificant contribution of vegetation to the total radiometric signature of a given desert scene. Where more detailed information concerning the vegetation of arid regions is required, large scale imagery is appropriate.

N75-10557* Environmental Research Inst. of Michigan, Ann Arbor.

UTILITY OF ERTS FOR MONITORING THE BREEDING HABIT OF MIGRATORY WATERFOWL

Edgar W. Work, Jr., David S. Gilmer (Northern Prairie Wildlife Res. Center), and A. T. Klett (Northern Prairie Wildlife Res. Center) In NASA. Goddard Space Flight Center Third ERTS Symp., Vol. 2 May 1974 p 102-115 refs

CSCL 06C

Waterfowl breeding-ground surveys conducted each year by the Bureau of Sport Fisheries and Wildlife extend over a vast regivn of the United States and Canada. Data from these surveys are used to estimate waterfowl production by means of a mathematical model. Counts of May and July ponds are some the variables used in this model. Annual production estimates are used to predict fall flights of ducks. This information is then used for establishing waterfowl hunting regulations. Work to date

indicates that satellite remote sensing techniques hold considerable promise for the accurate and rapid assessment of waterfowl breeding habitat, especially changes in pond numbers and distribution. Development of an operational system utilizing satellite sensors as a primary source of data appears to be a realistic goal for the future.

N75-10558* National Aeronautics and Space Administration. Lyndon B. Johnson Space Center, Houston, Tex. AGRICULTURE, FORESTRY, RANGE RESOURCES Robert B. MacDonald In its Third ERTS Symp., Vol. 2 May 1974 p 116-125 CSCL O8F

The necessary elements to perform global inventories of agriculture, forestry, and range resources are being brought together through the use of satellites, sensors, computers, mathematics, and phenomenology. Results of ERTS-1 applications in these areas, as well as soil mapping, are described. A.L.

N75-10577# Mathematical Applications Group, Inc., Elmsford, N Y

AN EXTENSION OF THE COMBINATORIAL GEOMETRY TECHNIQUE FOR MODELING VEGETATION AND TERRAIN FEATURES Final Contract Report

Joan Brooks, Ragini S. Murarka, Daniel Onuoha, Frank H. Rahn, and Herbert A. Steinberg Jun. 1974 52 p refs (Contract DAAD05-73-C-0537; DA Proj. 1G6-62708-D-05) (AD-782883; BRL-CR-159) Avail: NTIS CSCL 08/2

An extended version of the MAGI (Mathematical Applications Group, Inc.) Combinatorial Geometry technique has been developed which is capable of realistic modeling of trees and bushes. The method has been demonstrated photographically using the SYNTHAVISION process developed by MAGI. Although the basic modeling techniques are those of the standard Combinatorial Geometry method, major extensions have been introduced to reduce the strains on computer memory and computer time that are inherent in a complex description. The method will eventually be capable of integrating the tree and bush model wath models of terrain features, camouflage nets and BRL-COMGEOM descriptions of vehicles. These features of the program are in progress but not yet complete.

Author (GRA)

N75-12277*# National Aeronautics and Space Administration. Ames Research Center, Moffett Field, Calif. SYSTEMS STUDY OF AN AUTOMATED FIRE WEATHER DATA SYSTEM

Kenji Nishioka Oct. 1974 70 p refs (NASA-TM-X-62402) Avail: NTIS HC \$4.25 CSCL 14B

A sensor system applicable to an automated weather station was developed. The sensor provides automated fire weather data which correlates with manual readings. The equipment and methods are applied as an aid to the surveillance and protection of wildlands from fire damage. The continuous readings provided by the sensor system make it possible to determine the periods of time that the wilderness areas should be closed to the public to minimize the possibilities of fire.

N75-12394*# South Dakota State Univ., Brookings. Remote Sensing Inst.

DEVELOP TECHNIQUES AND PROCEDURES, USING MULTISPECTRAL SYSTEMS, TO IDENTIFY FROM REMOTELY SENSED DATA THE PHYSICAL AND THERMAL CHARACTERISTICS OF PLANTS AND SOIL Monthly Progress Report, Sep. 1974

Victor I. Myers, Principal Investigator 20 Oct. 1974 2 p

(Contract NAS9-13337)

(E75-10012; NASA-CR-140654) Avail: NTIS HC \$3.25 CSCL

N75-12402*# Pacific Southwest Forest and Range Experiment Station, Berkeley, Calif.

INVENTORY OF FOREST AND RANGELAND RESOURCES, INCLUDING FOREST STRESS Bimonthly Progress Report, 16 Sep. - 15 Nov. 1974

Robert C. Aldrich, Frederick P. Weber, and Richard S. Driscoll, Principal Investigators 18 Nov. 1974 7 p EREP

(NASA Order T-4106-B)

(E75-10032; NASA-CR-140762; BMPR-14) Avail: NTIS HC \$3.25 CSCL 02F

N75-12403*# South Dakota State Univ., Brookings. Remote Sensing Inst.

DEVELOP TECHNIQUES AND PROCEDURES, USING MULTISPECTRAL SYSTEMS, TO IDENTIFY FROM REMOTELY SENSED DATA THE PHYSICAL AND THERMAL CHARACTERISTICS OF PLANTS AND SOIL Monthly Progress Report, Oct. 1974

Victor I. Myers, Principal Investigator 20 Nov. 1974 2 p

(Contract NAS9-13337)

(E75-10033; NASA-CR-140763) Avail: NTIS HC \$3.25 CSCL 08F

N75-12405*# Kansas Univ. Center for Research, Inc., Lawrence. Atmospheric Science Lab.

DETECTION OF MOISTURE AND MOISTURE RELATED PHENOMENA FROM SKYLAB Monthly Progress Report, Sep. 1974

Joe R. Eagleman, Principal Investigator and Norman Hardy Sep. 1974 7 p EREP (Contract NAS9-13273)

(E75-10035; NASA-CR-140772; TR-239-17) Avail: NTIS HC \$3.25 CSCL 08B

NC \$3.25 C3CL 00B

N75-12414*# Agricultural Research Service, Weslaco, Tex. IRRIGATION SCHEDULING, FREEZE WARNING AND SOIL SALINITY DETECTING Monthly Progress Report, Oct. 1974

Craig L. Wiegand, Principal Investigator Oct. 1974 3 p EREP (NASA Order T-4105-B)

(E75-10046; NASA-CR-140783; MPR-10) Avail: NTIS HC \$3.25 CSCL 02C

N75-12419*# Lockheed Electronics Co., Houston, Tex. Aerospace Systems Div.

FEASIBILITY STUDY ASCS REMOTE SENSING/COMPLIANCE DETERMINATION SYSTEM Final Report I. E. Duggan, T. C. Minter, Jr., B. H. Moore, and C. T. Nosworthy Jan. 1973 137 p refs (Contract NAS9-12200)

(NASA-CR-134288; EO-126) Avail: NTIS HC \$5.75 CSCL 05B

A short-term technical study was performed by the MSC Earth Observations Division to determine the feasibility of the proposed Agricultural Stabilization and Conservation Service Automatic Remote Sensing/Compliance Determination System. For the study, the term automatic was interpreted as applying to an automated remote-sensing system that includes data acquisition, processing, and management.

N75-12421*# Minnesota Univ., Minneapolis. Space Science Center.

A STUDY OF MINNESOTA FORESTS AND LAKES USING DATA FROM EARTH RESOURCES TECHNOLOGY SATELLITES Twenty-four Month Progress Report, Jul. 1974 30 Jun. 1974 246 p refs

(Grant NGL-24-005-263)

(NASA-CR-140865) Avail: NTIS HC \$7.50 CSCL 08F

Highlights of research and practical benefits are discussed for the following projects which utilized ERTS 1 data to provide municipal, state, federal, and industrial users with environmental resource information for the state of Minnesota: (1) forest disease detection and control: (2) evaluation of water quality by remote sensing techniques: (3) forest vegetation classification and management: (4) detection of saline soils in the Red River Valley; (5) snowmelt flood prediction: (6) remote sensing applications to hydrology; (7) Rice Creek watershed project; (8) water quality in Lake Superior and the Duluth Superior Harbor; and (9) determination of Lake Superior currents from turbidity patterns.

A.A.D.

N75-12423*# ECON, Inc., Princeton, N.J.
THE VALUE OF IMPROVED (ERS) INFORMATION BASED
ON DOMESTIC DISTRIBUTION EFFECTS OF U.S. AGRICUL-

TURE CROPS
David F. Bradford, Harry H. Kelejian, Richard Brusch, Jonathan
Gross, Herbert Fishman, and Daniel Feenberg 31 Oct. 1974
114 p refs

(Contract NASw-2558)

(NASA-CR-141046; Rept-74-2001-5) Avail: NTIS HC \$5.25 CSCL 02B

The value of improving information for forecasting future crop harvests was investigated. Emphasis was placed upon establishing practical evaluation procedures firmly based in economic theory. The analysis was applied to the case of U.S. domestic wheat consumption. Estimates for a cost of storage function and a demand function for wheat were calculated. A model of market determinations of wheat inventories was developed for inventory adjustment. The carry-over horizon is computed by the solution of a nonlinear programming problem. and related variables such as spot and future price at each stage are determined. The model is adaptable to other markets. Results are shown to depend critically on the accuracy of current and proposed measurement techniques. The quantitative results are presented parametrically, in terms of various possible values NR. of current and future accuracies.

N75-12518 Joint Publications Research Service, Arlington, Va. AUTOMATED REMOTE MEASURING SYSTEM IN AGRO-METEOROLOGY

V. A. Kovalenko, A. D. Kleshchenko, and O. V. Virchenko *In its* Meteorology and Hydrol., No. 9, 1974 (JPRS-63569) 2 Dec. 1974 p 110-116 refs Transl into ENGLISH from Meteorol. Gidrol. (Moscow), no. 9, 1974 p 93-97

An automated data gathering and processing system for aerophotometric investigations of the planted fields of farm crops was studied. A functional diagram is presented and its operation is described. A brief description is given for the individual assemblies of the system and the set of programmed moduli for processing the incoming information.

N75-13333*# Kansas Univ. Center for Research, Inc., Lawrence, Space Technology Center.

DETECTION OF MOISTURE AND MOISTURE RELATED PHENOMENA FROM SKYLAB Monthly Progress Report, Oct. 1974

Joe R. Eagleman, Principal Investigator and Richard Sloan Oct. 1974 4 p EREP (Contract NAS9-13273)

(E75-10007; NASA-CR-140586; TR-239-18) Avail: NTIS HC \$3.25 CSCL 08H

N75-13345*# American Univ., Washington, D.C. Drought Analysis Lab.

UTILIZATION OF ERTS-1 IMAGERY IN CULTIVATION AND SETTLEMENT SITE IDENTIFICATION AND CARRYING CAPACITY ESTIMATES IN UPPER VOLTA AND NIGER

Priscilla Reining Nov. 1973 172 p refs Original contains imagery. Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue, Sioux Falls, S. D. 57198 ERTS

(Contract NAS5-21970)

(E75-10029; NASA-CR-140570) Avail: NTIS HC \$6.25 CSCL 08B

N75-13361*# Northern Prairie Wildlife Research Center, Jamestown, N. Dak.
UTILIZATION OF SKYLAB (EREP) SYSTEM FOR APPRAISING CHANGES IN CONTINENTAL MIGRATORY BIRD HABITAT Monthly Progress Report, Nov. 1974
David S. Gilmer, Principal Investigator Nov. 1974 2 p EREP (NASA Order T-4114-8)
(E75-10061; NASA-CR-141001) Avail: NTIS HC \$3.25 CSCL 06C

N75-13362*# Agricultural Research Service, Weslaco, Tex. A STUDY OF THE EARLY DETECTION OF INSECT INFESTATIONS AND DENSITY/DISTRIBUTION OF HOST PLANTS Progress Report, Jun. 1974

William G. Hart, Sammy J. Ingle, and M. R. Davis, Principal Investigators 30 Jun. 1974 2 p EREP

(NASA Order T-4109-B)

(E75-10062; NASA-CR-141002; PR-17) Avail: NTIS HC \$3.25 CSCL 08F

The author has identified the following significant results. Significant results have been obtained in the identification of citrus, sugarcane, winter vegetables, irrigated pastures, and unimproved pastures which contain brush. Land without vegetation, lakes, roads, and waterways can also be determined. Different densities of vegetation covering some cultivated areas are apparent. The practical applications of these results are many. The abundance of host plants of pests can be determined. Avenues of entry of pests can be plotted, facilitating control or preventing entry of pest species. The boundaries of areas to be guarantined can be accurately established after viewing the S-190B data. Better cultural methods can be employed such as planning where to plant certain crops that indirectly are detrimental to those already growing. This would relate to such factors as pesticide drift or alternate hosts of major pests.

N75-13363*# North Carolina State Univ., Raleigh. School of Physical and Mathematical Sciences.

UTILIZATION OF EREP DATA IN GEOLOGICAL EVALUATION REGIONAL PLANNING, FOREST MANAGEMENT, AND WATER MANAGEMENT IN NORTH CAROLINA Quarterly Progress Report, Sep. - Nov. 1974

Charles W. Welby, Principal Investigator 13 Dec. 1974 3 p

(Contract NAS9-13321)

(E75-10063; NASA-CR-141003) Avail: NTIS HC \$3.25 CSCL 08B

N75-13503*# National Aeronautics and Space Administration.
Ames Research Center, Moffett Field, Calif.
BESMEX: BERING SEA MARINE MAMMAL EXPERIMENT

G. Carleton Ray and Douglas Wartzok Oct. 1974 55 p refs Prepared in cooperation with Johns Hopkins Univ. Original contains color illustrations

(NASA-TM-X-62399) Avail: NTIS HC \$4.25 CSCL 06C

Predictive ecological models are being studied for the management and conservation of the walrus, and the bowhead whale in the Bering Sea. The influence of sea ice on the distribution, and carrying capacity of the area for these two mammals is to be investigated with the primary target species being the walrus. Remote sensing and radio tracking is considered a requirement for assessing the walrus ecosystem. F.O.S.

N75-14192*# Kansas Univ. Center for Research, Inc., Lawrence. Atmospheric Science Lab.

DETECTION OF MOISTURE AND MOISTURE RELATED PHENOMENA FROM SKYLAB Monthly Progress Report, Nov. 1974

Joe R. Eagleman, Principal Investigator and Wen C. Lin Nov. 1974 14 p EREP (Contract NAS9-13273)

(E75-10073: NASA-CR-141098: TR-239-20) Avail: NTIS HC \$3.25 CSCL 08H

The author has identified the following significant results. The high correlations between radiometric temperature and soil moisture content are shown to remain quite high for independent footprints of the S194 sensor. Since an analysis based on overlapping footprints had previously been reported with a high correlation, it was necessary to verify that the correlation did not arise from dependent data.

N75-14196*# South Dakota State Univ., Brookings. Remote Sensing Inst.

DEVELOP TECHNIQUES AND PROCEDURES, USING MULTISPECTRAL SYSTEMS, TO IDENTIFY FROM REMOTELY SENSED DATA THE PHYSICAL AND THERMAL CHARACERISTICS OF PLANTS AND SOIL Monthly

Progress Report, Nov. 1974
Victor I. Myers, Principal Investigator 20 Dec. 1974 2 p

(Contract NAS9-13337)

(E75-10077; NASA-CR-141094) Avail: NTIS HC \$3.25 CSCL

N75-14202*# Agricultural Research Service, Weslaco, Tex. IRRIGATION SCHEDULING, FREEZE WARNING AND SOIL SALINITY DETECTING Monthly Progress Report, Dec. 1974

Craig L. Wiegand, Principal Investigator Dec. 1974 5 p

(NASA Order T-4105-B)

(E75-10084; NASA-CR-141108; MPR-12) Avail: NTIS HC \$3.25 CSCL 02C

N75-14241 London Univ. (England). Bedford Coll. RECOGNITION AND INTERPRETATION OF SPECTRAL SIGNATURES OF VEGETATION FROM AIRCRAFT AND SATELLITE IMAGERY IN WESTERN QUEENSLAND, AUSTRALIA

Monica M. Cole, E. S. Owen-Jones, N. D. E. Custance, and T. E. Beaumont *In* ESRO European Earth Resources Satellite Expts. May 1974 p 243-287 refs Original contains color illustrations (For availability see N75-14216 05-43)

The spectral signatures of vegetation communities recognized on multispectral photography, at scales of 1:15,000 and 1:5000, in the Mary Kathleen-Cloncurry and Dugald River areas of Australia were evaluated with reference of specific environmental parameters, notably bedrock mineralization, and were compared with those detectable on the ERTS-1 satellite imagery. Overlays of the spectral signature units including those subjected to enhancement techniques were prepared and computer output histograms of the geochemical and biogeochemical data generated. The distribution of vegetation associations, soil and rock types along each transect was classified by an unsuper-

vised learning type of analysis using a hierarchical fusion-cluster method. Each individual sample along the transect was specified by its reflectance on the eight spectral bands provided by the photographic imagery and clustering was applied in both the images and measurement space. The results of both methods were compared, when it was found that a hybrid method gave the best results.

Author (ESRO)

N75-14247 International Inst. for Aerial Survey and Earth Sciences, Enschede (Netherlands).

PRELIMINARY RESULTS OF THE INTERPRETATION OF ERTS-1 IMAGERY FOR SOIL SURVEYS IN THE MERIDA REGION, SPAIN

F. W. Hilwig, D. Goosen, and D. Katsieris In ESRO European Earth Resources Satellite Expts. May 1974 p 349-361 refs (For availability see N75-14216 05-43)

The potential of ERTS-1 imagery for soil surveying was evaluated. The imagery was related to the results of an aerial survey training and research project carried out in the Merida region in Spain over the past six years. Both black and white and color imagery was studied using various bands and a number of filter and illumination combinations. Preliminary results seem promising. Major landforms and some subdivisions could be easily recognized. Water bodies, river courses, extensive areas of Miocene clays, and more recent coarse-textured deposits could be delineated and existing soil maps at scales up to 1:100,000 could be updated. ERTS-1 imagery promises to be very valuable for generalized or exploratory surveys at small scales up to 1:500,000, valuable for reconnaissance surveys at scales up to 1:100,000, and useful for surveys at scales up to 1:50,000 in conjunction with conventional aerial photointerpretation.

Author (ESRO)

N75-14251 Bristol Univ. (England).

GROUND TRUTH MEASUREMENTS IN RELATION TO AIRCRAFT AND SATELLITE STUDIES OF AGRICULTURAL LAND USE AND LAND CLASSIFICATION IN BRITAIN

L. F. Curtis and A. J. Hooper (Min. of Agr., London) In ESRO European Earth Resources Satellite Expts. May 1974 p 405-415

Examples are given of ground truth collection in relation to ERTS-1 and aircraft sensing data for southwest Britain. These demonstrate that the concept of a ground truth site will vary according to the user's objective and the size of the area under study. The range of ground-truth data necessary for evaluation of remote sensing data collection techniques in respect of land-classification and agricultural land use studies are summarized. Estimates of the manpower requirements are given on the basis of two study programs. General problems of ground data collection are considered in respect of timing of observations. crop height, crop roughness, crop color, perspective, soil exposure, agricultural treatments, etc. Specific problems of data recording are discussed in relation to selected crops. Standardization of ground truth recording techniques for agricultural and land classification studies is described.

Author (ESRO)

N75-14252 Consiglio Nazionale delle Ricerche, Milan (Italy). Lab. per la Geofisica della Litosfera.

AGRICULTURAL RESOURCES INVESTIGATIONS IN NORTHERN ITALY, SOUTHERN FRANCE AND MADAGAS-CAR

R. Cassinis, M. Guyader (Centre d'Etudes Spatiales des Rayonnements, Toulouse), and S. GalliDeParatesi (Joint Res. Centre, Ispra. Italy) In ESRO European Earth Resources Satellite Expts. May 1974 p 417-437 Original contains color illustrations

Under a program of collaboration between the Commission of the European Communities and various specialized national institutes, the project is designed to utilize data from ERTS-B and aircraft flights for research into rice growing and forestry.

Ground experiments have been concerned mainly with the preparation of bio-climatological measurements and automation of their collection, as well as the methods of recording ground truth. By microdensitometric processing of ERTS-1 images of the Camargue, it was possible to identify rice fields before sowing. Owning a radiometry campaign with lysimeters, a series of aircraft flights with an IR camera, a four-channel radiometer, a two-channel thermal IR scanner, and thermographic surveys, it was possible to obtain details of several problems affecting rice fields in Northern Italy. The preliminary results form a basis for the exploitation of ERTS-B data.

Author (ESRO)

N75-14253* National Aeronautics and Space Administration, Washington, D.C.

FUTURE US ACTIVITIES BASED ON EXPERIENCE GAINED FROM ERTS-1 AND EREP

G. Thorley In ESRO European Earth Resources Satellite Expts. May 1974 p 439-447

Current and future U.S. activities in the fields of land use, agriculture, forestry, and water resources are summarized. ESRO

N75-14258# Army Foreign Science and Technology Center, Charlottesville, Va.

MORPHOGRAPHIC AND MORPHOMETRIC ANALYSIS OF VEGETATIVE COVER PHOTOGRAPHS IN AERIAL PHOTOGRAPHY FOR THE MAPPING OF VEGETATION

B. V. Vinogradov 1 Nov. 1973 11 p refs Transl, into ENGLISH from the book "Krupnomasshtabnoe Kartografirovanie Rastitelnosti" Novosibirsk, Nauka, 1970 ρ 96-101

(AD-785932: FSTC-HT-23-1296-73) Avail: NTIS CSCL 08/2 Countour boundaries and transition areas provide the greatest amount of information to the eye in the interpretation of photo images; this principle can be applied to the mapping of vegetation by aerial photography. The article discusses a variety of morphographic landform classifications, their relation to geobotanical interpretation, and means and methods for gathering and processing statistical information from aerial photographs. The value of statistical methods in simplifying field work is stressed.

GRA

N75-15097*# Kanner (Leo) Associates, Redwood City, Calif.
PRELIMINARY DATA ON THE UHF RADIOMETRIC
RESEARCH OF SOIL WETNESS, CONDUCTED IN THE
USSR, IN 1974

A. Ye. Basharinov, L. F. Borodin, G. I. Chukhray, and A. M. Shutko Washington NASA Jan. 1975 15 p Transl. into ENGLISH of "Predvarite! nyve materialy SVCH radiometricheskikh issledovaniy vlazhnosti pochv. provedennykh v SSSR v techeniye 1974 g" report, (Moscow), Acad. of Sci. USSR, 1975 15 p (Contract NASw-2481)

(NASA-TT-F-16151) Avail: NTIS HC \$3.25 CSCL 08M

Aircraft studies of the radiation characteristics of moist soils were performed in several regions of the U.S.S.R.. Radiometers installed on board an IL-14 aircraft were used for obtaining radiation measurements. The characteristics of the regions and their agricultural-meteorological conditions during these measurements are presented along with surface measurements and photographs. Data tables explaining the measurement results are included.

M.J.S.

· N75-15105*# Northern Prairie Wildlife Research Center, Jamestown, N. Dak.

UTILIZATION OF SKYLAB (EREP) SYSTEM FOR APPRAISING CHANGES IN CONTINENTAL MIGRATORY BIRD HABITAT Monthly Progress Report, Dec. 1974

David S. Gilmer, Principal Investigator Dec. 1974 2 p EREP (NASA Order T-4114-8) (E75-10087; NASA-CR-141151) Avail: NTIS HC \$3.25 CSCL

060

01 ARGICULTURE AND FORESTRY

N75-15115*# Wyoming Univ., Laramie. Remote Sensing

USEFULNESS OF SKYLAB COLOR PHOTOGRAPHY AND ERTS-1 MULTISPECTRAL IMAGERY FOR MAPPING RANGE VEGETATION TYPES IN SOUTHWESTERN WYOMING Special Report

Robert C. Gordon, Principal Investigator [1974] 56 p refs ERTS and EREP

(Contracts NAS5-21799; NAS9-13298)

(E75-10097; NASA-CR-141293) Avail: NTIS HC \$4.25 CSCL 08B

The author has identified the following significant results. Aerial photography at scales of 1:43,400 and 1:104,500 was used to evaluate the usefulness of Skylab color photography (scales of 1:477,979 and 1:712,917) and ERTS-1 multispectral imagery (scale 1:1.000,000) for mapping range vegetation types. The project was successful in producing a range vegetation map of the 68,000 acres of salt desert shrub type in southwestern Wyoming. Techniques for estimation of above-ground green biomass have not yet been confirmed due to the mechanical failure of the photometer used in obtaining relative reflectance measurement. However, graphs of log transmittance versus above-ground green biomass indicate that production estimates may be made for some vegetation types from ERTS imagery. Other vegetation types not suitable for direct ERTS estimation of green biomass may possibly be related to those vegetation types whose production has been estimated from the multispectral imagery.

N75-15125*# Agricultural Research Service, Weslaco, Tex. IRRIGATION SCHEDULING, FREEZE WARNING AND SOIL SALINITY DETECTING Monthly Progress Report, Jan. 1975

Craig L. Wiegand, Principal Investigator Jan. 1975 4 p EREP (NASA Order T-4105-B) (E75-10109; NASA-CR-140899; MPR-13) Avail: NTIS HC \$3.25 CSCL 02C

N75-15146# Army Foreign Science and Technology Center, Charlottesville Va

USE OF FOREST VEGETATION-AS AN INDICATOR IN THE INTERPRETATION AND LARGE SCALE MAPPING OF SOILS IN A FOREST ZONE

A. M. Berezin 19 Nov. 1973 14 p refs Transl into ENGLISH from the book "Krupnomasshtabnoe Kartografiya Rastitelnosti" Novosibirsk, 1970 p 102-110

(AD-785967; FSTC-HT-23-1297-73) Avail: NTIS CSCL 08/13

Field research into forest types and their relations to soil types, types of tertiary deposits and underlying rocks were correlated with interpretation of the same items from aerial photographs, in selected regions of the Ukraine and Karelia. Aerial spectral region photographs proved to be more suitable than black and white aerial photos. The relationship of alluvial-slopewash deposits to ravine black alder, thick fluvioglacial sands covered with foxberry pine forests and organogenic deposits with rosemary-sphagnum pine forests are the most clearly interpreted from aerial photos. Further investigations in other regions should allow generalization of the results and their introduction into forest zone soil and tertiary deposit mapping.

02

ENVIRONMENTAL CHANGES AND CULTURAL RESOURCES

Includes land use analysis, urban and metropolitan studies, environmental impact, air and water pollution, geographic information systems, and geographic analysis.

A75-11103 # Morning-evening asymmetry of the thermosphere during a geomagnetic disturbance (Utrenne-vecherniaia asimmetriia termosfery v period geomagnitnogo vozmushcheniia). V. V. Katiushina and Iu. A. Romanovskii (Glavnoe Upravlenie Gidrometeorologicheskoi Sluzhby SSSR, Institut Prikladnoi Geofiziki, Moscow, USSR). Akademiia Nauk SSSR, Doklady, vol. 217, Aug. 11, 1974, p. 1057-1060. 8 refs. In Russian.

A75-11211 * The earth as a radio source - Terrestrial kilometric radiation. D. A. Gurnett (Iowa, University, Iowa City, Iowa). Journal of Geophysical Research, vol. 79, Oct. 1, 1974, p. 4227-4238. 17 refs. Contracts No. NAS5-11074; No. NAS5-11431; No. N00014-68-A-0196-0009; Grant No. NGL-16-001-043.

Based on Imp 6 and 8 satellite observation data, a comprehensive study of terrestrial kilometric radiation is presented. In the light of these data, the earth appears to be a very intense planetary radio source, with a total power output comparable to the decametric radio emission from Jupiter. Terrestrial kilometric (i.e., about 50-500 kHz) radiation seems to originate from low altitudes in the auroral region.

M.V.E.

A75-11221 * Magnetopause rotational forms. B. U. O. Sonnerup (Dartmouth College, Hanover, N.H.) and B. G. Ledley (NASA, Goddard Space Flight Center, Greenbelt, Md.). *Journal of Geophysical Research*, vol. 79, Oct. 1, 1974, p. 4309-4314. 17 refs. Grant No. NGR-30-001-040.

Magnetic 'field data from the Goddard Space Flight Center magnetometer experiment on board Ogo 5 are analyzed by the minimum-variance technique for two magnetopause crossings, believed to provide the best evidence presently available of magnetopause rotational discontinuities. Approximate agreement with predictions from MHD and first-order orbit theory is found, but available low-energy electron data suggest the presence of significant non-MHD effects. The paper also illustrates an improved method for data interval selection, a new magnetopause hodogram representation, and the utility of data simulation. (Author)

A75-12037 * # Skylab induced environment. R. J. Naumann (NASA, Marshall Space Flight Center, Huntsville, Ala.). American Institute of Aeronautics and Astronautics and American Geophysical Union, Conference on Scientific Experiments of Skylab, Huntsville, Ala., Oct. 30-Nov. 1, 1974, AIAA Paper 74-1225. 7 p. 10 refs.

This induced environment consists of particles and gases which emanate from the spacecraft in orbit and form the 'contamination cloud'. The presence of the cloud can present certain problems connected with scattering effects and the deposition of condensibles on optical surfaces. Measures taken to investigate these effects involved the placement of quartz crystal microbalance deposition monitors at certain locations to determine molecular film build-up during the flight.

G.R.

A75-12803 * On evaluating compliance with air pollution levels 'not to be exceeded more than once a year'. H. E. Neustadter and S. M. Sidik (NASA, Lewis Research Center, Energy Conversion and Materials Science Div., Cleveland, Ohio). Air Pollution Control Association, Journal, vol. 24, June 1974, p. 559-563. 11 refs.

Under a sampling program operated in conformity with EPA guidelines, calculational procedures are described for estimating the corresponding ambient pollution level. The statistical independence of the data, the commutativity of expectation and exponential operations, and the statistical variability of the calculated results are quantitatively evaluated. It is shown that the recommended sampling frequencies of every third to every sixth day are inadequate in determining conformity when the standards are expressed as levels not to be exceeded more than once a year.

T.S.

A75-13576 Technology today for tomorrow; Proceedings of the Eleventh Space Congress, Cocoa Beach, Fla., April 17-19, 1974. Volumes 1 & 2. Congress sponsored by the Canaveral Council of Technical Societies. Cape Canaveral, Fla., Canaveral Council of Technical Societies, 1974. Vol. 1, 199 p.; vol. 2, 177 p. Price of two volumes, \$21.25.

The topics discussed are related to an evaluation of Skylab payoffs, the benefits of space exploration, earth resources programs, the space shuttle, the utilization of foreign technology, and the marine sciences. Questions of planetary exploration are investigated, giving attention to the science of the Viking missions, Pioneer missions to Jupiter, the Mariner Jupiter/Saturn 1977 mission, and ballistic missions to the comet Encke in 1980. Subjects considered in the area of meteorology are related to an industry view of earth observation programs, technology for television meteorologists, rainfall estimations from geosynchronous satellite imagery, and an expansion of the meteorological satellite potential.

G.R.

A75-13577 * Skylab earth resources program. V. R. Wilmarth (NASA, Johnson Space Center, Houston, Tex.). In: Technology today for tomorrow; Proceedings of the Eleventh Space Congress, Cocoa Beach, Fla., April 17-19, 1974. Volume 1.

Cape Canaveral, Fla., Canaveral Council of Technical Societies, 1974, p. 1-1 to 1-13.

A unique space platform for studying the ocean, the land, and atmospheric phenomena of the planet earth was provided by the successful launch of Skylab on May 14, 1973. The crew members surveyed selected portions of this planet with an array of photographic, infrared, and microwave sensors. These sensors formed the Earth Resources Experiment Package (EREP). Approximately 40,000 photographs and 220,000 feet of high density magnetic tape were acquired with EREP. Details regarding the sensors are discussed along with the results obtained in the investigations.

G.R.

A75-13757 # Thermal emission of agglomerations as recorded on environmental satellite imagery. H. Kaminski (Bochum, Sternwarte, Bochum, West Germany). International Astronautical Federation, International Astronautical Congress, 25th, Amsterdam, Netherlands, Sept. 30-Oct. 5, 1974, Paper 74-139. 56 p. 29 refs.

The high-resolution IR radiometer of the satellites NOAA 2 and NOAA 3 makes it possible to determine the heat accumulation and emission in and from large towns. The film density of IR images obtained from the two satellites during the time from March 1973 and June 1974 was used to correlate the surface radiation temperature of the agglomeration and its environment for a number of cities in Europe, including London, Paris, Antwerp, Brussels, Duisburg, Frankfurt, Bremen, Hamburg, and Berlin.

G.R.

A75-13760 # Comparison of existing earth resources data processing facilities. G. P. Fishman (General Electric Co., Philadelphia, Pa.). International Astronautical Federation, International Astronautical Congress, 25th, Amsterdam, Netherlands, Sept. 30-Oct. 5, 1974, Paper 74-144. 16 p.

The paper provides readers potentially interested in establishing their own capability for accessing ERTS data, with an overview of the ground systems currently in operation in the U.S., Canada, and Brazil, and under active development in Italy and the Federal Republic of Germany. The reader is cautioned to look upon these facilities as examples only. There are many ways to implement ERTS ground systems ranging from small, relatively inexpensive options to large, complex, and costly implementations. Each implementation plan should be carefully analyzed to fit the needs and budget of the prospective user. F.R.L.

A75-14841 * # Nimbus 4 observations of changes in total ozone and stratospheric temperatures during a sudden warming. A. Ghazi (NASA, Goddard Space Flight Center, Greenbelt, Md.). Journal of the Atmospheric Sciences, vol. 31, Nov. 1974, p. 2197-2206. 14 refs.

A75-16675 * # The detection of dust storms over land and water with satellite visible and infrared measurements. W. E. Shenk and R. J. Curran (NASA, Goddard Space Flight Center, Greenbelt, Md.), Monthly Weather Review, vol. 102, Dec. 1974, p. 830-837, 8

The position of a large area of dust was successfully located over the Sahara Desert and off the west coast of Africa with infrared measurements from the Temperature Humidity Infrared Radiometer (THIR) and visible information from the Image Dissector Camera System (IDCS) on Nimbus 4 between 20 and 25 April 1970. Over land, near midday, the dust areas were associated with lower equivalent blackbody temperatures (TBB's) than in the surrounding areas due to the mixing of the dust with the cooler air just above the surface. Near midnight dust areas appeared to be located where the TBB's were higher than the surrounding regions. The TBB contrast was insufficient for dust detection over the ocean, but the contrast in the visible was great enough to track the movement of the dust for several days off the African west coast. Measurements from the THIR 6.5-7.2 micron channel were useful for detecting the presence of cirrus clouds which could have caused a misinterpretation of the 10.5-12.5 micron TBB's. (Author)

Ice particulates in the mesosphere. E. T. Chesworth and L. C. Hale (Pennsylvania State University, University Park, Pa.). Geophysical Research Letters, vol. 1, Dec. 1974, p. 347-350. 14 refs. Grants No. NGR-39-009-218; No. DA-ARO(D)-31-124-72-G158.

The observations of noctifucent clouds, the measurements of hydrated and of large immobile ions, and the light-scattering layer detected by the OGO-6 satellite suggest the presence of particulates, probably ice, in the mesosphere. The correlation between temperature and positive ion conductivity where the vapor pressure over ice becomes greater than atmospheric pressure in the stratopause region indicates the presence of ice crystals throughout the mesosphere over a wide range of latitudes during all seasons. Between one and ten billion ice crystals per cubic meter of order 10 nanometers in diameter can dominate ionization loss processes in the mesosphere, and can explain a variety of experimental observations, including observed solar angle dependence and seasonal variability of electron (Author) density.

A75-17102 # Some features of the urban environment of Tokyo by remote sensing. I. Tsuchiya (Meteorological Research Institute, Tokyo, Japan). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 3-12. 5 refs.

Observations of surface temperature of the urban area of Tokyo were performed making use of several airborne remote sensing

methods on bright sunny afternoons during summer. The surface temperature of a river with a vast amount of water has the lowest temperature, natural or near natural forests have the next lowest temperature, an urban area with many highways is hotter than a suburban area with greenery, and a main street with heavy traffic is the hottest. There are many cases requiring +5 to +10 C corrections to the airborne records on heated roads and buildings. Urban air circulation assumes very complicated regime due to compounded heated and cooled areas of various scale under the developed thermal inversion during the sunshiny days in summer. Such circulation causes a very complicated flow of polluted air resulting in several densely polluted spots in the suburban area. (Author)

A75-17105 * # Environmental studies of Iceland with ERTS-1. imagery. R. S. Williams, Jr. (U.S. Geological Survey, EROS Program Office, Reston, Va.), A. Boovarsson (Icelandic Surveying Department, Reykjavík, Iceland), S. Frioriksson, I. Thorsteinsson (Agricultural Research Institute, Reykjavik, Iceland), G. Palmason, S. Rist, K. Saemundsson (National Energy Authority, Iceland), H. Sigtryggsson (Icelandic Meteorological Service, Reykjavík, Iceland). and S. Thorarinsson (University of Iceland, Reykjavik, Iceland). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 31-81, 104 refs. Research supported

by the U.S. Geological Survey, NASA Order S-70243-G.

Imagery from the ERTS-1 satellite can be used to study geological and geophysical phenomena which are important in relation to Iceland's natural resources. Multispectral scanner (MSS) imagery can be used to map areas of altered ground, intense thermal emission, fallout from volcanic eruptions, lava flows, volcanic. geomorphology, erosion or build-up of land, snow cover, the areal extent of glaciers and ice caps, etc. At least five distinct vegetation types and barren areas can be mapped using MSS false-color composites. Stereoscopic coverage of iceland by side-lapping ERTS imagery permits precise analysis of various natural phenomena.

A.T.S.

A75-17112 * # A remote sensing study of Pacific hurricane Ava. D. Ross, J. McFadden (NOAA, Miami, Fla.), B. Au (U.S. Navy, Naval Research Laboratory, Washington, D.C.), and W. Brown (California Institute of Technology, Jet Propulsion Laboratory, Pasadena, Calif.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 163-180. 13

Aircraft, Skylab, NOAA-2, ATS-3, and NIMBUS-5 recently obtained a variety of measurements of Pacific hurricane Ava. These measurements are unusually broad in scope and include satellite observed passive microwave emissivities at 13.9 and 19.5 GHz, active microwave scattering cross-sections at 13.9 GHz, and near infrared and visible images. Essentially simultaneous aircraft measurements of wind speed, waves, whitecaps, 1.4 and 13-15 GHz passive microwave emissivities, 1.4 GHz active microwave images, sea surface temperatures, pressure fields, and aerosol size distributions were also obtained. A brief description of sensors and platforms is presented along with some in-depth details of results obtained. (Author)

A75-17114 * # Simultaneous active and passive microwave response of the earth - The Skylab radscat experiment. R. K. Moore, J. P. Claassen, A. C. Cook, D. L. Fayman, J. C. Holtzman, A. Sobti, F. T. Ulaby, J. D. Young (Kansas, University, Lawrence, Kan.), N. M. Hatcher (NASA, Johnson Space Center, Houston, Tex.), and W. E. Spencer, In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 189-217. 22 refs.

The Skylab microwave radiometer-scatterometer experiment S-193 was designed to collect simultaneous information on the radiometric emission and the backscatter from the earth from orbital altitude for the first time. A primary driving force was ascertaining the response from the oceans of both active and passive systems to variations in wind, and the ability of the passive system to aid in calibrating the active system. Numerous overland applications were also explored. Preliminary results of backscatter and emission from sea and atmosphere are shown for hurricane Ava and for two flights across the Gulf of Mexico. The results tend to confirm the most recent theoretical prediction of an approximately square-law relation between scattering coefficient for horizontal polarization and windspeed, with a somewhat smaller dependence for vertical polarization. Overland measurements show that scatterometer signals at 30 deg increase and radiometer signals decrease for terrain that has recently experienced rainfall. (Author)

A75-17129 # On the detectability of atmospheric carbon monoxide by microwave remote sensing. J. Fulde and E. Schanda (Bern, Universität, Berne, Switzerland). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 465-469. 5 refs.

The feasibility of tracing stratospheric and mesospheric carbon monoxide by microwave radiometric sensing of its 115-GHz line of the J sub 0-1 transition has been studied. A carefully designed balloon or airborne radiometer at 15 km altitude should be able to detect the excess temperature of 0.3 K due to CO at a zenith angle of 87 deg.

(Author)

A75-17130 • # Determination of the aerosol content in the atmosphere from ERTS-1 data. M. Griggs (Science Applications, Inc., La Jolla, Calif.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 471-481. 22 refs. Contract No. NAS5-21860.

The ERTS-1 satellite offered the opportunity of determining the feasibility of monitoring the atmospheric aerosol optical thickness on a global basis, as suggested by theoretical studies, which showed relationships between the upwelling earth-atmosphere radiance and the aerosol optical thickness, and between contrast and the aerosol optical thickness. These relationships were investigated at two test sites, using the multispectral scanner (MSS) radiance data, with ground-truth observations of the aerosol optical thickness being made with a Volz photometer at the time of the satellite overpasses. The results indicate that the MSS channels, 4, 5, and 6, centered at 0.55, 0.65, and 0.75 micron have comparable sensitivity, and that the optical thickness can be determined to within plus or minus 10% with the assumed measurement errors of the MSS. (Author)

A75-17131 • # The effect of atmospheric water vapor on automatic classification of ERTS data. D. E. Pitts, W. E. McAllum (NASA, Johnson Space Center, Houston, Tex.), and A. E. Dillinger (Lockheed Electronics Co., Inc., Houston, Tex.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 483-497. 9 refs.

Absorption by atmospheric water vapor changes the spectral signatures collected by multispectral scanners if channels are not chosen to avoid the atmospheric water bands. For ERTS (Earth Resources Technology Satellite), the Multispectral Scanner band 7 (MSS 7, .8 to 1.1 micron) is the only band significantly affected. Line-by-line atmospheric absorption calculations showed that this

effect can multiply the intensity by factors ranging from .77 to 1.0. If horizontal gradients in atmospheric water exist between training fields and the rest of the scene, errors are introduced in automatic classification of the imagery. The degradation of the classification of corn and soybeans was determined by using actual ERTS data and simulating the absorption effects on the MSS 7 band. (Author)

A75-17133 # Design concepts for land use and natural resource inventories and information systems. R. L. Shelton (Michigan State University, Ann Arbor, Mich.) and E. E. Hardy (Cornell University, Ithaca, N.Y.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 517-535, 5 refs.

A75-17136 # Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis. V. A. Milazzo (U.S. Geological Survey, Reston, Va.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 569-585. 8 refs.

Work undertaken on a three-sensor land use data evaluation for a portion of the Phoenix area is reported. Analyses between land use data generated from 1970 high-altitude photography and that detectable from ERTS and Skylab, especially in terms of changes in land use indicate that ERTS and Skylab imagery can be used effectively to detect and identify areas of post-1970 land use change, especially those documenting urban expansion at the rural-urban fringe. Significant preliminary findings on the utility of ERTS and Skylab data for metropolitan land use analysis, substantiated by evaluations with 1970 and 1972 ground control land use data are reported. (Author)

A75-17137 * # Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado. J. E. Nichol (Aston, University, Birmingham, England). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 589-596. 8 refs. Research supported by the Boulder Area Growth Study Commission and Boulder County Department of Planning; Grant No. NGL-06-003-200.

A75-17149 # Interactive machine assessment of critical land resources using ERTS-1 data. W. W. Kuhlow and L. T. Fisher (Wisconsin, University, Madison, Wis.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 759-769. NSF-supported research.

The University of Wisconsin's Space Science and Engineering Center is currently developing a highly interactive man-machine system called McIDAS (Man-Computer Interactive Data Access System) created primarily for quick access to large volumes of digital data generated by satellites such as ATS-III, SMS and ERTS-1. The system is presently operable and several research projects are being carried out as further system developments proceed. This paper will describe the McIDAS system and some of its features which allow superior man-machine interaction. Impressions and results of investigators experienced in conventional remote sensing data analysis but completely unfamiliar with this system or other digital systems are presented. (Author)

02' ENVIRONMENTAL CHANGES AND CULTURAL RESOURCES

A75-17153 * # Implementation of an advanced table look-up classifier for large area land-use classification. C. Jones (NASA, Johnson Space Center, Earth Resources Laboratory, Bay St. Louis, Miss.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 813-824. 6 refs.

Software employing Eppler's improved table look-up approach to pattern recognition has been developed, and results from this software are presented. The look-up table for each class is a computer representation of a hyperellipsoid in four dimensional space. During implementation of the software Eppler's look-up procedure was modified to include multiple ranges in order to accommodate hollow regions in the ellipsoids. In a typical ERTS classification run less than 6000 36-bit computer words were required to store tables for 24 classes. Classification results from the improved table look-up are identical with those produced by the conventional method, i.e., by calculation of the maximum likelihood decision rule at the moment of classification. With the new look-up approach an entire ERTS MSS frame can be classified into 24 classes in 1.3 hours, compared to 22.5 hours required by the conventional method. The new software is coded completely in FORTRAN to facilitate transfer to other digital computers.

A75-17157 * # Effect of atmospheric haze and sun angle on automatic classification of ERTS-1 data. J. Potter and M. Shelton (Lockheed Electronics Co., Inc., Houston, Tex.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 865-874. 7 refs. Contract No. NAS9-12200.

The effect of variations in sun angle and haze level on the accuracy of automatic classification of Earth Resources Technology Satellite-1 (ERTS-1) data was studied by classifying ERTS imagery in which such variations were computer-simulated. It was found that relatively small changes in sun angle and haze level can substantially reduce classification accuracy. (Author)

A75-17162 * # Scanning thermal plumes. F. L. Scarpace, R. P. Madding, and T. Green, III (Wisconsin, University, Madison, Wis.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 939-961. Research supported by the Wisconsin Department of Natural Resources, Wisconsin Power and Light Co., Wisconsin Public Service Corp., Wisconsin Electric Power Co., U.S. Department of Commerce, and NASA.

In order to study the behavior and effects of thermal plumes associated with the condenser cooling of power plants, thermal line scans are periodically made from aircraft over all power plants along the Wisconsin shore of Lake Michigan. Simultaneous ground truth is also gathered with a radiometer. Some sequential imagery has been obtained for periods up to two hours to study short term variations in the surface temperature of the plume. The article concentrates on the techniques used to analyze thermal scanner data for a single power plant which was studied intensively. The calibration methods, temperature dependence of the thermal scanner, and calculation of the modulation transfer function for the scanner are treated. It is concluded that obtaining quantitative surface-temperature data from thermal scanning is a nontrivial task. Accuracies up to plus or minus 0.1 C are attainable.

A75-17165 # Correlation of multispectral imagery with water analysis - The Ross Barnett Reservoir Remote Sensing Project. D. L. Wertz, W. T. Mealor, M. L. Steele, and J. W. Pinson (Southern Mississippi, University, Hattiesburg, Miss.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 995-999.

A75-17182 * # Boundaries of ERTS and aircraft data within which useful water quality information can be obtained. W. G. Egan (Grumman Aerospace Corp., Bethpage, N.Y.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1319-1343. 5 refs. NASA-sponsored research. Calibration procedures have been devised and applied to ERTS-1, multispectral, true color, and false color imagery. The results indicate that the ERTS and multispectral imagery are correlated with optical in situ measurements of the harbor water. Correlation is extended to true and false color imagery through in situ optical measurements of the harbor water. The best photometric accuracy is achieved with multispectral aerial imagery and the use of bulk MSS tape. The aircraft green photographic and ERTS-1 MSS-4 bands have been found most suitable for monitoring the scattered light levels under the conditions of this investigation. The application of satellite or aircraft for optical remote sensing depends upon the physical scale and frequency of sensing since both sensor systems generally have sufficient photometric sensitivity. The chemical parameters of the harbor water were found to be correlated to the optical properties for two stations investigated in detail.

A75-17188 * # Land use classification and ground truth correlations from simultaneously acquired aircraft and ERTS-1 MSS data. A. J. Richardson, M. R. Gautreaux, R. J. Torline, and C. L. Wiegand (U.S. Department of Agriculture, Weslaco, Tex.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1423-1440. 12 refs. NASA Order S-70251-AG.

An analytical formula is developed which describes the relationships between the sporadic cumulus coverage, the sun height, and the additional coverage of the earth's surface by cloud shadows. Based on this formula, the sun height of about 40 to 45 deg is optimized for natural resources survey from satellites synchronous with the sun. (Author)

A75-17205 * # Remote measurements of water pollution with a lidar polarimeter. T. C. Sheives, J. W. Rouse, Jr., and W. T. Mayo, Jr. (Texas A & M University, College Station, Tex.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1695-1708. 7 refs. U.S. Department of Transportation Contract No. CG-34071-A; Grant No. NGL-44-001-001.

This paper examines a dual polarization laser backscatter system as a method for remote measurements of certain water quality parameters. Analytical models for describing the backscatter from turbid water and oil on turbid water are presented and compared with experimental data. Laser backscatter field measurements from natural waterways are presented and compared with simultaneous ground observations of the water quality parameters: turbidity, suspended solids, and transmittance. The results of this study show that the analytical models appear valid and that the sensor investigated is applicable to remote measurements of these water quality parameters and oil spills on water.

(Author)

Multidisciplinary applications of ERTS and A75-17234 * # Skylab data in Ohio, D. C. Sweet, P. G. Pincura (Department of Economic and Community Development, Columbus, Ohio), C. J. Meier (Department of Natural Resources, Columbus, Ohio), G. B. Garrett (Ohio Environmental Protection Agency, Columbus, Ohio), L. O. Herd (U.S. Department of Transportation, Washington, D.C.), J. M. Dowdy (Ohio State University, Columbus, Ohio), D. M. Anderson (Ohio Biological Survey, Columbus, Ohio), G. E. Wukelic, J. G. Stephan, and H. E. Smail (Battelle Columbus Laboratories, Columbus, Ohio). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Ann Arbor, Mich., Proceedings, Volume 3. Environmental Research Institute of Michigan, 1974, p. 2093-2106. 8 refs. NASA-supported research.

Experimental studies of ERTS-1 and Skylab earth resources data, in combination with correlative aircraft and on-site data, for environmental quality, land use, and resource management applications in Ohio show several areas of operational promise. Prime data use candidates demonstrated to date include definition and enforcement of surface mining (all minerals) legislation; Lake Erie modeling/management; land use classification and mapping studies at state, regional, and localized levels; and resources' inventories particularly of forested areas on both regional (multicounty) and localized scales.

(Author)

A75-17667 Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV. E. P. Mazets, S. V. Golenetskii, V. N. Il'inskii, Iu. A. Gur'ian, and T. V. Kharitonova (Akademiia Nauk SSSR, Fiziko-Tekhnicheskii Institut, Leningrad, USSR). (ZHETF Pis'ma v Redaktsiiu, vol. 20, July 20, 1974, p. 77-80.) JETP Letters, vol. 20, July 20, 1974, p. 32-34. 14 refs. Translation.

The Kosmos-461 satellite carried experiments to study the diffuse background of X rays and gamma rays in the range 28 keV-4.1 MeV. The procedures developed for eliminating the influence of atmospheric radiation and of radioactivity induced in the detector by cosmic rays are described. It is concluded that the diffuse background can not be represented by a single law in the range investigated. Measurements by Apollo 15 and high-altitude balloons are judged to overestimate the gamma-ray flux in the 10-30 MeV range.

A.T.S.

A75-17932 # The potentialities for determining content of minor gas constituents in the atmosphere by satellites (O vozmozhnostiakh opredeleniia soderzhaniia malykh gazovykh komponentow atmosfery so sputnikov). V. V. Rozanov and Iu. M. Timofeev (Leningradskii Gosudarstvennyi Universitet, Leningrad, USSR). Meteorologiia i Gidrologiia, Nov. 1974, p. 87-91. 23 refs. In Russian.

The use of oblique measurements of atmospheric transmittance as a base for determinations of the concentrations of minor gas constituents (CO, CH4, NO, N2O, NH3, etc.) in the atmosphere is briefly discussed. A simple, approximate formula, not requiring any knowledge of the fine structure of absorption bands, is given for computing vertical profiles of gas concentrations from transmittance data.

P.T.H.

A75-18054 Recent progress in urban mapping. J. P. Burns (U.S. Geological Survey, National Center, Reston, Va.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings. Falls Church, Va., American Society of Photogrammetry, 1974, p. 45-49.

Accurate large-scale maps of urban areas are urgently needed. They are the first step in a logical solution to severe socioeconomic problems caused by population explosion in these regions. Fragmented urban governmental structure makes urban mapping uncoordinated, repetitive and wasteful unless it is undertaken by a centrally-coordinated Federal effort. Several pilot projects have provided knowledge of specific urban map needs on which a national map series can be based. An early 1973 program decided upon a 1:2400 scale in order to reach a broad user cross-section and orthophoto mapping for detail and ease in updating. At Fort Wayne,

Ind., 440 half-square-mile orthophoto map bases were produced, and the Schut method was applied in aerotriangulation. USGS-monitored non-federal projects are also under way.

S.J.M.

A75-18057 Remote sensing and land use inventory and mapping. J. R. Anderson (U.S. Geological Survey, Reston, Va.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings. Falls Church, Va., American Society of Photogrammetry, 1974, p. 66-71. 7 refs.

In the Geography Program of the U.S. Geological Survey, we have been working experimentally for more than two years to develop a program which will provide needed data on current land use efficiently and in a timely way. A two-level resource-oriented land use classification system for use with remote sensor data has been developed. This system can be expanded by State, local governments, or by Federal agencies to the third and fourth levels for greater detail. The classification system and new methods of collecting current land use data using remote sensor source data have been successfully tested. (Author)

A75-18058 The application of ERTS-derived information to the regional land use planning process. J. L. Clapp, R. W. Kiefer, W. W. Kuhlow, and B. J. Niemann, Jr. (Wisconsin, University, Madison, Wis.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings. Falls Church, Va., American Society of Photogrammetry, 1974, p. 72-87.

The need for the development and implementation of methods for the detection, inventory and monitoring of land resource variables is reflected in present and pending federal and state legislation. Data has been extracted from ERTS imagery, RB-57 color infrared imagery and best available conventional sources for a 10,000 square kilometer test area in eastern Wisconsin. The data is compared on a spatial basis for a 300 square kilometer portion of the test area and the effect of the data source on land use decisions is examined. Based upon these investigations it is concluded that ERTS can provide an operational data source for land use planning at the policy level. (Author)

A75-18059 * Land use classification with ERTS-1 data in the Houston area test site. J. E. Dornbach (NASA, Johnson Space Center, Houston, Tex.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings. Falls Church, Va., American Society of Photogrammetry, 1974, p. 88-98.

A75-18061 Remote sensing for regional resource analysis.

O. W. Mintzer and J. M. Dowdy (Ohio State University, Columbus, Ohio). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings.

Falls Church, Va., American Society of Photogrammetry, 1974, p. 137-154. 12 refs.

In improving levels of living, we become increasingly involved in activities which provide additional economic growth. These activities include the development of natural resources to make them more accessible for productive and beneficial uses. In order to accomplish these goals we are turning to remote sensing technology which is useful for many data acquisition tasks. The technique of using ERTS imagery, airphotos and mini-format aerial color photography is demonstrated as the synergistic remote sensing approach for inventories of regional resources. (Author)

A75-18064 The application of ERTS-1 data to the Corps of Engineers Civil Works Mission. S. C. Sollers (U.S. Army, Office of the Chief of Engineers, Washington, D.C.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings. Church, Va., American Society of Photogrammetry, 1974, p. 181-203.

02 ENVIRONMENTAL CHANGES AND CULTURAL RESOURCES

In support of its civil works function, the Army Corps of Engineers is constantly engaged in applied research to observe natural processes and ecological relationships in the interest of preservation of natural resources. Remotely sensed data facilitates this research. The application of remotely sensed imagery results in the saving of time and money and increased precision in gross evaluation of large ecosystems and provides a convenient method to monitor changes within such a system. Several projects are now underway employing remotely sensed data acquired from the NASA Earth Resources Technology Satellite (ERTS-1). (Author)

A75-18069 Application of ERTS data to environmental inventory and assessment. J. D. O'Neal and J. R. Burns (U.S. Army, Engineer Topographic Laboratories, Fort Belvoir, Va.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings. Falls Church, Va., American Society of Photogrammetry, 1974, p. 256-261.

The use of ERTS imagery to supplement aerial photography and collateral source material has expedited the preparation of land use and vegetative cover maps included in environmental resources inventory studies. The studies are being prepared by the Engineer Agency for Resources Inventories (EARI), an element of the U.S. Army Engineer Topographic Laboratories, to satisfy civil works program requirements of Corps of Engineers division and district offices for basic input for project planning and preparation of environmental impact statements. The paper describes recent experience gained by EARI in support of programs of district offices of the Lower Mississippi Valley Division and the Ohio River Division of the Corps. (Author)

A75-18070 Land use mapping using ERTS-1 imagery for The Four Rivers Resource Conservation and Development District in Kansas. R. V. Shaklee and T. L. Talmon (Kansas, University, Lawrence, Kan.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings. Falls Church, Va., American Society of Photogrammetry, 1974, p. 262-267.

A75-18108 # New methods for studying the dynamics of the entire earth. E. Tengstrom (Uppsala, Universitet, Uppsala, Sweden). In: Geodesy and physics of the earth; Proceedings of the Second International Symposium, Potsdam, East Germany, May 7-11, 1973. Part 1. Potsdam, East Germany, Deutsche Akademie der Wissenschaften, Zentralinstitut für Physik der Erde, 1974, p. 39-46.

Methods involving new theoretical procedures including the collaboration of scientists in geodesy, geophysics, and astronomy in a unified approach are required to study geodynamical problems related to a deformable earth with elastoviscous properties and discontinuities of density, elasticity, and plasticity. Attention is given to astrogeodetic measurements, gravity field measurements, earth tide records, and aspects of very-long base-line interferometry. G.R.

A75-18151 Conference on Weather Forecasting and Analysis, 5th, St. Louis, Mo., March 4-7, 1974, Preprints. Conference sponsored by the American Meteorological Society. Boston, American Meteorological Society, 1974. 301 p. Members, \$15.; nonmembers, \$20.

Topics discussed include automated weather systems of the future; the formulation, expression, dissemination, and use of weather forecasts; the use of satellite data in weather forecasting; observations of precipitation and clouds; mesoscale and synoptic-scale forecasting and analysis; air pollution and urban effects; applications and deficiencies of synoptic meteorological material; ceiling and visibility forecasting; planetary boundary layer phenomena; and convective activity.

A.B.K.

A75-18153 # Observing and forecasting local effects from satellite data. F. C. Parmenter (NOAA, National Environmental Satellite Service, Washington, D.C.). In: Conference on Weather Forecasting and Analysis, 5th, St. Louis, Mo., March 4-7, 1974, Preprints.

Boston, American Meteorological Society, 1974, p. 46-49.

Many mesoscale cloud patterns are strongly influenced by terrain, geography and urbanization. Continuous satellite data over the U.S. have allowed meteorologists to observe the establishment and dissipation of many small-scale cloud formations due to local effects heretofore known mainly to residents of the locale. The distribution of cloudiness to the lee of the Great Lakes, along the East Coast, and over the mountainous states is discussed. (Author)

A75-19457 • A Shuttle/SpaceLab RF environment survey facility. R. E. Taylor (NASA, Goddard Space Flight Center, Greenbelt, Md.), R. E. Prince, and D. N. McGregor (Operations Research, Inc., Silver Spring, Md.). In: National Telecommunications Conference, San Diego, Calif., December 2-4, 1974, Record.

New York, Institute of Electrical and Electronics Engineers, Inc., 1974, p. 318-326.

This paper describes some of the basic system considerations of a proposed experiment for monitoring and measuring the global electromagnetic radiation environment. RF spectrum coverage for this spaceborne survey facility is anticipated to be quite broad and will include selected frequency bands ranging from UHF to millimeter waves. By establishing this RF environment survey facility, a broad data base of useful electromagnetic field intensity information will be developed. This, in turn, should help both the regulatory agencies and user community in terms of future frequency planning and system design by avoiding unwarranted interference. It is proposed that this experiment be flown as a future test bed during the Shuttle/Spacelab era for continuously gathering and updating information on earth-emitted electromagnetic emissions on a global scale. (Author)

N75-10469# Sandia Corp., Albuquerque, N.Mex.
MODERN TRENDS IN ENVIRONMENTAL TESTING

A. J. Clark, Jr. 1 Feb. 1974 16 p refs Presented at the 20th Ann. Tech. Meeting and Equipment Exposition of the Inst. of Environ. Sci., Washington, D. C., 28 Apr. 1974 Sponsored by AEC

(SLA-73-5931; Conf-740415-1) Avail: NTIS HC \$3.25

From 20th annual technical meeting and equipment exposition of the Institute of Environmental Sciences; Washington, District of Columbia, U.S.A. (28 Apr. 1974).

NSA

N75-10535*# Boeing Co., Kent, Wash.
QUANTITATIVE DETERMINATION OF STRATOSPHERIC
AEROSOL CHARACTERISTICS Monthly Report, Sep. 1974
David L. Tingey, Principal Investigator Sep. 1974 1 p EREP
(Contract NAS9-13303)
(E74-10808; NASA-CR-140502) Avail: NTIS HC \$3.25 CSCL

N75-10538*# Tennessee Univ., Knoxville. Dept. of Electrical Engineering.

ERTS-1 IMAGERY INTERPRETATION TECHNIQUES IN THE TENNESSEE VALLEY Final Report, Sep. 1972 - Mar. 1974
Robert E. Bodenheimer, Principal Investigator Apr. 1974 59 p refs Original contains imagery. Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue. Sioux Falls, S. D. 57198 ERTS

(Contract NAS5-21875)

(E74-10811; NASA-CR-140505; TR-EE/CS-74-5) Avail: NTIS HC \$4.25 CSCL 08B

The author has identified the following significant results. The feasibility of delineating major soil associations and land

uses through computerized analyses is discussed. Useful and potential applications in detecting landscape change and land use mapping are described. Recommendations for improving the data processing effort in a multidisciplinary program are presented.

N75-10545*# Dartmouth Coll., Hanover, N.H. Dept. of Geography

INVESTIGATION OF LAND USE OF NORTHERN MEGALOP-OLIS USING ERTS-1 IMAGERY Final Report, Jun. 1972 -Jul. 1973

Robert B. Simpson, David T. Lindgren, David J. Ruml, and William Goldstein Washington NASA Sep. 1974 63 p refs Original contains color illustrations

(Contract NAS5-21749)

(NASA-CR-2459; S13-116) Avail: NTIS HC \$4.25 CSCL O8B

Primary objective was to produce a color-coded land use map and digital data base for the northern third of Megalopolis. Secondary objective was to investigate possible applications of ERTS products to land use planning. Many of the materials in this report already have received national, dissemination as a result of unexpected interest in land use surveys from ERTS. Of special historical interest is the first comprehensive urban-type land use map from space imagery, which covered the entire state of Rhode Island and was made from a single image taken on 28 July 1972.

N75-10551* Alaska Univ., Fairbanks. ENVIRONMENTAL SURVEYS IN ALASKA BASED UPON ERTS DATA

John M. Miller In NASA. Goddard Space Flight Center Third ERTS Symp., Vol. 2 May 1974 p 12-40

CSCL 08B

Alaskan applications of ERTS data are summarized. Areas discussed are: (1) land use; (2) archaeology; (3) vegetation mapping; (4) ice reporting and mapping; (5) fermafrost; (6) mineral and oil exploration; (7) geological surveys; (8) seismology; (9) geological faults and structures; (10) hydrology and water resources; (11) glaciology; (12) water circulation in Cook Inlet; and (13) fish and mammal populations.

A.L.

N75-10560* National Aeronautics and Space Administration. Mississippi Test Facility, Bay Saint Louis.

LAND USE ANP MAPPING

Armond T. Joyce *In its* Third ERTS Symp., Vol. 2 May 1974 p 138-146 CSCL 08B

The ERTS program provides data that can be used to derive information relative to the actual use of the land resource, in a practical and timely manner. ERTS data provide coverage of total land areas, and its repetitive nature enables the detection and monitoring of changes taking place in land use. Generally, the techniques and the procedures used to extract information from ERTS data may be categorized as pertaining to either the interpretations of ERTS imagery or to the use of digital data and computer techniques. Examples are given of the use of ERTS-1 data for land use classification in: (1) New England areas: (2) Chesapeake Bay and Washington, D.C.; (3) Mississippi Gulf Coast; (4) Los Angeles, California; (5) Houston, Texas; and (6) Phoenix, Arizona.

N75-10563*# Texas A&M Univ., College Station. Remote Sensing Center.

A STUDY OF A DUAL POLARIZATION LASER BACKSCATTER SYSTEM FOR REMOTE IDENTIFICATION AND MEASUREMENT OF WATER POLLUTION

Thomas Carlyle Sheives May 1974 148 p refs (Grant NGL-44-001-001; Contract DOT-CG-34017-A) (NASA-CR-140665; RSC-53) Avail: NTIS HC \$5.75 CSCL 171 Remote identification and measurement of subsurface water turbidity and oil on water was accomplished with analytical models which describe the backscatter from smooth surface turbid water, including single scatter and multiple scatter effects. Lidar measurements from natural waterways are also presented and compared with ground observations of several physical water quality parameters.

N75-10572*# Miami Univ., Coral Gables, Fla. Clean Energy Research Inst.

APPLICATION OF REMOTE SENSING FOR PREDICTION AND DETECTION OF THERMAL POLLUTION

T. Nejat Veziroglu and Samuel S. Lee Oct. 1974 100 p refs (Contract NAS10-8498)

(NASA-CR-139182) Avail: NTIS HC \$4.75 CSCL 13B

The first phase is described of a three year project for the development of a mathematical model for predicting thermal pollution by use of remote sensing measurements. A rigid-lid model was developed, and results were obtained for different wind conditions at Biscayne Bay in South Florida. The design of the measurement system was completed, and instruments needed for the first stage of experiment were acquired, tested, and calibrated. A preliminary research flight was conducted.

Author

N75-10638# California Univ., Livermore. Lawrence Livermore

COMPARISON OF THE SOUTHERN AND NORTHERN HEMISPHERE GENERAL CIRCULATION CHARACTERISTICS AS DETERMINED BY SATELLITE OZONE DATA

J. E. Lovill Mar. 1974 51 p refs Presented at the 1st Spec. Assembly of the Intern. Assoc. of Meteorol. and Atmospheric Phys., Melbourne, 14-25 Jan. 1974 Sponsored by AEC (UCRL-75556; Conf-740106-4) Avail: NTIS HC \$4.25

Satellite total ozone data were analyzed for the Southern Hemisphere (autumn and winter) and the Northern Hemisphere (spring and summer). A strong correlation was found between global meridional gradients of total ozone and the wand velocity in baroclinic zones. The ozone wave maxima correspond to regions of stratospheric heating. A large region of low total ozone was observed in the four months averages in equatorial regions, extending from Southeast Asia to the central Pacific; a large area of high equatorial ozone occurs from the central Atlantic across Africa. This ozone distribution suggests a large double cell motion near the equator, such that there is a large scale descending motion in the Atlantic and African sectors, as well as a slowly ascending motion in the Southeast Asian and Pacific sectors. A relatively tight ozone gradient was also seen in the Atlantic Ocean between 40 deg N and 50 deg N, a region which corresponds to the high density aircraft flight corridor between the eastern United States and the European region.

Author

N75-11413*# National Aeronautics and Space Administration.
Goddard Space Flight Center, Greenbelt, Md.

EXPLORATION FOR FOSSIL AND NUCLEAR FUELS FROM ORBITAL ALTITUDES

Nicholas M. Short Nov. 1974 59 p

(NASA-TM-X-70781; X-923-74-322) Avail: NTIS HC \$4.25 CSCL 08G

Results from the ERTS program pertinent to exploration for oil, gas, and uranium are discussed. A review of achievements in relevant geological studies from ERTS, and a survey of accomplishments oriented towards exploration for energy sources are presented along with an evaluation of the prospects and limitations of the space platform approach to fuel exploration, and an examination of continuing programs designed to prove out the use of ERTS and other space system in exploring for fuel resources.

N75-11476# World Meteorological Organization, Geneva (Switzerland).

OBSERVATION AND MEASUREMENT OF ATMOSPHERIC POLLUTION

1974 675 p refs Partly in ENGLISH, FRENCH and SPANISH Proc. of the TECOMAP Conf., Helsinki, 30 Jul. - 4 Aug. 1973; sponsored in part by the World Health Organ. (WMO-368; Rept-3; ISBN-92-63-10368-2) Avail: NTIS HC

(WMO-368; Rept-3; ISBN-92-63-10368-2) Avail: NTIS H \$15.25

Papers are presented on: requirements for measurement of air pollutants for forecasting of pollution potential; measurement and analysis of pollutants on urban, regional, and global scales; turbidity and radiation measurement as indicators of pollution; and method of standardization and intercomparison.

N75-11498 World Meteorological Organization, Geneva (Switzerland).

A TECHNIQUE FOR MEASUREMENT OF SULPHUR DIOXIDE AND SULPHATE CONCENTRATIONS FROM AN AIRCRAFT

D. A. Johnson and D. H. F. Atkins *In its* Observation and Meas. of Atmospheric Pollution 1974 p 216-222 refs

An airborne air sampling is described which, using a dry filtration technique, enables sulfur dioxide and sulfate concentration to be determined over distances of 40 km or more. ESRO

N75-11505 World Meteorological Organization, Geneva (Switzerland)

CORRELATION SPECTROSCOPY TECHNIQUES FOR THE REGIONAL AND GLOBAL MEASUREMENTS OF AIR POLLUTION

J. H. Davies In its Observation and Meas. of Atmospheric Pollution 1974 p 282-296 refs

The remote sensing approach to air quality monitoring is reviewed, and optical correlation techniques applied to remote sensing of gaseous atmospheric constituents are discussed. Correlation spectroscopy is described, and results of studies using the method for aircraft and balloon measurements of sulfur and nitrogen dioxide are presented. The application of airborne and satellite mounted remote sensors to atmospheric monitoring is also discussed.

N75-11510 World Meteorological Organization, Geneva (Switzerland).

GEOPHYSICAL MONITORING FOR CLIMATIC CHANGE. THE NOAA PROGRAM

D. H. Pack, R. Fegley, G. Herbert, D. Hoyt, W. Komhyr, J. Miller, and C. Turner *In its* Observation and Meas. of Atmospheric Pollution 1974 p 334-345 refs

The overall objective of the NOAA's program is to predict the effects of natural or artificial emissions into the atmosphere on climate. The processes by which the monitoring program was designed, locations for additional observations selected, the rationale for the primary measurement program and a brief discussion of the intrumentation, the quality control and data aspects required for the program are outlined.

 ${\bf N75-11521}^{f *}$ World Meteorological Organization, Geneva (Switzerland).

GLOBAL MEASUREMENTS OF AIR POLLUTION FROM SATELLITES

L. L. Acton, E. R. Bartle, M. Griggs, G. D. Hall, W. D. Hesketh, C. B. Ludwig, W. Malkmus, and H. Reichle In its Observation and Meas. of Atmospheric Pollution 1974 p 434-442 refs

(Contract NAS1-12048)

The conceptual design of an FOV nadir radiometer was examined for its applicability to monitoring the radiation process in the atmosphere as it relates to aerosol behavior. The instrument employs a gas filter correlation technique and is suitable for transportation onboard satellite.

N75-11523 World Meteorological Organization, Geneva (Switzerland).

MEASUREMENT OF TRACE STRATOSPHERIC GASES BY INVERTING THERMAL INFRARED LIMB RADIANCE PROFILES

John C. Gille, James M. Russell, III, Paul L. Bailey, and Frederick B. House *In its* Observation and Meas. of Atmospheric Pollution 1974 p 455-462 refs

An infrared technique for the remote measurement by satellite of minor constituents in the upper troposphere, stratosphere, and lower mesosphere is described. The spectral features used are explained, and the quality of results expected are discussed. A summary of expected capabilities from such an experiment is presented, with consideration of instrumentation requirements.

N75-11527 World Meteorological Organization, Geneva (Switzerland).

GROUND-BASED MEASUREMENTS OF SOLAR RADIA-TION BY GEOPHYSICAL MONITORING FOR CLIMATIC CHANGE (USA)

Douglas V. Hoyt and Gary A. Herbert In its Observation and Meas. of Atmospheric Pollution 1974 p 506-515 refs (For

A progress report is presented on the geophysical monitoring for climatic change (GMCC) program which is to provide baseline observations of the incident solar radiation. The field and calibration instrumentation (multichannel pyroheliometers, pyranometers, and radiometers) are described in detail, and aspects of the data analysis are discussed.

N75-11529* World Meteorological Organization, Geneva (Switzerland).

MICROWAVE ROTATIONAL SPECTROSCOPY: A PHYSICAL TECHNIQUE FOR SPECIFIC POLLUTANT MONITORING

Lawrence W. Hrubesh In its Observation and Meas. of Atmospheric Pollution 1974 p 526-537 refs

(NASA Order L-75902; Contracts EPA-IAG-D179(D))

Evidence is presented that microwave rotational spectroscopy can be developed for use in air pollution monitoring, with the employment of a diode-cavity spectrometer capable of detecting small concentrations of NO2, SO2, H2CO, and HN3 gas with _very.high.specificity.—The-theory-behind the technique is discussed together with its application and experimental results. ESRO

N75-11531 World Meteorological Organization, Geneva (Switzerland).

MEASUREMENTS OF METEOROLOGICAL PARAMETERS AND AIR POLLUTION SPREAD OVER STOCKHOLM, SWEDEN, BY MEANS OF REMOTE SENSING AND AIRBORNE INSTRUMENTATION

Lars E. Olsson, Christer Persson, Arne Haagaard, Hans Ottersten, Lennart Granat, and Claes Traegaardh *In its* Observation and Meas. of Atmospheric Pollution 1974 p 545-556 refs Sponsored partly by the Swed. Environ. Protect. Board and the Swed. Board for Tech. Develop.

A comparison is made of different available measurement techniques suitable for studies of mesoscale air pollution dispersion over Stockholm, Sweden. The measurement program is outlined and the aircraft instrumentation briefly described. The results are presented and discussed.

N75-11540 World Meteorological Organization, Geneva (Switzerland).

QUALITY CONTROL IN GLOBAL AND REGIONAL AIR CHEMISTRY MONITORING NETWORKS: LESSONS OF THE EUROPEAN AIR CHEMISTRY NETWORK

M. P. Paterson and R. S. Scorer In its Observation and Meas. of Atmospheric Pollution 1974 p 647-655 refs

A preliminary analysis was made of annual data on the Ca/Mg ratio from the continental area of the European Air

Chemistry Network. The results of this analysis are given and it is concluded that much of the data is wrong, due to faults in the chemical analysis.

N75-11552* Tohoku Univ., Sendai (Japan). RADIATIVE TRANSFER OF VISIBLE RADIATION IN TURBID ATMOSPHERE

Giichi Yamamoto and Masayuki Tanaka In Calif. Univ. Proc. of the UCLA Intern. Conf. on Radiation and Remote Probing of the Atmosphere 1974 p 74-134 refs

CSCL 04A

Methods are presented for solving radiative transfer problems; they include the doubling method and the closely related matrix method, iterative method, Chandrasekhar's method of discrete ordinates, and Monte Carlo method. To consider radiation transport through turbid atmosphere, an atmospheric model was developed characterizing aerosols by parameters. Intensity and polarization of radiation in turbid atmospheres is discussed, as well as lower atmospheric heating due to solar radiation absorption by aerosols.

J.A.M.

N76-11553* Mainz Univ. (West Germany). SCATTERING AND ABSORPTION FROM POLY-DISPERSED AEROSOLS

K. Bullrich, R. Eiden, G. Eschelbach, K. Fischer, G. Haenel, and J. Heintzenberg In Calif. Univ. Proc. of the UCLA Intern. Conf. on Radiation and Remote Probing of the Atmosphere 1974 p. 135-161 refs CSCL 04A

Results from numerical computations of the diffuse sky radiances and degree of polarization are given. These results were obtained by solving the equation of radiative transfer in the formulation of Eschelbach. In these computations, multiple scattering and absorption by aerosol particles were considered. In addition, results from experimental measurements of the complex refractive index of aerosol samples at different relative humidities are presented. From the radiation flux divergences which were computed based on the determined properties of atmospheric aerosol particles, atmospheric heating rates were derived which were found to be comparable to the heating rates by water vapor.

N75-11554* Leiden Univ. Observatory (Netherlands). MULTIPLE SCATTERING IN CLOUD LAYERS; SOME RESULTS

H. C. VanDeHulst In Calif. Univ. Proc. of the UCLA Intern. Conf. on Radiation and Remote Probing of the Atmosphere 1974 p 162-195 refs CSCL 04A

Theoretical methods are discussed for calculating radiative effects of aerosols. Experimental determination is emphasized for relevant aerosol parameters on a global basis to arrive at realistic estimates of heating and cooling. Internal radiation fields in very thin and very thick slabs are reviewed. Phase functions, polarization, emission by internal sources, and path length distribution are also considered.

J.A.M.

N75-11555* Ljubljana Univ. (Yugoslavia). Dept. of Physics. SOME ANALYTICAL RESULTS FOR RADIATIVE TRANSFER IN THICK ATMOSPHERES

I. Kuscer and N. J. McCormick In Calif. Univ. Proc. of the UCLA Intern. Conf. on Radiation and Remote Probing of the Atmosphere 1974 p 196-226 refs Sponsored in part by NSF

CSCL 04A

Singular eigenmode expansions are a convenient analytical tool with which to study problems of monochromatic radiative transfer in thick or semi-infinite atmospheres. Some closed-form solutions are presented for anisotropic scattering, with the neglect of polarization effects. A basic ingredient for applications to the semi-infinite medium is Chandrasekhar's H-function, which is best defined through the Wiener-Hopf factorization. The solutions of the Milne and albedo problems are discussed in order that the method of matched asymptotic approximations may be used to

describe the solution for a thick atmosphere adjacent to a diffusely reflecting ground. Expressions for the emerging distributions are quoted. A possible extension of the general scheme to problems involving polarization is indicated.

N75-11556* California Univ., Los Angeles. Dept. of Physics. LECTURES ON THE SCATTERING OF LIGHT

David S. Saxon In its Proc. of the UCLA Intern. Conf. on Radiation and Remote Probing of the Atmosphere 1974 p 227-308 refs CSCL 04A

The exact (Mie) theory for the scattering of a plane wave by a dielectric sphere is presented. Since this infinite series solution is computationally impractical for large spheres, another formulation is given in terms of an integral equation valid for a bounded, but otherwise general array of scatterers. This equation is applied to the scattering by a single sphere, and several methods are suggested for approximating the scattering cross section in closed form. A tensor scattering matrix is introduced, in terms of which some general scattering theorems are derived. The application of the formalism to multiple scattering is briefly considered.

N75-11557* Leningrad (A. A. Zhdanov) State Univ. (USSR). INFLUENCE OF THE ATMOSPHERE ON SPECTRAL RADIANCE AND CONTRASTS OF NATURAL FORMATIONS MEASURED FROM SPACE

K. Ya. Kondratyev, A. A. Buznikov, O. B. Vasilyev, and O. I. Smokty In Calif. Univ. Proc. of the UCLA Intern. Conf. on Radiation and Remote Probing of the Atmosphere 1974 p 309-336 refs CSCL 04A

The method of transfer functions is discussed that enable the elimination of atmospheric effects from spectral photometric data measured from spacecraft. Soyuz 7, 9 and Salyut spacecraft data were used. The results made it possible to estimate the variations in the transfer functions in the visible spectral region depending on the albedo of the underlying surface both for the uniform surfaces and near the boundary between two uniform underlying surfaces.

J.A.M.

N75-11558* California Univ., Los Angeles. Dept. of Meteorology.

PARTICULATE SIZES FROM POLARIZATION MEASURE-MENTS

Jacob G. Kuriyan *In its* Proc. of the UCLA Intern. Conf. on Radiation and Remote Probing of the Atmosphere 1974 p 337-366 refs Sponsored in part by DOT and NASA

CSCL 04A

Radiation measurements from a ground based polarimeter were used to infer the optical properties of atmospheric particles. The inherent nonuniqueness in model calculations is discussed. The inverse problem in atmospheric optics is described, along with incident and emergent beams. Emergent radiation was calculated for all possible particulate distributions, and results were catalogued.

J.A.M.

N75-11559* Stanford Research Inst., Mento Park, Calif. Atmospheric Sciences Lab.

LIDAR OBSERVATIONS OF ATMOSPHERIC PARTICULATE CONTENT

R. T. H. Collis, P. B. Russell, E. E. Uthe, and W. Viezee In Calif. Univ. Proc. of the UCLA Intern. Conf. on Radiation and Remote Probing of the Atmosphere 1974 p 367-394 refs

CSCL 04A

The role of lidar in atmospheric studies concerned with radiative energy transfer and remote sensing is summarized. The application of lidar to climatic change investigations is proposed.

J.A.M.

N75-11560° National Center for Atmospheric Research, Boulder, Colo.

METHODS OF CALCULATING INFRARED TRANSFER: A REVIEW

02 ENVIRONMENTAL CHANGES AND CULTURAL RESOURCES

John C. Gille In Calif. Univ. Proc. of the UCLA Intern. Conf. on Radiation and Remote Probing of the Atmosphere 1974 p 395-430 refs Sponsored by NSF

CSCL 04A

Methods for calculating the transfer of infrared radiation in planetary atmospheres are summarized. Direct spectral integration is compared with high precision measurements, suggesting that calculational techniques are sufficient for molecular atmospheres. Infrared transfer is used to study the atmospheric energy budget and atmospheric structures.

J.A.M.

N75-12391*# Environmental Research Inst. of Michigan, Ann Arbor.

STUDY OF ATMOSPHERIC EFFECTS IN SKYLAB DATA Quarterly Progress Report, 1 Jun. - 31 Aug. 1974 Frederick J. Thomson, Principal Investigator 6 Nov. 1974 3 p EREP

(Contract NAS9-13272)

(E75-10006; NASA-CR-140728; ERIM-101700-17-L; QPR-6) Avail: NTIS HC \$3.25 CSCL 05B

N75-12393*# Boeing Co., Kent, Wash.
QUANTITATIVE DETERMINATION OF STRATOSPHERIC
AEROSOL CHARACTERISTICS Monthly Report, Oct. 1974
David L. Tingey, Principal Investigator Oct. 1974 1 p EREP
(Contract NAS9-13303)
(E75-10010: NASA-CR-140656) Avail: NTIS HC \$3.25 CSCL
04A

N75-12397*# Cornell Univ., Ithaca, N.Y. Dept. of Natural Resources.

EVALUATION OF SKYLAB IMAGERY AS AN INFORMATION SERVICE FOR INVESTIGATING LAND USE AND NATURAL RESOURCES Progress Report

Ernest_E. Hardy.-Principal-Investigator [1974] 18 p EREP (Contract NAS9-13364)

(E75-10018; NASA-CR-140732) Avail: NTIS HC \$3.25 CSCL 05B

N75-12398*# Geological Survey, Reston, Va.
URBAN AND REGIONAL LAND USE ANALYSIS: CARETS
AND CENSUS CITIES EXPERIMENT PACKAGE
Monthly
Progress Report, Jul. - Aug. 1974
Robert Alexander, Principal Investigator and Harry F. Lins, Jr.

20 Aug. 1974 4 p EREP

(NASA Order T-5290-B) (E75-10019; NASA-CR-140733) Avail: NTIS HC \$3.25 CSCL 08B

The author has identified the following significant results. The most significant finding has been the ability of the S-190B data to produce land use maps not far removed from the quality of high altitude aircraft photography generated maps.

N75-12399*# Environmental Research Inst. of Michigan, Ann

STUDY OF RECREATIONAL LAND AND OPEN SPACE USING SKYLAB IMAGERY Monthly Progress Report, Sep. 1974

Irvin J. Sattinger. Principal Investigator 15 Oct. 1974 2 p EREP (Contract NAS9-13283) (E75-10020; NASA-CR-140734; ERIM-103300-35-L) Avail: NTIS HC \$3.25 CSCL 088

N75-12400*# National Oceanic and Atmospheric Administration, Washington, D.C.
A CLOUD PHYSICS INVESTIGATION UTILIZING SKYLAB DATA Quarterly Progress Report, Jul. - Sep. 1974

John Alishouse, Herbert Jacobowitz, and David Wark, Principal Investigators Sep. 1974 4 p EREP (NASA Order T-4715-8)

(E75-10021; NASA-CR-140735; QPR-6) Avail: NTIS HC \$3.25 CSCL 04A

N75-12404*# Environmental Research Inst. of Michigan, Ann Arbor.
STUDY OF RECREATIONAL LAND AND OPEN SPACE USING SKYLAB IMAGERY Monthly Progress Report, Oct. 1974
Irvin J. Sattinger, Principal Investigator 20 Nov. 1974 5 p refs EREP (Contract NAS9-13283)

(E75-10034; NASA-CR-140771; ERIM-103300-37-L) Avail: NTIS HC \$3.25 CSCL 08B

N75-12406*# Environmental Research and Technology, Inc., Lexington, Mass.

EXPERIMENTAL EVALUATION OF ATMOSPHERIC EFFECTS ON RADIOMETRIC MEASUREMENTS USING THE EREP OF SKYLAB Quarterly Progress Report, Aug. - Oct. 1974

David T. Chang, Principal Investigator Oct. 1974 14 p EREP (Contract NAS9-13343)

(E75-10036; NASA-CR-140773; QPR-6) Avail: NTIS HC \$3.25 CSCL 048

N75-12425*# ECON, Inc., Princeton, N.J.
THE ROLE OF ERTS IN THE ESTABLISHMENT AND OF A
NATIONWIDE LAND COVER INFORMATION SYSTEM

Philip Abram and Jay Tullos 31 Oct. 1974 302 p refs Original contains color illustrations (Contract NASw-2558)

(NASA-CR-141045; Rept-74-2001-7) Avail: NTIS HC \$9.25 CSCL 05B

The economic potential of utilizing an ERTS type satellite in the development, updating, and maintenance of a nation-wide land cover information system in the post-1977 time frame was examined. Several alternative acquisition systems were evaluated for land cover data acquisition, processing, and interpretation costs in order to determine, on a total life cycle cost basis, under which conditions of user demand (i.e., area of coverage, frequency of coverage, timeliness of information, and level of information detail) an ERTS type satellite would be cost effective, and what the annual cost savings benefits would be. It was concluded that a three satellite system with high and low altitude aircraft and ground survey team utilizing automatic interpretation and classification techniques is an economically sound proposal.

Author

N75-12459# Varian Associates, Palo Alto, Calif.
AIRBORNE VAPOR SURVEILLANCE SYSTEM Final Report
James T. Arnold and Paul Robbiano Jun. 1974 45 p refs
(Contract DAAD05-70-C-0197)

(AD-785585; LWL-CR-03P68B) Avail: NTIS CSCL 07/4

The report describes a transportable membrane separatormass spectrometer system for the detection of trace vapors in ambient atmosphere. It employs a Uewellyn membrane separator, a quadrupole mass filter, and an on line system for data processing and display. Design electronics, and software are discussed, and installations in a mobile van and a fixed-wing aircraft are described. GRA

N75-12482*# National Aeronautics and Space Administration. Marshall Space Flight Center, Huntsville, Ala.

APPLICATION OF A SINGLE LASER DOPPLER SYSTEM TO THE MEASUREMENT OF ATMOSPHERIC WINDS William C. Cliff and Robert M. Huffaker Oct. 1974 23 p refs (NASA-TM-X-64891) Avail: NTIS HC \$3.25 CSCL 04B

The feasibility of employing a single laser Doppler velocimeter (LDV) system to remotely measure one-, two-, and three-dimensional velocity components in atmospheric flow fields is presented. A focused continuous wave CO2 laser emitting at the 10.6 mu wavelength is used as the laser source. Scan configurations employed by the LDV system were single-point, two-point, conical, and spiral conical. Test results are presented, which include favorable comparisons of velocity components measured by conventional anemometry and the LDV system. The feasibility using a single-beam LDV employing a conical scan technique for measuring two- and three-dimensional mean winds. Measurements to 300 m in dense fogs showed the ability of the LDV system to operate in dense fogs.

N75-12463*# Scientific Translation Service, Santa Barbara, Calif. DUST STORMS ACCORDING TO DATA OF SPACE RESEARCH

A. A. Grigoryev and V. B. Lipatov Washington NASA Nov. 1974 39 p refs Transl into ENGLISH of the book "Pylnyye Buri po Dannym Kosmicheskikh Issledovaniy" Moscow, Hydrometeorological Press, 1974 p 1-31 (Contract NASw-2483)

(NASA-TT-F-16021) Avail: NTIS HC \$3.75 CSCL 08E

The main regions of dust storms in the arid and sub-arid zones and the transfer paths of the dust materials were studied. Relationships between the dust storms and synoptic character and composition of the underlying surfaces are analyzed. The feasibility of studying the dynamics of these phenomena using space imagery is shown. The possibility of using microphotometry from TV pictures is considered for estimating the structure of dust storms.

Author

N75-12467# Defence Research Establishment Ottawa (Ontario). Research and Development Branch.

AN EXAMINATION OF WHITEOUTS AND TRANSMISSION THROUGH ICE FOG IN THE VISIBLE AND NEAR INFRARED

R. J. Brown, J. D. R. Pattman, and D. J. DiFruscio Jul. 1974 20 p. refs

(DREO-TN-74-18) Avail: NTIS HC \$3.25

Closed circuit television was investigated as a means of improving contrast during whiteout conditions at Yellowknife, N.W.T., in December, 1973. Cameras sensitive in the visible and near infrared portions of the electromagnetic spectrum were used. It is predicted that the reduction of snow reflectance in the near infrared might improve contrast perception in this region over that in the visible. Further studies included an investigation of the ability of these cameras to penetrate ice fog and the measurements of sky illuminance using a four spectral channel radiometer. No appreciable advantage over unaided eye observations was detected, and, in most cases, binoculars provided the most effective visual aid.

N75-12546*# Alabama Univ., Huntsville.
DETERMINATION OF POINTS OF ENTRY FOR POTENTIAL
CONTAMINANTS INTO LIMESTONE AQUIFERS USING

THERMAL INFRARED IMAGERY Interim Report F. L. Doyle 31 Oct. 1974 19 p refs (Contract NAS8-30216)

(NASA-CR-120540) Avail: NTIS HC \$3.25 CSCL 061

Lineations were identified involving the application of ERTS imagery to geologic and hydrologic problems. Interpretation of the southwest Madison County area is discussed. The tracing of the Beech Grove lineament to the northern boundary of Madison County. Alabama raises the question of its relationship to the trend of lineations in southwestern Madison County. The use of thermography as an indication of soil moisture is reviewed. The effect of soil moisture on surface temperature and the relationship between soil moisture and ground water are examined. J.A.M.

N75-13063# Tennessee Univ., Knoxville.
INVESTIGATION OF THE UTILITY OF KINETICALLY
CONTROLLED AND CATALYTIC REACTIONS TO PERFORM
ENVIRONMENTAL DETERMINATIONS BY PARALLEL
SPECTROPHOTOMETRIC ANALYSIS Ph.D. Thesis

J. L. Bowling Mar. 1974 102 p refs Sponsored by AEC (TID-26594) Avail: NTIS HC \$5.25

The analytical performance of the GeMSAEC spectrophotometric analyzer and the principles of parallel analysis which it employs were evaluated with respect to several important environmental species. The GeMSAEC system used was interfaced with a small computer for data collection and treatment. The production of formaldehyde from the selective cleavage of the double bond of 4-allyl 2-methoxy phenol (eugenol) by ozone permitted the specific determination of ozone in air samples. A rapid ion exchange separation converted the methylene blue spot test for selenium into a selective and sensitive method for quantitative determination of trace selenium. A modified GeMSAEC system was used for chemiluminescence studies. The use of EDTA as a general masking agent permitted the determination of chromium using the transition metal catalyzed reduction of luminol by basic hydrogen peroxide. The ability of the GeMSAEC to perform rapid, accurate, and reproducible Author (NSA) analyses was demonstrated.

N75-13334*# National Aeronautics and Space Administration. Lyndon B. Johnson Space Center, Houston, Tex.
THE ERTS-1 INVESTIGATION. VOLUME 7: ERTS-1 LAND-USE ANALYSIS OF THE HOUSTON AREA TEST SITE Final Report, Jul. 1972 - Jun. 1973
R. Bryan Erb, Principal Investigator Jun. 1974 110 p refs

Original contains imagery. Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue, Sioux Fall, S. D. 57198 ERTS

(E75-10009; NASA-TM-X-58124) Avail: NTIS HC \$5.25 CSCL 08B

N75-13337*# Brevard County Planning Dept., Titusville, Fla.
PLANNING APPLICATIONS IN EAST CENTRAL FLORIDA
Progress Report, 1 Aug. - 30 Sep. 1974

John W. Hannah, Principal Investigator, Garland L. Thomas, Fernando Esparza (NASA, Kennedy Space Center), and James J. Millard (NASA, Kennedy Space Center) 30 Sep. 1974 21 p FRTS

(Contract NAS5-21847)

(E75-10015; NASA-TM-X-72076) Avail: NTIS HC \$3.25 CSCL 08B

N75-13339*# Nebraska Univ., Lincoln.
SATELLITE PICTURES CAN HELP GUIDE SANDHILLS
IRRIGATION

James V. Drew, Principal Investigator, David T. Lewis, Leslie F. Sheffield, and Paul M. Seevers 27 Sep. 1974, 3 p. Repr. from Nebraska Farmer, 7 Sep. 1974, p. 38-40 Original contains

imagery. Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue, Sioux Falls, S. D. 57198

(Contract NAS5-21756)

(E75-10023; NASA-CR-140737) Avail: NTIS HC \$3.25 CSCL 08H

N75-13343*# Environmental Research Inst. of Michigan, Ann

OIL POLLUTION DETECTION, MONITORING AND LAW **ENFORCEMENT Quarterly Progress Report, Aug. 1974** Robert Horvath, Principal Investigator 22 Oct. 1974 2 p FREP

(Contract NAS9-13281)

(E75-10027; NASA-CR-140740; ERIM-101800-14-P) Avail: NTIS HC \$3.25 CSCL 08J

N75-13356*# Geological Survey, Reston, Va.

URBAN AND REGIONAL LAND USE ANALYSIS: CARETS AND CENSUS CITIES EXPERIMENT PACKAGE Monthly Progress Report, Oct. 1974

Robert Alexander, Principal Investigator and Harry F. Lins, Jr.

Oct. 1974 2 p EREP (NASA Order T-5290-B)

(E75-10056; NASA-CR-140964) Avail: NTIS HC \$3.25 CSCL

08B

N75-13369# Council for Scientific and Industrial Research, Pretoria (South Africa).

RESEARCH IN EARTH RESOURCES UTILIZATION. PHYSICAL ENGINEERING, AND POLLUTION] Annual Report, 1973

1974 87 p refs

(AR-29) Avail: NTIS HC \$4.75

The 1973 retrospective account of the various cooperative research activities provides photographs and descriptions of advancements in the following fields: (1) chemistry and physics: (2) geomagnetism; (3) astronomy; (4) telecommunications; (5) mechanical and electrical engineering; (6) water and food research; (7) air pollution; (8) road and building technology; and (9) timber and textile research. The information systems that were designed to disseminate the data gained from the research are also described A.A D

N75-13372*# National Aeronautics and Space Administration. Marshall Space Flight Center, Huntsville, Ala. COMPUTER AND PHOTOGRAMMETRIC GENERAL LAND

R. R. Jayroe, P. A. Larsen, and C. W. Campbell Oct. 1974 114 p refs Original contains color illustrations

USE STUDY OF CENTRAL NORTH ALABAMA

(NASA-TR-R-431: M-130) Avail: NTIS HC \$5.25 CSCL 08B The object of this report is to acquaint potential users with two computer programs, developed at NASA, Marshall Space Flight Center. They were used in producing a land use survey and maps of central north Alabama from Earth Resources Technology Satellite (ERTS) digital data. The report describes in detail the thought processes and analysis procedures used from the initiation of the land use study to its completion, as well as a photogrammetric study that was used in conjunction with the computer analysis to produce similar land use maps. The results of the land use demonstration indicate that, with respect to computer time and cost, such a study may be economically and realistically feasible on a statewide basis. Author

N75-13419# Air Force Avionics Lab., Wright-Patterson AFB.

AN APPLICATION OF INFRARED REMOTE SENSING TECHNIQUES TO ECOLOGICAL PROBLEMS

L. W. Crouch and R. D. Mower Jul. 1974 48 p (AD-786028; AFAL-TR-74-98) Avail: NTIS CSCL 13/2

The report and compendium of aerial infrared imagery was extracted from an R and D laboratory test of an Air Force infrared scanner. The purpose of the R and D test was to demonstrate the ability of one Air Force scanner to display thermal differentials found in a ground scene. The flight test area included a series of waterways and contiguous shorelines which illustrate ecological thermal phenomena. Specific examples of imagery were extracted from the R and D test program data for inclusion in this report. A summary of pertinent infrared technology is presented which relates sensor performance to atmospheric conditions, target and background characteristics, as well as image information content. In addition, brief photographic interpretation comments are included for each infrared image.

Author (GRA)

N75-13437# Suomalainen Tiedeakatemia, Helsinki (Finland). IONOSPHERIC MEETING

Oct. 1973 12 p Meeting held at Sodankyla, 14-16 May 1973

(Rept-3) Avail: NTIS HC \$3.25 CSCL 03B

Techniques in ionogram scaling are reviewed, along with parameters and difficulties in latitude ionospheric sounding. Blank and fixed gain ionograms are proposed for publication. Experiments on ionogram sequencing, fixed frequency recordings, interferometers, and polarimeters are presented.

N75-13452*# Chicago Univ., III. Dept. of the Geophysical

REMOTE SOUNDING OF STRATOSPHERIC TEMPERA-TURES USING HIGH RESOLUTION RADIANCE MEASURE-MENTS FROM THE IRIS-D M.S. Thesis

William O. Gallery Dec. 1974 119 p refs (Grant NGR-14-001-194)

(NASA-CR-141130) Avail: NTIS HC \$5.25 CSCL 04A

Remote sounding of stratospheric temperatures up to 3.2 millibars is attempted using high resolution (unapodized) radiance measurements in the 15 micron CO2 band from the infrared Interferometer Spectrometer on Nimbus 4. Inversions are performed using the Chahine relaxation technique. Radiance data and simultaneous in situ temperature profiles are obtained from the Rocket/Nimbus Sounder Comparison. Numerical tests with synthetic radiance data show that the uncertainty in the retrieved temperatures due to random instrument noise is about 1.1 K when averaged over layers about 10 km thick. However, comparison of the measured radiances with the radiances calculated from the in situ profiles show the calculated radiances to be systematically higher than the measured radiances. The evidence indicates that systematic errors exist in both the radiance and the in situ measurements. Author

N75-13458# California Univ., Livermore. Lawrence Livermore Lab

ENVIRONMENTAL DEFINITION PROGRAM Final Report, period ending 30 Jun. 1974

J. B. Knox and J. E. Lovill 25 Jul. 1974 12 p Sponsored in part by DNA

(Contract W-7405-eng-48)

(UCID-16565) Avail: NTIS HC \$3.25

Satellite monitoring of the stratosphere and troposphere is conducted to provide thousands of vertical profiles each day of temperature, water vapor, ozone, ice and liquid cloud contents, and various discrete radiational emission spectra. The data acquired and computer codes being developed to analyze the data are discussed NSA

N75-13479 Freie Univ., Berlin (West Germany). Inst Fuer Meteorologie.

METEOROLOGICAL DATA. VOLUME 145 NO. 1: EUROPEAN CLIMATOLOGY 1974 METEOROLOGICAL SATELLITE DATA FROM THE BERLIN ZONE OF RECEP-TION, PART 1, 1ST QUARTER [METEOROLOGISCHE ABHANDLUNGEN. BAND 145, HEFT 1: DAS EUROPAEI-SCHE WETTERBILD 1974, WETTERSATELLITEN-DATEN AUS DEM BERLINER EMPFANGSBEREICH, TEIL 1, 1. VIERTELJAHR)

1974 199 p In GERMAN

Avail: Issuing Activity

63 1 1

The present volume continues the daily European weather maps for the first quarter, 1974. Collected are APT photographs from ESSA 8 and film strips from the NOAA 2 scanning radiometer. Transl, by G.G.

N75-13483 Freie Univ., Berlin (West Germany). Inst. for Meteorology

METEOROLOGICAL DATA. VOLUME 139, NO. 4: THE EUROPEAN CLIMATOLOGY, 1973. METEOROLOGICAL SATELLITE DATA FROM THE BERLIN ZONE OF RECEP-TION, PART 4, FOURTH QUARTER [METEOROLOGISCHE ABHANDLUNGEN, BAND 139, HEFT 4: DAS EUROPAEIS-CHE WETTERBILD 1973 WETTERSATELLITEN-DATEN AUS DEM BERLINER EMPFANGSBEREICH, TEIL 4, 4. VIERTEL-JAHR]

1973 202 p In GERMAN Avail: Issuing Activity

The present volume continues the daily European weather maps for the 4th quarter in 1973. Collected are APT photographs from ESSA 8 and film strips from the NOAA 2 scanning Transl. by G.G. radiometer.

N75-14198 *# Boeing Co., Kent, Wash.

QUANTITATIVE DETERMINATION OF STRATOSPHERIC AEROSOL CHARACTERISTICS Monthly Report, Nov. 1974 David L. Tingey, Principal Investigator Nov. 1974 1 p EREP (Contract NAS9-13303)

(E75-10079; NASA-CR-141092) Avail: NTIS HC \$3.25 CSCL 04A

N75-14201 *# Ohio Dept. of Economic and Community Development, Columbus.

THE APPLICATION OF SKYLAB EREP DATA FOR LAND USE CLASSIFICATION AND MAPPING IN THE CLEVELAND AND COLUMBUS AREAS

David C. Sweet, Principal Investigator, Paul Pincura, George E. Wukelinc, Joachim G. Stephan, Harry E. Smail, and Thomas F. Ebbert Apr. 1974 28 p Prepared in cooperation with Battelle Columbus Labs., Ohio Original contains color imagery. Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue, Sioux Falls, S. D. 57198 EREP (NASA Order C-21372-C)

(E75-10083; NASA-CR-141107) Avail: NTIS HC \$3.75 CSCL 088

N75-14226 Telespazio, S.p.A., Rome (Italy). DIGITAL LAND-USE CLASSIFICATION USING AN ERTS-1 IMAGE

B. Ratti and F. Capozza In ESRO European Earth-Resources Satellite Expts. May 1974 p 93-105 refs

Initial work on digital land use classification was carried out utilizing an ERTS-1 MSS frame for an area centered on Rome. The high variability of ground cover types, inherent in the region under investigation, had led to use clustering (i.e. unsupervised) techniques rather than supervised techniques, which call for acquisition of substantial amounts of ground-truth information to be effective. In this initial approach, a simplified clustering algorithm was used. The ground cover types were divided into 12 classes, plus one class for points not assigned to any of these 12. The region classified covers approximately 1000 sq km. The results were very encouraging. Further work is being carried out to improve the clustering procedures, with accompanying improvements in classification accuracy. Author (ESRO)

N75-14245 Strasbourg Univ. (France).

ERTS 1 CONTRIBUTIONS TO THE ECOGENETICAL KNOWLEDGE OF THE LLANOS OF THE ORINOCO (COLOMBIA AND VENEZUELA) [APPORTS DE ERTS-1 A NOTRE CONNAISSANCE ECOGENETIQUE DES LLANOS DE L'ORENOQUE (COLOMBIE ET VENEZUELA)

J. Tricart In ESRO European Earth Resources Satellite Expts. May 1974 p 317-324 refs in FRENCH; ENGLISH summary

Satellite multispectral photographs were used for reconnaissance of the Orinoco Llanos. These photographs, combined with verification on the ground, provide a small-scale overall view which helps to bring out the main features of the ecogenesis. The sandy deposits built up along the rivers up to about 6 deg N were affected by strong wind action, which formed slightly bow-shaped longitudinal dunes lying in a direction between ENE-WSW and NNE-SSW. At the western and southern extremities of the region the dunes form folds on the southern banks of the rivers, invading the edge of sand-clay interfluves formed from older materials. The dunes were partially drowned by Holocene silt-clay material and probably date from the pre-Flandrian regression. Dune plant cover degradation, by cattle during flooding, leads to wind reactivation. Any rational development of this area should take these phenomena into account.

Author (ESRO)

N75-14248 Centre d'Etudes Phytosociologiques et Ecologiques. Montpellier (France).

QUALITATIVE STUDY OF SOME ERTS-1 IMAGES OF THE GOLION PROJECT. ECOLOGICAL ASPECTS OF TER-RESTRIAL ENVIRONMENTS [ETUDE QUALITATIVE DE QUELQUES IMAGES ERTS-1 DU PROJET GOLION. ASPECTS ECOLOGIQUES DES MILIEUX TERRESTRES]

G. Caballe, B. Lacaze, and G. Long In ESRO European Earth Resources Satellite Expts. May 1974 p 363-370 refs In FRENCH; ENGLISH summary Sponsored jointly by CNES and Direc des Rech. et Moyens d'Essais

The main results of an initial study of the ERTS-1 imagery (MSS, 4 channels) of the GOLION Project southern (France) are presented. The aim is to demonstrate that, from an objective analysis of the images on a scale of 1:1,000,000 studied channel-by-channel or by comparison, it is possible to discern homogeneous or isophenic zones which contribute to global thematic data (chiefly geomorphology, vegetation and land use). Over a test area of 1200 sq km, 24 zones were determined on the basis of the image criteria and their variations. The conclusions from this qualitative approach will serve as basic assumptions for a quantitative, multispectral study using digital data. Because of the erratic nature of the images received, it was not possible to make an effective time-sequence study. The value of ecographic satellite images for the definition of ecological regions is commented upon. Author (ESRO)

N75-14250 Reading Univ. (England). Dept. of Geography. A COMPARISON OF THE CAPABILITIES OF ERTS-1 AND THE SKYLARK EARTH-RESOURCES ROCKET FOR RE-SOURCES SURVEYING IN CENTRAL ARGENTINA

J. R. G. Townshend, R. A. G. Savigear, C. O. Justice, J. R. Hardy, D. S. H. Drennan, and C. J. Bray In ESRO European Earth Resources Satellite Expts. May 1974 p 381-403 refs Orginal contains color illustrations

(Contract AT/2035/025)

An evaluation of the potential of photographs from the earth-resources Skylark rocket was made in a joint Anglo-Argentinian project. Part of this program consists of a resources evaluation, and comparisons were made with imagery of the same area obtained from ERTS-1. These two space platform resource systems have several differences which affect their utility for different resource-evaluation tasks. With high resolution emulsion and long focal length cameras, the resolution of the Skylark photography is significantly better than that of ERTS imagery. Comparisons of the two sets of imagery were made for a variety of features.

Author (ESRO)

N75-14331# Deutsche Forschungs- und Versuchsanstalt fuer Luft- und Raumfahrt, Oberpfaffenhofen (West Germany). Inst. fuer Flugfunk und Mikrowellen.

THEORETICAL INVESTIGATIONS ON THE DETERMINATION OF THE VERTICAL TEMPERATURE DISTRIBUTION IN THE ATMOSPHERE ASSUMING MULTISPECTRAL THERMAL RADIATION MEASUREMENTS AT 60 GHz [THEORETISCHE UNTERSUCHUNGEN IM HINBLICK AUF DIE BESTIMMUNG DER TEMPERATURVERTEILUNG IN DER ATMOSPHAERE UNTER DER ANNAHME MULTISPEKTRALER MESSUNGEN DER THERMISCHEN STRAHLUNG BEI 60 GHz]

Friedrich Malota and Volker Stein Bonn Bundesmin, fuer Forsch, und Technol. Jul. 1974 71 p refs In GERMAN; ENGLISH summary Sponsored by Bundesmin, fuer Forsch, und Technol. (BMFT-FB-W-74-04) Avail: NTIS HC \$4.25; ZLDI, Munich 14,95 DM

Vertical atmospheric temperature distribution was derived from microwave thermal radiation measurements. The radiation transfer equation is discussed, notably correlation between environmental temperature and noise temperature, temperature distribution models and the influence of clouds on noise temperature. Information content and numerical solution methods of indirect temperature observations are dealt with, and noise temperature simulation calculations and inversion calculations for temperature profiles are presented.

N75-14366* National Aeronautics and Space Administration. Lyndon B. Johnson Space Center, Houston, Tex. SKYLAB ENVIRONMENTAL AND CREW MICROBIOLOGY

STUDIES

R. M. Brockett (School of Aerospace Med.), J. K. Ferguson, R. C. Graves, T. O. Groves, M. R. Henney, C. J. Hodapp, K. D. Kropp, J. L. McQueen (Natl. Cancer Inst.), B. J. Mieszkuc, F. J. Pipes et al. *In its* Proc. of the Skylab Life Sci. Symp., Vol. 1 Nov. 1974 p 121-143 refs

CSCL 06M

The results of some ground-based simulations have engendered theories that forecasted microbial simplification, intercrew transfer of microbial pathogens, autoinfections, and postflight microbial shock. In an effort to understand the effects of space flight, microbiological samples from multiple sites on the crewmembers were collected several times before, during, and after the space flights. The Skylab are related to analogous Apollo data and are discussed in a manner that will allow an evaluation of the validity of the hypotheses presented. Additionally, in-flight environmental samples were acquired from designated sites within the spacecraft and returned to earth for analysis. The resulting data were used to identify potential microbial problems for the maintenance of a habitable environment in the orbital workshop.

N75-15100 *# Cornell Univ. New York. Dept. of Natural Resources.

EVALUATION OF SKYLAB IMAGERY AS AN INFORMATION SERVICE FOR INVESTIGATING LAND USE AND NATURAL RESOURCES Progress Report

Ernest E. Hardy, Principal Investigator 31 Dec. 1974 8 p EREP (Contract NAS9-13364) (E75-10106: NASA-CR-141372) Avail: NTIS HC \$3.25 CSCL 08F

N75-15104*# Tokyo Univ. (Japan).
INVESTIGATION OF THE ENVIRONMENTAL CHANGE PATTERN OF JAPAN Final Report, Jul. 1972 - Aug. 1973
Takakazu Maruyasu, Principal Investigator Oct. 1973 . 58 p
Sponsored by NASA Original contains imagery. Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue. Sioux Falls. S. D. 57198 ERTS (E75-10086; NASA-CR-141150) Avail: NTIS HC \$4.25 CSCL 048

The author has identified the following significant results. * P -> b f . 8 ERTS-1 imagery clearly identifies the relationships between the status of erosion, effluent patterns affected by the coastal current, and the cultural construction activities. Simple photographic techniques can be used for detecting water mass distribution separately from cloud cover and also noise caused by reflected sunlight from wave surfaces. Polluted water does not diffuse continuously into the oceanic water, but forms masses in the water in the Kuroshio area. The polluted or turbid water in the area just north of the Tomogashima Channel, the south outlet of the Osaka Bay, shows that the northward tidal current runs in a clockwise eddy at the tidal period when the imagery was taken. Such an eddy-like pattern of tidal current had never been revealed by conventional oceanographic data. A front between an oceanic water mass and a polluted water mass runs in a NW-SE direction in the central part of the Osaka Bay. The patterns of turbid water discharged from the Kii River and Yoshino River show a northward tidal current in the North Kii Straits. The pattern of lighter turbid or polluted water located in the northwest region of the North Kii straits suggests the existence of a clockwise eddy in the straits.

N75-15109*# Geological Survey, Reston, Va.
URBAN AND REGIONAL LAND USE ANALYSIS: CARETS
AND CENSUS CITIES EXPERIMENT PACKAGE
Progress Report

Monthly

Robert Alexander, Principal Investigator, Daniel B. Gallagher, and Harry F. Lins. Jr. 20 Nov. 1974 3 p EREP (NASA Order T-5290-B)

(E75-10091; NASA-CR-141155) Avail: NTIS HC \$3.25 CSCL

08B

N75-15110*# Environmental Research Inst. of Michigan, Ann Arbor.

STUDY OF RECREATIONAL LAND AND OPEN SPACE USING SKYLAB IMAGERY Monthly Progress Report, Nov. 1974

Irvin J. Sattinger, Principal Investigator 18 Dec. 1974 3 p

(Contract NAS9-13283)

(E75-10092; NASA-CR-141156; ERIM-103300-40-L) Avail: NTIS HC \$3.25 CSCL 08B

The author has identified the following significant results. Despite almost uniform surface temperature conditions in the study area, the thermal imagery did illustrate the following possible uses: (1) Surface temperatures relative to 0 C reveal whether the snow and ice cover is wet and the melt pattern. This information is useful in hydrologic monitoring of runoff timing and rate, as well as indicating trafficability conditions on the snow. (2) When the surface temperature of snow and ice is below freezing, it may serve as an indicator of spatial variation of air temperatures. This information may be statified calculating the spatial variation of surface radic.

ing synoptic weather condition changes or local microclimatic effects. (3) Frozen inland lakes with less than about three or four inches of snow over the ice may be differentiated from surrounding snow covered land areas; this is not always feasible in visible wavelength imagery. The feasibility of this application decreases as the ice thickness increases.

N75-15114*# Environmental Research Inst. of Michigan, Ann

DETERMINATION OF THE EARTH'S AEROSOL ALBEDO USING SKYLAB DATA Quarterly Progress Report, Aug. -Dec. 1974

Robert E. Turner, Principal Investigator 13 Jan. 1975 2 p EDED

(Contract NAS9-13279)

(E75-10096; NASA-CR-141292; ERIM-102200-14-L) Avail: NTIS HC \$3.25 CSCL 04A

N75-15117*# National Oceanic and Atmospheric Administration, Washington, D.C.

A CLOUD PHYSICS INVESTIGATION UTILIZING SKYLAB DATA Quarterly Progress Report, Oct. - Dec. 1974

John Alishouse, Herbert Jacobowitz, and David Wark, Principal Investigators Dec. 1974 5 p EREP

(NASA Order T-4715-B) (E75-10099; NASA-CR-141295: QPR-7) Avail: NTIS HC \$3.25 CSCL 04A

N75-15118*# Geological Survey, Reston, Va.

URBAN AND REGIONAL LAND USE ANALYSIS: CARETS AND CENSUS CITIES EXPERIMENT PACKAGE Monthly **Progress Report**

Robert Alexander, Principal Investigator and Harry F. Lins, Jr. 27 Dec. 1974 4 p EREP

(NASA Order T-5290-B)

(E75-10100; NASA-CR-141296) Avail: NTIS HC \$3.25 CSCL 08B

N75-15121*# Maine State Highway Dept., 8angor. MULTIDISCIPLINARY ANALYSIS FOR HIGHWAY ENGINEERING PURPOSES Progress Report

Ernest G. Stoeckeler, Raymond G. Woodman, and Robert S. Farrell, Principal Investigators 6 Nov. 1974 7 p EREP (Contract NAS9-13359)

(E75-10103; NASA-CR-141304; PR-4) Avail: NTIS HC \$3.25 CSCL 13B

N75-15123*# Colorado School of Mines, Golden. Dept. of Geology

GEOLOGIC AND MINERAL AND WATER RESOURCES INVESTIGATIONS IN WESTERN COLORADO, USING SKYLAB EREP DATA Monthly Progress Report, Dec. 1974 Keenan Lee, Principal Investigator 20 Jan. 1975 4 p EREP (Contract NAS9-13394)

(E75-10105; NASA-CR-141306) Avail: (NTIS HC \$3.25 CSCL 08G

N75-15126*# Jet Propulsion Lab., Calif. Inst. of Tech., Pasadena. THE AVAILABILITY OF LOCAL AERIAL PHOTOGRAPHY IN SOUTHERN CALIFORNIA

renda Sledge, Charles K. Paul, and Albert J. 74 J26ip. Walter Aller Landini 1 L

(Contract NAS7-100)

(NASA-CR-141314; JPL-SP-43-14) Avail: NTIS HC \$3.75 CSCL 14E

Some of the major photography and photogrammetric suppliers and users located in Southern California are listed. Recent trends in aerial photographic coverage of the Los Angeles basin area are also noted, as well as the uses of that imagery. Author

N75-15135# California Univ., Berkeley. School of Forestry and Conservation

USEFULNESS OF REMOTE SENSING TECHNIQUES FOR THE ENVIRONMENTAL MONITORING OF THE SACRAMENTO-SAN JOAQUIN DELTA Final Report

William C. Draeger 1 Feb. 1974 55 p refs Sponsored by Calif. Dept. of Water Resources

Avail: NTIS HC \$4.25

Information is presented on the capabilities of using remote sensing technology in an environmental monitoring program for the Sacramento-San Joaquin delta. Remote sensing literature pertaining to water quality measurement was reviewed to define the current state-of-the-art and to ascertain the possibility of evaluating the water quality parameters. Based on the findings of the review, a number of experimental aerial photo flights over the delta were conducted and coordinated with on-thewater measurements. Water quality parameters, the measurement accuracies needed, and the required frequency or timing of measurement are discussed. It was found that annual, smallscale color photographic coverage and periodic hand-held photography from a light plane permit subjective evaluations of direct on-the-water measurements. M.J.S.

N75-15199*# Miami Univ., Coral Gables, Fla. Dept of Mechanical and Industrial Engineering.

FEASIBILITY OF REMOTE SENSING FOR DETECTING THERMAL POLLUTION. PART 1: PART 2: IMPLEMENTATION PLAN FEASIBILITY STUDY.

T. Nejat Veziroglu and Samuel S. Lee Dec. 1973 184 p refs (Contract NAS10-8402)

(NASA-CR-134453) Avail: NTIS HC \$7.00 CSCL 13B

A feasibility study for the development of a three-dimensional generalized, predictive, analytical model involving remote sensing, in-situ measurements, and an active system to remotely measure turbidity is presented. An implementation plan for the development of the three-dimensional model and for the application of remote sensing of temperature and turbidity measurements is outlined. J.M.S.

N75-15202# California Univ., Livermore. Lawrence Livermore

ATMOSPHERIC RELEASE ADVISORY CAPABILITY: RESEARCH AND PROGRESS

J. B. Knox and T. V. Crawford 8 May 1974 33 p Presented at 2d AEC Pollution Control Conf., Albuquerque, New Mexico, 16 Apr. 1974 Revised

(UCRL-75644-Rev-2; Conf-740406-16) Avail: NTIS HC \$3.75

The purpose of the Atmospheric release advisory capability is to provide for real-time predictions of effects and dose-toman of atmospheric releases of hazardous materials as rapidly and accurately as possible, to provide environmental surveillance and monitoring information in real-time, and to provide regional transport modeling capabilities. Planned technical milestones are given.

03 GEODESY AND CARTOGRAPHY

Includes mapping and topography.

A75-10076 An electronic drawing aid for map compiling. R. R. Real, Y. Fujimoto, H. Moeller, and M. Paulun (National Research Council, Div. of Physics, Ottawa, Canada). *IEEE Transactions on Geoscience Electronics*, vol. GE-12, Oct. 1974, p. 134-139. 5 refs.

A versatile digital electronic drawing aid is described for assisting an operator in scribing map originals from stereo-orthophotos. These photographs permit the user to see an accurate, stable, and uniformly scaled stereo image of the earth's terrain. Direct measurement, planning and mapping within the stereo image is achieved with a simple instrument called a Stereocompiler. The drawing aid described has options that the operator of the Stereocompiler may use to facilitate the plotting of planimetry, contours, symbols, and the typescribing of alphanumeric annotation on map manuscripts while viewing the stereo image. (Author)

A75-11861 On the longitudinal localization of the substorm active region and its changes during the substorm. V. A. Sergeev (Leningradskii Gosudarstvennyi Universitet, Leningrad, USSR). Planetary and Space Science, vol. 22, Sept. 1974, p. 1341-1343, 10 refs.

During the expansive phase of a substorm the active region frequently changes its situation by a jump, both to the evening and morning sectors. The typical interval between successive activizations is often greater than about 10-20 min. It is emphasized that in such events the definition of a substorm phase for all the magnetosphere is ambiguous, because the conditions characteristic for different phases can be realized in different parts of the magnetosphere simultaneously.

(Author)

A75-11975 Solar wind and geomagnetic activity. V. Formisano, G. Moreno, and C. Signorini (CNR, Laboratorio per il Plasma nello Spazio, Frascati, Italy). *Nuovo Cimento, Sezione B*, vol. 23 B, Oct. 11, 1974, p. 473-488. 26 refs. Research supported by the Considio Nazionale delle Ricerche.

Interplanetary plasma and magnetic-field parameters, measured by the ESRO satellite HEOS-1, have been correlated with the D sub st, taken as an index of the overall geomagnetic activity. It is found that two different states exist for the interaction between the solar wind and the magnetosphere: they correspond to quiet periods (D sub st positive) and to disturbed conditions (main phase of geomagnetic storms - i.e., D sub st negative). In disturbed conditions the magnetosphere appears to be much more sensitive to variations of the solar-wind dynamical pressure than in periods of positive D sub st. (Author)

A75-14646 # Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars.

K. Arnold (Deutsche Akademie der Wissenschaften, Zentralinstitut für Physik der Erde, Potsdam, East Germany). Gerlands Beiträge zur Geophysik, vol. 83, no. 4, 1974, p. 249-269. 6 refs.

An investigation is conducted regarding the distance between a terrestrial observing station and a laser reflector on the moon, taking into account the functional relations between this distance and various parameters. Attention is given to the coordinates of the station, the parameters of polar motion, the deceleration of the earth's rotation, and the space coordinates of the reflector. The feasibility of a determination of the individual parameter values on the basis of the distance measurements is studied.

G.R.

A75-14683 Coherent optics potential applications to mapping. R. D. Leighty (U.S. Army, Engineer Topographic Laboratories, Fort Belvoir, Va.). (American Society of Photogrammetry and Society of Photo-Optical Instrumentation Engineers, Tutorial Seminar on Coherent Optics in Mapping, Rochester, N.Y., Mar. 27-29, 1974.) Optical Engineering, vol. 13, Sept.-Oct. 1974, p. 440-450. 96 refs.

Some of the roles for the potential application of coherent optical techniques in mapping systems are discussed. Other references on this subject are integrated into the overall framework of coherent optics in mapping processes. Potentials are defined in the eight areas of photogrammetric data reduction, image processing, optical memories, pattern recognition, performance prediction, holographic terrain displays, all-weather mapping, and field applications.

T.S.

A75-15435 • Mapping soil features from multispectral scanner data. S. J. Kristof and A. L. Zachary (Purdue University, West Lafayette, Ind.). *Photogrammetric Engineering*, vol. 40, Dec. 1974, p. 1427-1434. Grant No. NGL-15-005-112.

In being able to identify quickly gross variations in soil features, the computer-aided classification of multispectral scanner data can be an effective aid to soil surveying. Variations in soil tone are easily seen as well as variations in features related to soil tone, e.g., drainage patterns and organic matter content. Changes in surface texture also affect the reflectance properties of soils. Inasmuch as conventional soil classes are based on both surface and subsurface soil characteristics, the technique described here can be expected only to augment and not replace traditional soil mapping. (Author)

A75-16676

Coherent optics in mapping; Proceedings of the Tutorial Seminar and Technology Utilization Program, Rochester, N.Y., March 27-29, 1974. Seminar sponsored by the American Society of Photogrammetry and Society of Photo-optical Instrumentation Engineers. Edited by N. Balasubramanian (Rochester, University, Rochester, N.Y.) and R. D. Leighty (U.S. Army, Engineer Topographic Laboratories, Fort Belvoir, Va.). Palos Verdes Estates, Calif., Society of Photo-optical Instrumentation Engineers (SPIE Proceedings. Volume 45), 1974. 305 p. \$30.

The present work gathers together studies dealing with, first, the fundamental concepts of coherent optics and photogrammetry in themselves, and second, the combination of coherent optics technology with photogrammetric mapping techniques. Some of the topics covered include: photogrammetric and reconnaissance applications of coherent optics, holographic moving map display, the mensuration and mapping of holograms and holographic stereomodels, terrain measurement from holographic stereograms, a system for photogrammetric data reduction using coherent optics, and cloud screening from aerial photography applying coherent optical pattern recognition techniques.

P.T.H.

A75-17125 # Plutonic structure of the earth's crust on space images. V. I. Makarov, S. F. Skobelev, V. G. Trifonov, P. V. Florenskii, and Iu. K. Shchukin (Akademiia Nauk SSSR, Geologicheskii Institut, Moscow, USSR). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1.

1-1-19, 1974, Proceedings. Volume 1.

Examples from tectonically varied territories are analyzed to show that the plutonic structure of the earth's crust can be studied through the interpretation of black-and-white space photographs. Images with decreasing resolving power seem to reveal the structural elements of ever deeper layers. Applications of space photography to the study of deep seismically active zones and to oil and gas exploration are indicated.

A.T.S.

A75-17139 # Mapping of the 1973 Mississippi River floods by the NOAA-2 satellite. D. R. Wiesnet, D. F. McGinnis, and J. A. Pritchard (NOAA, Hillcrest Heights, Md.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 621-627. 6 refs.

A75-17201 # Single flight stereo radar capabilities. G. E. Carlson and G. L. Bair (Missouri, University, Rolla, Mo.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1633-1647. 6 refs. Contract No. N00014-69-A-0141-0008. NR Project 387-069.

An improved single-flight technique for obtaining stereo radar image pairs is briefly described, as well as a previously proposed single-flight technique and a previously implemented two-flight technique. The results of tradeoff and theoretical performance analyses are presented for the improved single-flight technique, and theoretical performance comparisons are made with the two previous techniques. The results of an error analysis for the improved single-flight technique are presented and compared with similar results for the other two techniques. (Author)

A75-17224 * # Surface compositional mapping in the Wind River Range and Basin, Wyoming by multispectral techniques applied to ERTS-1 data. B. Salmon and R. K. Vincent (Michigan, Environmental Research Institute, Ann Arbor, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 2005-2012. Contract No. NASS-21783.

A75-17228 # Elements of the deep structure of the earth's crust on the space images of the East Caucasus. V. G. Trifonov (Akademiia Nauk SSSR, Geologicheskii Institut, Moscow, USSR). In: International Symposium on Remote Sensing of Environment, 9th, Ann-Arbor, Mich., April-15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 2051-2053. 8 refs.

A75-18055 Automatic contouring on the Gestalt Photomapper 2-1. B. G. Crawley (Gestalt International, Ltd., Vancouver, Canada). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings.

Falls Church, Va., American Society of Photogrammetry, 1974, p. 50-56.

The Gestalt Photomapper II distorts the scanned images from a stereo-pair of aerial diapositives to simulate an objective view from infinity. The distortion is done under controlled conditions established by the model, and in accordance with the X parallaxes sensed by an electronic correlator. When the images have been perfectly registered, the height matrix in the correlator's memory provides the data-base for the extraction of elevation contours for the scanned images. With GPM II, a systematic profile of the model in a patch-like manner, produces an orthophoto and a contour sheet of the overlap areas. (Author)

A75-18062 * Small scale thematic mapping - A case for radar imagery. F. M. Henderson (New York, State University, Albany, N.Y.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings.

Falls Church, Va., American Society of Photogrammetry, 1974, p. 155-169. 24 refs. Contract No. NAS9-10261.

Small scale thematic maps (1:250,000 and smaller) of physical and cultural phenomena manifested on the landscape are a major concern to scientists and investigators in diverse disciplines. A strip of K-band radar imagery consisting of a traverse from eastern Minnesota to northern Utah was employed to evaluate the potential of radar imagery for small scale land use mapping. In the course of this investigation, it was discovered that certain borders derived from radar imagery were compatible with borders found on the nonland use thematic maps used for comparison. Specifically, numerous borders and regions of small scale maps of landforms, soils, vegetation, and geology are found to be similar to the radar land use regions. Although far from conclusive it appears that radar imagery can be employed in the small scale mapping of landforms and possibly for mapping physiognomic or economic vegetation. (Author)

A75-18106 Geodesy and physics of the earth; Proceedings of the Second International Symposium, Potsdam, East Germany, May 7-11, 1973. Parts 1 & 2. Symposium sponsored by the International Union of Geodesy and Geophysics. Edited by H. Kautzleben (Deutsche Akademie der Wissenschaften, Zentralinstitut für Physik der Erde, Potsdam, East Germany) and E. Buschmann. Potsdam, East Germany, Deutsche Akademie der Wissenschaften, Zentralinstitut für Physik der Erde (Zentralinstitut für Physik der Erde (Zentralinstitut für Physik der Erde, Veröffentlichungen, No. 30. Parts 1 & 2), 1974. Pt. 1, 215 p.; pt. 2, 306 p. In English and German.

The dynamics of the entire earth are considered, taking into account the earth rotation and gravity variations, the effect of the core on the earth's rotation, the dynamics of the outer core, and the distribution of the gravimeter factor obtained from earth-tide observations in Europe. Other topics discussed are related to recent earth crust movements, the figure of the earth and the gravity field, and special problems of geodesy and geophysics. Attention is given to new methods for studying the dynamics of the entire earth and to problems and consequences of modern hypotheses on the global tectonics for geodesy and gravimetry.

G.R.

A75-18109 # Problems and consequences of modern hypotheses on the global tectonics for geodesy and gravimetry. H. Kautzleben (Deutsche-Akademie der-Wissenschaften, Zentralinstitut für Physik der Erde, Potsdam, East Germany). In: Geodesy and physics of the earth; Proceedings of the Second International Symposium, Potsdam, East Germany, May 7-11, 1973. Part 1. Potsdam, East Germany, Deutsche Akademie der Wissenschaften, Zentralinstitut für Physik der Erde, 1974, p. 47-64. 68 refs.

A review is presented of the hypothesis of plate tectonics. The hypothesis provides a kinematic picture regarding the fundamental structures of the earth's tectonosphere and the phenomena occurring in it. The geodetic evidence of secular movement processes is discussed along with gravimetric investigations and problems which have to be considered in the investigation of the earth's rotational behavior.

G.R.

A75-18122 # The effect of the secular change of the terrestrial gravity field on elevation level differences (Der Einfluss der Säkularänderung des Erdschwerefeldes auf die nivellierten Höhenunterschiede). P. Biro (Budapesti Müszaki Egyetem, Budapest, Hungary). In: Geodesy and physics of the earth; Proceedings of the Second International Symposium, Potsdam, East Germany, May 7-11, 1973. Part 2. Potsdam, East Germany, Deutsche Akademie der Wissenschaften, Zentralinstitut für Physik der Erde, 1974, p. 315-322. In German.

The elevation difference in geodesy is the distance of the corresponding equipotential surfaces of the gravity potential. This parameter can be very accurately determined by repeated levellings and gravity measurements. Current scientific studies of recent

movements of the earth's crust are based on the assumption that the form and location of the equipotential surfaces will not change with time. An investigation of the involved relations shows that currently only relative movements can be determined.

A75-18123 # Applications of satellite geodesy in the exploration of the internal structure of the earth (Anwendungsmöglichkeiten der Satellitengeodäsie bei der Erforschung der inneren Struktur der Erde). G. Barta (Eotvos Lorand Tudomanyegyetem, Budapest, Hungary). In: Geodesy and physics of the earth; Proceedings of the Second International Symposium, Potsdam, East Germany, May 7-11, 1973. Part 2. Potsdam, East Germany, Deutsche Akademie der Wissenschaften, Zentralinstitut für Physik der Erde, 1974, p. 351-356. In German.

Magnetic and gravitational evidence concerning the asymmetry of the internal structure of the earth and its temporal variations are given in brief and further possibilities of the study are discussed. An even more accurate approximation of the geoidal figure can be obtained if the axes of the approximating rotation-symmetrical forms will be located outside the plane of the equator. Such an investigation of static and dynamic phenomena can provide a better understanding for some features of the material of the core. Further seismological investigations are needed for a precise location of the position of the inner core. The study of the mass movements of the core may result in a better knowledge of the tectonical processes of the earth's surface.

A75-18125 # On the determination of secular variation in the earth's gravity field and of secular polar motion. V. F. leremeev and M. I. lurkina (Tsentral'nyi Nauchno-Issledovatel'skii Institut Geodezii Aerofotos'emki i Kartografii, Moscow, USSR). In: Geodesy and physics of the earth; Proceedings of the Second International Symposium, Potsdam, East Germany, May 7-11, 1973. Part 2. Potsdam, East Germany, Deutsche Akademie der Wissenschaften, Zentralinstitut für Physik der Erde, 1974, p. 415-422. 8 refs.

The approach used in the determination of the secular variation in the earth's gravity field is based on the principles considered by Molodenskiy (1958). The problem can be solved if the components of the plane displacement of a point on the earth's surface and the elements of variation in the earth's gravity field can be expressed by means of smoothly varying functions of the plane coordinates. G.R.

A75-18128 # Use of terrestrial gravity data in combination with satellite results for determining geoid heights in Central Europe. W. Groten and R. Rummel (Darmstadt, Technische Hochschule, Darmstadt, West Germany). In: Geodesy and physics of the earth; Proceedings of the Second International Symposium, Potsdam, East Germany, May 7-11, 1973. Part 2. Potsdam, East Germany, Deutsche Akademie der Wissenschaften, Zentralinstitut für Physik der Erde, 1974, p. 441-454. 8 refs. Research supported by the Deutsche Forschungsgemeinschaft.

Possibilities of calculating the earth's local A75-18524 # gravitational field from satellite observations (Vozmozhnosti vychisteniia lokal'nogo gravitatsionnogo polia zemli po nabliudeniiam ISZ). Iu. Kh. Zhagar and N. V. Zhagar. In: Optimization of the treatment of satellite and time observations. Riga, Latviiskii Gosudarstvennyi Universitet, 1974, p. 75-83. In

Russian.

N75-10524# Informatics, Inc., Rockville, Md. SOVIET SATELLITE GEODESY: BASIS AND STATUS OF SOVIET LASER GEODESY AND LONG-BASE LINE IN-TERFEROMETRY Interim Report

Stuart G. Hibben and Eleanor M. Rowell 15 May 1974 262 p

(Contract F44620-72-C-0053)

(AD-781138/3; AFOSR-74-0962TR) Avail: NTIS HC \$6.50 **CSCL 08/5**

· Soviet developments in satellite geodesy are described in a survey of the literature, and by reports on observation and tracking stations, space triangulation networks, cameras used for satellite geodesy, laser geodesy, and long baseline interferometry. JIGRA

N75-10528*# Geological Survey, Reston, Va. OVERALL EVALUATION OF ERTS-1 IMAGERY FOR CARTOGRAPHIC APPLICATION Final Report, 1 Jul. 1972 - 30 Jun. 1974

Alden P. Covocoresses, Principal Investigator 1 Jul. 1974 69 p refs Original contains imagery. Original photography contains imagery. Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue, Sioux Falls, S. D. 57198 ERTS

(NASA Order S-70243-AG)

(E74-10800; NASA-CR-140494) Avail: NTIS HC \$4.25 CSCL 08B

The author has identified the following significant results. The conclusion reached is that an ERTS-type satellite has widespread cartographic application for scales of 1:250,000 and smaller. ERTS imagery also indicates those areas requiring revision at larger scales. For optimum cartographic application, ERTS must be flown continuously as temporal change (seasonal and long term) detection requires comparative coverage. ERTS is the first imagery system that lends itself to automated mapping wherein cartographic products may be produced in a matter of days rather than in months or years.

N75-10625# Commonwealth Scientific and Industrial Research Organization, Melbourne (Australia) Div. of Applied Geomechanics. THE PUCE PROGRAMME FOR TERRAIN EVALUATION FOR ENGINEERING PURPOSES. 2: PROCEDURES FOR TERRAIN CLASSIFICATION

K. Grant 1974 82 p refs (PB-233847/3; TP-19) Avail: NTIS HC \$7.25 CSCL 08F

Principles of terrain evaluation in accordance with the PUCE program for terrain evaluation for engineering purposes are outlined. A complete instruction manual is also provided on the art of terrain class classification conducted in accordance with the standard system incorporated as an integral part of the PUCE program. The manual includes as appendices the definitions of and numerical system for nomenclature of the terrain classes. class intervals for terrain parameters, and the Unified Soil Classification scheme. Also included as appendices are examples of standard terrain class description sheets, a glossary of standard terms to be used in terrain classification, and examples extracted from the complete classification of one area in Australia.

Author (GRA)

N75-10641# Woods Hole Oceanographic Institution, Mass. CRUISE DATA REPORT. CHAIN 115 LEG 2, SOUTHLANT **EXPEDITION**

D. A. Richards, R. C. Groman, and S. R. Gegg Aug. 1974 25 p

(Contract N00014-66-C-0241; NR Proj. 083-004) (AD-784421; WHOI-74-59) Avail: NTIS CSCL 08/10

The report summarizes the geological and geophysical activities which were undertaken during Leg 2 (Dakar to Cape Town) of R/V CHAIN Cruise 115 (SOUTHLANT Expedition). A series of charts and profiles display the navigational, bathymetry, total field magnetic anomaly, and free air gravity anomaly measurements, as well as the locations of coring, camera and hydrographic stations. Author (GRA)

N75-10869# Joint Publications Research Service, Arlington, Va.

STATUS AND PROSPECTS OF DEVELOPMENT OF SPACE **GEODESY**

A. A. Izotov 18 Oct. 1974 16 p Transl. into ENGLISH from Geod. i Kartografiya (Moscow), no. 5, 1974 p 10-18 (JPRS-63231) Avail: NTIS HC \$3.25

The developments in space geodesy for studying the earth Author and circumterrestrial space are reviewed.

N75-12395*# lowa Univ., lowa City. Dept. of Geology. EXPERIMENT TO EVALUATE FEASIBILITY OF UTILIZING SKYLAB-EREP REMOTE SENSING DATA OF TECTONIC ANALYSIS OF THE BIGHORN MOUNTAINS REGION, WYOMING-MONTANA Quarterly Progress Report, 1 Jul. - 30 Sep. 1974

Richard A. Hoppin, Principal Investigator 20 Oct. 1974 6 p

(Contract NAS9-13313)

(E75-10014; NASA-CR-140650) Avail: NTIS HC \$3.25 CSCL 08G

N75-12413*# Geological Survey, Denver, Colo.
REMOTE SENSING GEOPHYSICS FROM SKYLAB Monthly
Report, Jun. 1974
Kenneth Watson, Principal Investigator and H. A. Pohn Jun.
1974 2 p EREP
(NASA Order T-6555-B)
(E75-10045; NASA-CR-140782) Avail: NTIS HC \$3.25 CSCL
08G

N75-13338*# California Earth Science Corp., Santa Monica. FAULT TECTONICS AND EARTHQUAKE HAZARDS IN THE PENINSULAR RANGES, SOUTHERN CALIFORNIA Monthly Progress Report, Sep. 1974
Paul M. Merifield and Donald L. Lamar, Principal Investigators

Paul M. Merifield and Donald L. Lamar, Principal Investigators 5 Oct. 1974 4 p. EREP

(Contract NAS2-7698)

(E75-10022; NASA-CR-140736; MPR-16) Avail: NTIS HC \$3.25 CSCL 08G

The author has identified the following significant results. ERTS and Skylab images reveal a number of prominent lineaments in the basement terrane of the Peninsular Ranges, Southern California. The major, well-known, active, northwest trending, right-slip faults are well displayed, but northeast and west to west-northwest trending lineaments are also present. Study of large-scale airphotos followed by field investigations have shown that several of these lineaments represent previously unmapped faults. Pitches of striations on shear surfaces of the northeast and west trending faults indicate oblique-slip movement; data are insufficient to determine the net-slip. These faults are restricted to the pre-Tertiary basement terrane and are truncated by the major northwest trending faults; therefore, they may have formed in response to an earlier stress system. Future work should be directed toward determining whether the northeast and west trending faults are related to the presently active stress system or to an older inactive system, because this question relates to the earthquake risk in the vicinity of these faults.

N75-13340*# Smithsonian Astrophysical Observatory, Cambridge, Mass.

MAPPING OF THE MAJOR STRUCTURES OF THE AFRICAN RIFT SYSTEM

P. A. Mohr, Principal Investigator $\,$ 29 Oct. 1974 $\,$ 90 p $\,$ refs ERTS $\,$

(Contract NAS5-21748)

(E75-10024; NASA-CR-140738; SAO-Special-Rept-361;

SAO-403-004) Avail: NTIS HC \$4.75 CSCL 08B

The author has identified the following significant results. The ERTS-1 satellite imagery has facilitated a major advance in accurate mapping and better understanding of the African Rift valleys. A unified scheme of mapping of the whole rift system, from Malawi to Ethiopia, has been accomplished. The structures revealed by the imagery are discussed in the light of known ground truth for the northern half of the African Rift System. The ERTS-revealed structures confirm the likelihood of a drift of Arabia away from Africa and impose constraints on the drift vectors. Numerous features have been newly recognized from

the imagery: the gently curvilinear plan of virtually all the African Rift valleys, the pervasive but not overriding degree of influence of Precambrian structures on rift faulting, the deep structural control of many rift volcanic centers, the occurrence of several unsuspected calderas both in the rifts and on the plateaus, and massive cauldron subsidence phenomena previously unknown to eastern Africa. Lithological mapping in eastern Africa can be greatly refined using the ERTS imagery, and new or revised subdivisions of the Precambrian appear possible and, in some places, necessary. Valleys excavated in the Ethiopian highlands by Pleistocene glaciers are newly recognized.

N75-13352*# Geological Survey, Denver, Coto.
REMOTE SENSING GEOPHYSICS FROM SKYLAB Monthly
Report, Sep. 1974
Kenneth Watson, Principal Investigator and H. A. Pohn Sep.
1974 3 p EREP
(NASA Order T-6555-B)
(E75-10052; NASA-CR-140954) Avail: NTIS HC \$3.25 CSCL
08G

N75-13359*# Naval Research Lab., Washington, D.C. TERRAIN PROPERTIES AND TOPOGRAPHY FROM SKYLAB ALTIMETRY Monthly Progress Report, Oct. 1974
Allan Shapiro, Principal Investigator 6 Dec. 1974 1 p EREP (NASA Order T-4716-B) (E75-10059; NASA-CR-140999) Avail: NTIS HC \$3.25 CSCL 08F

N75-13364*# Battelle Columbus Labs., Ohio.
CALIBRATION AND EVALUATION OF SKYLAB ALTIMETRY
FOR GEODETIC DETERMINATION OF THE GEOID Progress
Report, 1 Sep. - 30 Nov. 1974
A. G. Mourad, Principal Investigator, S. Gopalapillai, and M. Kuhner
12 Dec. 1974 18 p EREP
(Contract NAS9-13276)
(E75-10064; NASA-CR-141004; PR-19) Avail: NTIS
HC \$3.25 CSCL 08E

N75-13375# Army Engineer Waterways Experiment Station, Vicksburg, Miss.

AUTOMATED SELECTION AND LOCATION OF TERRAIN FEATURES ON THE BASIS OF SPECTRAL CHARACTERISTICS

Albert N. Williamson, Jr. 26 Aug. 1974 16 p (AD-785698) Avail: NTIS CSCL 08/2

The report details data processing methods for converting multispectral scanner images, transmitted from a satellite, by application of computer techniques.

N75-13432# Geophysical Observatory, Sodankyla (Finland). GEOPHYSICAL VARIATIONS AT SODANKYLA DURING A GEOMAGNETIC STORM, 17-18 DECEMBER 1971
Tauno Turunen and M. Mukunda Rao May 1974 21 p refs

(Rept-9) Avail: NTIS HC \$3.25

Extremely bright auroras could be seen at Sodankyla and throughout Scandinavia on the night of 17 - 18 December 1971. A detailed study was undertaken of the geophysical phenomena observed at Sodankyla during this storm. The all-sky camera records, magnetograms, riometer records, and ionograms obtained at Sodankyla were analyzed in detail and published data from neighboring stations were examined to elucidate the extent of the geomagnetic storm and its effects on the upper atmosphere. It was noted that the occurrence of active auroral forms overhead was invariably followed by an increase of riometer absorption, and by changes in sporadic E. The analysis of this storm indicates that at least three substorms occurred during the main phase of the storm.

N75-13454# Bureau of Mineral Resources, Geology and Geophysics, Canberra (Australia).

AUSTRALIAN AND SOVIET GRAVITY SURVEYS ALONG THE AUSTRALIAN CALIBRATION LINE

P. Wellman, Yu. O. Boulanger, B. C. Barlow, S. N. Scheglov, and D. A. Coutts 1974 177 p refs

(Bull-161; ISBN-0-642-00853-1) Avail: NTIS HC \$7.00

The Australian calibration line (ACL), with a total gravity interval of 3 Gal, was established in 1970 between Laiagam in Papua New Guinea and Hobart in Tasmania, and during 1973 the Australian Bureau of Mineral Resources and the U.S.S.R. Geodesy and Cartography Survey made joint observations along the full length of the ACL. Measurements made with eight Soviet GAG-2 gravity meters established a gravity scale for Australia to an accuracy of 2.5 parts in 10 to the 5th power. This scale and a datum of 979671.86 mGal for Sydney A were adopted for Australia in 1973. The complete data are shown in tabular form.

N75-13498# Joint Publications Research Service, Arlington,

ANALYSIS OF METHODS FOR THE GEODETIC SUPPORT OF COMPLEX SEA GEOPHYSICAL SURVEYS

I. F. Glumov, V. P. Glumov, and A. V. Smirnov 10 Dec. 1974 17 p refs Transl. into ENGLISH from Izv. Vyshikh Uchebn. Zavedenii, Geod. i Aerofotosemka (Moscow), no. 3, 1974 p 17-26

(JPRS-63621) Avail: NTIS HC \$3.25

Methods for determining position at sea and the range of applicability of each method in complex sea geophysical surveys are analyzed.

Author

N75-14193*# California Earth Science Corp., Santa Monica. FAULT TECTONICS AND EARTHQUAKE HAZARDS IN THE PENINSULAR RANGES, SOUTHERN CALIFORNIA Monthly Progress Report, Nov. 1974

Paul M. Merifield, Principal Investigator 5 Dec. 1974 2 p

(Contract NAS2-7698)

(E75-10074: NASA-CR-141097: MPR-18) Avail: NTIS HC \$3.25 CSCL 08E

N75-14197*# Geological Survey, Reston, Va.
CARTOGRAPHIC EVALUATION OF SKYLAB-A S-192
SCANNER IMAGES Quarterly Progress Report, 1 Aug. 31 Oct. 1974

John D. McLaurin, Principal Investigator 31 Oct. 1974 3 p EREP

(NASA Order T-4111-B)

(E75-10078; NASA-CR-141093) Avail: NTIS HC \$3.25 CSCL ORB

N75-14235 Hunting Surveys, Ltd., Boreham Wood (England).
NATURAL-RESOURCES MAPPING OF ETHIOPIA FROM
ERTS-1 IMAGERY

L. P. White *In* ESRO European Earth-Resources Satellite Expts. May 1974 p 179-184 Original contains color illustrations

The complete ERTS coverage of Ethiopia was used to examine its potential for mapping a variety of natural resources including soils and landforms, vegetation, and land use. The high planimetric accuracy of the imagery has proved invaluable in map compilation and in a number of cases major discrepancies were found in geographical detail between the imagery and the 1,000,000 base maps used for reference. The study confirms the experience of many investigators that the ERTS imagery can be used for a range of natural resources mapping to reconnaissance level, but the supplementary ground information necessary varies considera-

bly for each application. As well as the basic resource mapping capability, the sequential cover allows monitoring of dynamic situations, even of very subtle expression.

Author (ESRO)

N75-14239 Technische Universitaet, Clausthal-Zellerfeld (West Germany).

ERTS-1 DATA ON AFAR TECTONICS

P. Kronberg, M. Schoenfeld, R. Guenther, P. Tsombos, and D. Bannert (Federal Inst. for Earth Res., Hanover) In ESRO European Earth Resources Satellite Expts. May 1974 p 217-229 :refs Original contains color illustrations i

(Grant WRK-213)

The Afar Triangle (Ethiopia), where the rift systems of the Red Sea and Gulf of Aden seem to meet, was one of the key areas for recent investigations of crustal-spreading problems. The evaluation of ERTS images led to the first detailed structural map covering the whole of the Afar and the adjacent areas. A lithological map was also compiled from ERTS data, covering the Afar and its surrounding areas. Correlations of ERTS data with data from photogeological mapping and from geological, petrological, and geophysical field observations were made. The evaluation of the ERTS-1 images has highlighted new aspects of the tectonics of the Afar and its geological history.

Author (ESRO)

N75-14261# Environmental Research Inst. of Michigan, Ann Arbor.

THE REMOTE IDENTIFICATION OF TERRAIN FEATURES AND MATERIALS AT A CALIFORNIA TEST SITE: AN INVESTIGATIVE STUDY OF TECHNIQUES Interim Report, Sep. 1972 - Dec. 1973

R. K. Vincent, R. D. Dillman, and P. G. Hasell, Jr. Apr. 1974 64 p refs

(Contract FH-11-7136)

(PB-235991/7; ERIM-196200-11-T; FHWA-RD-74-27) Avail: NTIS HC \$4.25 CSCL 08M

The collection, processing; and analysis of multispectral imagery gathered in airborne mapping of a site in the vicinity of Halloran Springs, California in the fall of 1972 are discussed. The general purpose is to develop techniques for automatically identifying selected terrain features and natural materials in remotely sensed imagery. Objectives were to automatically classify both soils of different textures or parent materials and rock outcrops based on (1) their spectral reflectance characteristics at ultraviolet, visible, and infrared wavelengths, and (2) their emissivity differences as noted in multiple thermal bands. The first objective was achieved; the second was not, primarily because of the poor performance of an experimental multiband thermal-wavelength detector.

N75-14263# Environmental Research Inst. of Michigan, Ann Arbor.

THE REMOTE IDENTIFICATION OF TERRAIN FEATURES AND MATERIALS AT PENNSYLVANIA TEST SITES: AN INVESTIGATIVE STUDY OF TECHNIQUES Interim Report, Apr. 1969 - Dec. 1972

Franklin G. Sadowski, Thomas W. Wagner, Philip G. Hasell, Jr., and Frederick J. Thomson Mar. 1974 93 p refs

(Contract FH-11-7136)

(PB-235992/5; ERIM-196200-5-T; FHWA-RD-74-9) Avail: NTIS HC \$4.75 CSCL 08M

The collection, processing, and analysis of multispectral imagery gathered in airborne mapping is reported. The specific objective was to automatically classify soils of different parent materials by surface spectral reflectance and emittance characteristics. Processing techniques included multispectral statistical pattern recognition, ratio processing, and single channel processing. Analog and digital procedures were utilized. The discrimination of bare soils was accomplished, but classification of soils according to parent material yielded unpromising results. An average spectral difference was noted between limestone-derived soils and sandstone- and shale-derived soils in the 11 channel visible and

near-IR spectral region. An apparent capability to improve the classification of soils despite albedo variation was demonstrated by using the ratio of five channels from the same region. Scan-angle variations indicated sufficient differences in the bidirectional reflectance characteristics to warrent correction of the data prior to processing. (Modified author abstract) GRA

N75-14335# Ohio State Univ., Columbus. Dept. of Geodetic Science.

SOME FIRST ACCURACY ESTIMATES FOR APPLICATIONS OF AERIAL GRADIOMETRY

Helmut Moritz Jul. 1974 78 p refs

(Contract F19628-72-C-0120; AF Proj. 8607)

(AD-786418; DGS-209; AFCRL-TR-74-0317; SR-18)

NTIS CSCL 08/5

Estimates of accuracies obtainable in the geodetic use of ... aerial gradiometry are given. For simplicity, mainly the second vertical gradient T sub zz is considered; the use of other gradients and their optimal combination by least-squares collocation is also outlined. GRA .

N75-15141# Center for Naval Analyses, Arlington, Va. Warfare Analysis Group

THE GLOBAL COMPUTER PROGRAM AND GLOBEMAP: A SPECIAL PURPOSE THREE-DIMENSIONAL PLOTTER SUBROUTINE

Lewis W. Linton Jul. 1974 45 p

(Contract N00014-68-A-0091)

(AD-785869; NWAG-Research-Contrib-266) Avail: NTIS CSCL 08/5

GLOBALL manipulates several smaller computer programs to calculate global ranges and surface areas. GLOBEMAP provides a three-dimensional map background upon which the ranges and areas may be presented. This paper is not intended to be a user's manual but to describe characteristics of these models Author (GRA) and possible applications of them.

N75-15143# Army Foreign Science and Technology Center. Charlottesville, Va.

THE NEXT PROBLEMS IN THE AREA OF THE IMPLEMEN-TATION OF PLANNING FOR CARTOGRAPHICAL-GEODETIC WORK

L. A. Kashin 21 Feb. 1974 12 p Transl. into ENGLISH from Geodeziya i Kartografiya (Moscow), no. 7, 1972 p 1-5 (AD-785882; FSTC-HT-23-1558-73) Avail: NTIS CSCL 08/5

Geodesy and cartography in the Soviet Union are discussed. A brief history of the improvements and changes over the past few years is given.

N75-15147# Defense Mapping Agency Aerospace Center, St.

FULLY AUTOMATED COMPILATION OF GRAVITY AND GEOPHYSICAL MAPS

Luman E. Wilcox, Joe M. Estes, and Walter Czarnecki Sep. 1974 69 p refs

(AD-785838; DMAAC-TP-74-002) Avail: NTIS CSCL 08/2 A wide variety of gravity and geophysical maps can be produced by fully automated methods. The locations of gravity measurements, gravity anomaly values, gravity anomaly contours, continental outlines, title, legend, labels, and other data to be shown on the maps are computer generated, then drawn by automatic plotting equipment. This procedure is used to prepare charts showing the density and distribution of gravity data holdings. contoured gravity anomaly maps and other graphics needed for gravity data evaluation and mean anomaly prediction purposes, maps of gravity station elevations used as vertical control intopographic map compilation, and maps of glacial ice thickness for geophysical analysis. (Modified author abstract)

N75-15228# Kakioka Magnetic Observatory (Japan). REPORT OF THE GEOMAGNETIC AND GEOELECTRIC **OBSERVATIONS, 1972 (RAPID VARIATIONS)** 1973 76 p

(Rept-13) Avail: NTIS HC \$4.75

Results of observations of the Post-IGC, carried out by the Kakioka Magnetic Observatory in close cooperation with its two branch magnetic observatories, are presented. Standardization and comparison services of absolute magnetic instruments are operated by the electro-magnetic standard magnetometer A-56 and the proton procession magnetometer MO-P. Results of observations of rapid variations and records of the rapid-run induction magnetograms and rapid-run tellurigrams are included. Author

N75-15237# Army Foreign Science and Technology Center, Charlottesville. Va.

OPTICAL QUANTUM GENERATORS IN GEODETIC MEAS-UREMENTS

M. T. Prilepin and A. N. Golubev 1974 182 p. refs Transl. into ENGLISH of the book "Opticheskie Kvantovye Generatory v Geodezicheskikh Izmereniyakh" Moscow, 1972 168 p (AD-787281; FSTC-HT-23-1170-73) Avail: NTIS CSCL 08/5

The book considers the theoretical and practical problems of the application of optical quantum generators in contemporary geodetic measurements. It also offers an analysis of completed experiments and describes laser geodetic apparatus and measuring systems and methods. The book is made up of four chapters. The first chapter will familiarize the reader with certain problems connected with the application of optical quantum generators. The second chapter is devoted to the use of lasers in terrestrial range finder systems. Pulse, phase, and interference methods of measuring distances using optical quantum generators are discussed. The third chapter discusses laser systems for measuring large distances (to the earth's artificial satellites and to the moon) and gives an analysis of the accuracy of such systems and describe experiments which have been conducted. In the fourth chapter other geodetic applications of optical quantum generators are discussed. GRA

04 GEOLOGY AND MINERAL RESOURCES

Includes mineral deposits, petroleum deposits, spectral properties of rocks, geological exploration, and lithology.

A75-10437 Oil exploration needs for digital processing of imagery. F. F. Sabins, Jr. (Chevron Oil Field Research Co., La Habra, Calif.). (American Society of Photogrammetry, Annual Convention, St. Louis, Mo., Mar. 11, 1974.) Photogrammetric Engineering, vol. 40, Oct. 1974, p. 1197-1200.

Survey of the applications of digital image processing in the oil industry. The four major types of imaging sensors employed in oil exploration are aerial photography, thermal infrared, side-looking radar, and Earth Resource Technology Satellite (ERTS). ERTS imagery, because of its availability in a digital format and its use in reconnaissance studies, is shown to have the greatest potential in future digital processing. Well suited for exploration reconnaissance, ERTS imagery is characterized by broad regional coverage, acceptable resolution, and minimum image distortion. A discussion of the objectives and requirements of future image processing is included.

A75-11068 * A simulation of the San Andreas fault experiment. R. W. Agreen and D. E. Smith (NASA, Goddard Space Flight Center, Greenbelt, Md.). *Journal of Geophysical Research*, vol. 79, Oct. 10, 1974, p. 4413-4417. 9 refs.

The San Andreas fault experiment (Safe), which employs two laser tracking systems for measuring the relative motion of two points on opposite sides of the fault, has been simulated for an 8-yr observation period. The two tracking stations are located near San Diego on the western side of the fault and near Quincy on the eastern side; they are roughly 900 km apart. Both will simultaneously track laser reflector equipped satellites as they pass near the stations. Tracking of the Beacon Explorer C spacecraft has been simulated for these two stations during August and September for 8 consecutive years. An error analysis of the recovery of the relative location of Quincy from the data has been made, allowing for model errors in the mass of the earth, the gravity field, solar radiation pressure, atmospheric drag, errors in the position of the San Diego site, and biases and noise in the laser systems. The results of this simulation indicate that the distance of Quincy from San Diego will be determined each year with a precision of about 10 cm. Projected improvements in these model parameters and in the laser systems over the next few years will bring the precision to about 1-2 cm by 1980. (Author)

A75-17108 # Geologic interpretation of ERTS-1 satellite images for East Aswan area, Egypt. E. M. El Shazly (Academy of Scientific Research and Technology, Cairo, Egypt), M. A. Abdel-Hady (Oklahoma State University, Stillwater, Okla.; Academy of Scientific Research and Technology, Cairo, Egypt), M. A. El Ghawaby, and I. A. El Kassas (Atomic Energy Establishment, Cairo, Egypt). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 105-117. 12 refs. Research supported by Oklahoma State University and NSF.

A75-17109 # Geologic interpretation of ERTS-1 satellite images for West Aswan area, Egypt. E. M. El Shazly (Academy of Scientific Research and Technology, Cairo, Egypt), M. A. Abdel-Hady (Oklahoma State University, Stillwater, Okla.; Academy of Scientific Research and Technology, Cairo, Egypt), M. A. El Ghawaby, and I. A. El Kassas (Atomic Energy Establishment, Cairo,

Egypt). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings, Volume 1.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 119-131. 6 refs. Research supported by Oklahoma State University and NSF.

A75-17126 # Development of research in the USSR on the use of radar images for geological purposes. V. B. Komarov, V. A. Starostin, and B. P. Niavro (Ministry of Geology of the USSR, Leningrad; Akademiia Nauk SSSR, Moscow, USSR). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1.

Ann Arbor, Mich., Environmental Research Institute

of Michigan, 1974, p. 441-444.

Side-looking airborne radar is widely used in the USSR for geological surveys. The radar images obtained are influenced by physical properties of the reflecting surface, such as density, and seasonal factors such as moisture content, vegetation, and snow cover. Correlation analysis is used to determine the dependence of the radar image on the surface characteristics. Correlation relationships established so far indicate that it is possible to determine quantitatively the moisture content and the density of loose dry-phase deposits from radar surveys.

A.T.S.

A75-17207 # Crude and refined petroleum oil structured luminescence signatures induced by UV laser or lamp and their remote sensing applications. H. G. Gross and M. Muramoto (McDonnell Douglas Astronautics Co., Huntington Beach, Calif.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1747-1759, 6 refs. Research supported by the McDonnell Douglas Astronautics Independent Research and Development Funds.

A75-17215 # Geologic interpretation of infrared thermal images in East Qatrani area, Western Desert, Egypt. E. M. El-Shazly (Academy of Scientific Research and Technology, Cairo, Egypt), M. A. Abdel-Hady (Oklahoma State University, Stillwater, Okla.; Academy of Scientific Research and Technology, Cairo, Egypt), and M. A. Morsy (Atomic Energy Establishment, Cairo, Egypt). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1877-1889. 13 refs. Research supported by the Oklahoma State University and NSF.

A75-17220 # Remote sensing of rock type in the visible and near-infrared. J. W. Salisbury and G. R. Hunt (USAF, Terrestrial Sciences Leboratory, Bedford, Mass.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1953-1958. 9 refs.

Visible and near-infrared spectra of minerals and rocks have been measured and evaluated in terms of remote sensing applications. We conclude that there are some differences in the spectral behavior of different rock types in the visible and near-infrared. These differences are, however, difficult or impossible to utilize in a generalized remote sensing effort in which the composition of all rocks is to be mapped. Instead, this spectral region lends itself best to precise and particular applications, such as enhancing the visibility of a rock unit with a known and distinctive spectral signature, or enhancing the contrast between rock units, or between rocks and the vegetative background. (Author)

A75-17222 # Application of radar imagery to environmental geologic mapping of Texas. P. J. Cannon (Texas, University, Austin, Tex.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings.

Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1981-1988.

A75-17225 # Unsupervised mapping of geologic features and soils in California. R. D. Dillman and R. K. Vincent (Michigan, Environmental Research Institute, Ann Arbor, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 2013-2025. 10 refs.

This paper describes a photographic and computer implemented method of analyzing and mapping the distribution of geologic materials at a test site near Halloran Springs, in Southern California. Laboratory spectral curves of selected rocks and minerals were used to calculate the ratio of reflectances using six bands, including two thermal bands. These laboratory ratio values were used to predict characteristic colors to be found on a color composite of three ratio images and also to train an analogue computer for automatic scene classification. In comparison with a geologic map prepared by ground survey, correct classification was as high as 77.9% using the ratio color composite. (Author)

A75-17226 # Remote sensing to detect the toxic effects of metals on vegetation for mineral exploration. N. P. Press (Nigel Press Associates, London, England). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 2027-2038. 26 refs. Research supported by the American Smelting and Refining Co., Barringer Research, Ltd., and Rio Tinto Zinc Corp.

A75-17229 # The use of space photos for search of oil and gas fields. P. V. Florenskii (Akademiia Nauk SSSR, Geologicheskii Institut; Moskovskii Institut Neftekhimicheskoi i Gazovoi Promyshlennosti, Moscow, USSR). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 2055-2058. 7 refs.

A75-18065 Automatic mapping of strip mine operations from spacecraft data. R. H. Rogers, L. E. Reed (Bendix Corp., Aerospace Systems Div., Ann Arbor, Mich.), and W. A. Pettyjohn (Ohio State University, Columbus, Ohio). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings.

Falls Church, Va., American Society of Photogrammetry, 1974, p. 204-219. 5 refs.

Computer techniques were applied to process ERTS tapes acquired over coal mining operations in southeastern Ohio on Aug. 21, 1972 and Sept. 3, 1973. ERTS products obtained included geometrically correct map overlays showing stripped earth, partially reclaimed earth, water, and natural vegetation. Computer generated tables listing the area covered by each land-water category in square kilometers and acres were also produced. By comparing these mapping products, the study demonstrates the capability of ERTS to monitor changes in the extent of stripping, success of reclamation, and the secondary effects of mining on the environment. (Author)

N75-10530*# Colorado School of Mines, Golden. Dept. of Geology.

GEOLOGIC AND MINERAL AND WATER RESOURCES INVESTIGATIONS IN WESTERN COLORADO, USING SKYLAB EREP DATA Monthly Progress Report, Sep. 1974 Keenan Lee, Principal Investigator Sep. 1974 4 p EREP (Contract NAS9-13394)

(E74-10802; NASA-CR-140496) Avail: NTIS HC \$3.25 CSCL 08G

N75-10536*# New York State Museum and Science Service.

ASSESSMENT OF ERTS-1 IMAGERY AS A TOOL FOR REGIONAL GEOLOGICAL ANALYSIS IN NEW YORK STATE Final Report

Yngvar W. Isachsen, Principal Investigator, Robert H. Fakundiny, and Stephen W. Forster 28 May 1974 181 p refs Original contains imagery. Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue, Sioux Falls, S. D. 57198 ERTS

(Contract NAS5-21764)

(E74-10809; NASA-CR-140503) Avail: NTIS HC \$7.00 CSCL 08G

The author has identified the following significant results. Linear anomalies dominate the new geological information derived from ERTS-1 imagery, total lengths now exceeding 26,500 km. Maxima on rose diagrams for ERTS-1 anomalies correspond well with those for mapped faults and topographic lineaments. Multi-scale analysis of linears shows that single topographic linears at 1:2,500,000 may become dashed linears at 1:1,000,000 aligned zones of shorter parallel, en echelon, or conjugate linears at 1:5000,000, and shorter linears lacking any conspicuous zonal alignment at 1:250,000. Field work in the Catskills suggests that the prominent new NNE lineaments may be surface manifestations of dip slip faulting in the basement, and that it may become possible to map major joint sets over extensive plateau regions directly on the imagery. Most circular features found were explained away by U-2 airfoto analysis but several remain as anomalies. Visible glacial features include individual drumlins, drumlinoids, eskers, ice-marginal drainage channels, glacial lake shorelines, sand plains, and end moraines.

N75-10537*# Tennessee Univ., Knoxville. Dept. of Geography.

GEOGRAPHIC APPLICATIONS OF ERTS-1 IMAGERY TO LANDSCAPE CHANGE Final Report

John B. Rehder, Principal Investigator Dec. 1973 112 p refs Original contains color imagery. Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue, Sioux Falls, S. D. 57198 ERTS (Contract NASS-21726)

(E74-10810; NASA-CR-140504) Avail: NTIS HC \$5.25 CSCL

The author has identified the following significant results. ERTS-1 has proven to be an effective earth-orbiting monitor of landscape change. Its regional coverage for large areal monitoring has been effective for the detection and mapping of agricultural plowing regions, for general forest cover mapping, for flood mapping, for strip mine mapping, and for short-lived precipitation mapping patterns. Paramount to the entire study has been the temporal coverage provided by ERTS. Without the cyclic coverage on an 18 day basis, temporac coverage would have been inadequate for the detection and mapping of strip mining landscape change, the analysis of agricultural landscape change based on plowing patterns, the analysis of urban-suburban growth changes, and the mapping of the Mississippi River floods. Cost benefits from ERTS are unquestionably superior to aircraft systems in regard to large regional coverage and cyclic temporal parameters. For the analysis of landscape change in large regions such as statewide areas or even areas of 10,000 square miles, ERTS is of cost benefit consideration. Not only does the cost of imagery favor ERTS but the reduction of man-hours using ERTS has been in the magnitude of 1:10.

N75-10552* Geological Survey, Atlanta, Ga.
GEOLOGIC EVALUATION AND APPLICATIONS OF ERTS-1
IMAGERY OVER GEORGIA

S. M. Pickering and R. C. Jones In NASA. Goddard Space Flight Center Third ERTS Symp., Vol. 2 May 1974 p 41-49

CSCL 08G

Satellite imagery and other remote sensing tools and techniques have provided a powerful tool to assist geologic research; significantly increased the mapping efficiency of field geologists; shown new lineaments associated with known shear and fault zones; delineated new structural features; provided a tool to reevaluate tectonic history; helped to locate potential ground-water sources and areas of aquifer recharge; defined areas of geologic hazards; shown areas of heavy siltation in major reservoirs; and, by close interval repetition, aided in monitoring surface mine reclamation activities and the environmental protection of the intricate marshland system. The Georgia Geological Survey has been engaged in regional mapping for the new state geologic map. ERTS-1 images enlarged to compatible mapping scales have increased field geologic mapping efficiency by at least 25%. There are a number of areas where data from ERTS-1 imagery has allowed a notably higher level of precision than has been available with any amount of field work on the ground.

N75-10553* Earth Satellite Corp., Washington, D.C.
AN EVALUATION OF THE SUITABILITY OF ERTS DATA FOR THE PURPOSES OF PETROLEUM EXPLORATION John R. Everett and Gerald Petzel (Eason Oil Co.) In NASA. Goddard Space Flight Center Third ERTS Symp., Vol. 2 May 1974 p 50-61 CSCL 08G

This investigation was undertaken to determine the types and amounts of information valuable to petroleum exploration that are extractable from ERTS data and to determine the cost of obtaining the information from ERTS relative to costs using traditional or conventional means. In particular, it was desirable to evaluate this new petroleum exploration tool in a geologically well-known area in order to assess its potential usefulness in an unknown area. In light of the current energy situation, it is felt that such an evacuation is important in order to best utilize technical efforts with customary exploration tools, by rapidly focusing attention on the most promising areas in order to reduce the time required to go through the exploration cycle and to maximize cost savings. The Anadarko Basin lies in western Oklahoma and the panhandle of Texas (Figure 1). It was chosen as a test site because there is a great deal of published information available on the surface and subsurface geology of the area, there are many known structures that act as traps for hydrocarbons, and it is similar to several other large epicontinental sedimentary basins.

N75-10556* Ohio State Univ., Columbus. AUTOMATED STRIP MINE AND RECLAMATION MAPPING FROM ERTS

Wayne A. Pettyjohn, Robert H. Rogers (Bendix Corp., Ann Arbor, Mich.), and Larry E. Reed (Bendix Corp., Ann Arbor, Mich.) In NASA. Goddard Space Flight Center Third ERTS Symp., Vol. 2 May 1974 p 87-101 CSCL 081

In response to the urgent need for a faster and more economical means of generating strip mine and reclamation maps, a study was conducted to evaluate the suitability of using ERTS computer compatible tape for automatic mapping. The procedure uses computer target spectral recognition techniques as a basis for classification. The area encompassed by this investigation includes five counties in eastern Ohio that comprise nearly 7,500 square kcometers (3,000 square miles). The counties have been disrupted by coal mining since the early 1800's, and strip mining has been practiced in all of them. The environmental effects of strip mining are also discussed. A.L.

N75-10561* National Aeronautics and Space Administration. Goddard Space Flight Center, Greenbelt, Md. MINERAL RESOURCES, GEOLOGICAL STRUCTURES, AND LANDFORM SURVEYS

p 147-167 refs

Nicholas M. Short In its Third ERTS Symp., Vol. 2 May 1974 CSCL 08E

Since March 1973 there has been a shift in ERTS results in geology from the initial show-and-tell stage to a period in which scientific studies predominated, and now to an emphasis on effective applications having economic benefits and clearcut relevance to national needs. Many years will be spent on geological tasks resulting from ERTS alone; reconnaissance mapping in inaccessible regions, map revisions, regional or synoptic analysis of crustal fractures, assessment of dynamic surficial processes, systematic search for mineral wealth, use of sophisticated enhancement techniques, recognition of potentia: geologic hazards, and many more applications that still need to be defined.

Author

N75-10887*# Scientific Translation Service, Santa Barbara, Calif. ORBITAL GEOLOGY

V. Brukhanov Washington NASA Oct. 1974 5 p Transl. into English from Izv. (Moscow), no. 173, 25 Jul. 1974 p. 2 (Contract NASw-2483)

(NASA-TT-F-16000) Avail: NTIS HC \$3.25 CSCL 03B

The use of satellites in space to determine geological features on the earth is analyzed and discussed with emphasis on geological advances in space technology.

N75-11407*# California Earth Science Corp., Santa Monica. LINEAMENTS IN BASEMENT TERRANE OF THE PENIN-SULAR RANGES, SOUTHERN CALIFORNIA

P. M. Merifield, Principal Investigator and D. L. Lamar 27 Sep,_ 1974 26 p refs Orginal contains imagery Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue, Sioux Falls, S. D. 57198 EREP

(Contracts NAS2-7698; DI-14-08-0001-13911)

(E75-10003; NASA-CR-140725; TR-74-1) Avail: NTIS HC \$3.75 CSCL 08G

The author has identified the following significant results. ERTS and Skylab images reveal a number of prominent lineaments in the basement terrane of the Peninsular Ranges, Southern California. The major, well-known, active, northwest trending, right-slip faults are well displayed; northeast and west to west-northwest trending lineaments are also present. Study of large-scale airphotos followed by field investigations have shown that several of these lineaments represent previously unmapped faults. Pitches of striations on shear surfaces of the northeast and west trending faults indicate oblique slip movement; data are insufficient to determine the net-slip. These faults are restricted to the pre-tertiary basement terrane and are truncated by the major northwest trending faults. They may have been formed in response to an earlier stress system. All lineaments observed in the space photography are not due to faulting, and additional detailed geologic investigations are required to: determine the nature of the unstudied lineaments, and the history and net-slip of fault-controlled lineaments.

N75-11408*# California Earth Science Corp., Santa Monica. FAULT TECTONICS AND EARTHQUAKE HAZARDS IN THE PENINSULAR RANGES, SOUTHERN CALIFORNIA Montly Progress Report, Nov. 1974

Paul M. Merifield, Principal Investigator 5 Nov. 1974 2 p EREP

(Contract NAS2-7698)

(E75-10005; NASA-CR-140727; MPR-17) Avail: NTIS HC \$3.25 CSCL 08G

N75-11436* Scientific Translation Service, Santa Barbara, Calif. GEOLOGICAL AND GEOPHYSICAL STUDIES FROM SPACE AND PROSPECTS FOR THEIR DEVELOPMENT

G. B. Gonin, A. I. Vinogradova, B. V. Shilin, and N. A. Yakovlev In its Space Iconics (NASA-TT-F-798) Sep. 1974 p 174-181 Transl. into ENGLISH of the publ. "Kosmicheskaya Ikonika" Moscow, Nauka Press, 1973 p 147-152

CSCL 08G

04 GEOLOGY AND MINERAL RESOURCES

The possibilities of using images obtained by devices which detect the reflected and self-radiation of the the earth's surface and atmosphere in other parts of the spectrum, other than visible, were analyzed. Considered were: (1) infrared; (2) radiothermal; and (3) magnetic field.

N75-11437* Scientific Translation Service, Santa Barbara, Calif. EXPERIENCE IN INTEGRATED GEOLOGICAL AND GEOGRAPHICAL INTERPRETATION OF SPACE PHOTOGRAPHS OBTAINED IN THE USSR

B. V. Vinogradov and A. A. Grigoryev *In its* Space Iconics (NASA-TT-F-798) Sep. 1974 p 182-189 refs Transl into ENGLISH of the publ. "Kosmicheskaya Ikonika" Moscow, Nauka Press, 1973 p 153-158 CSCL 14E

Two examples of regional interpretation of imagery acquired from satellite photographic and television systems are discussed. The imagery was taken aboard the Zond 7 and Kosmos satellites, and depicted: (1) sand and dust storm activity in the eastern Mediterranean Sea area; and (2) landscape and structural properties of territory of 1 southwestern United States and northwestern Mexico.

A.L.

N75-11438* Scientific Translation Service, Santa Barbara, Calif.
USE OF TV PICTURES OBTAINED FROM WEATHER
SATELLITES TO STUDY THE GEOLOGICAL STRUCTURE
OF THE EARTH

I. I. Bashilova In its Space Iconics (NASA-TT-F-798) Sep. 1974 p 190-194 Transl into ENGLISH of the publ. "Kosmicheskaya Ikonika" Moscow, Nauka Press, 1973 p 159-162 CSCL 08G

Results are presented of Soviet investigations into the use of television imagery obtained from meteorological satellites for the interpretation of tectonics, lithology, and structural properties of the earth. The area investigated encompassed the eastern part of Iran, and the western regions of Afghanistan and Pakistan. Interpretation of the features identified during the analysis was accomplished by comparing these features with physical, geographic, geological, and tectonic maps. It was concluded that television pictures of the earth from space can be used for geological and structural analysis.

A.L.

N75-11458# Bureau of Natural Gas, Washington, D.C.
OFFSHORE INVESTIGATION: PRODUCIBLE SHUT-IN
LEASES AS OF JANUARY 1974, PHASE 2
Jul. 1974 30 p Original contains color illustrations

Avail: NTIS HC \$3.75

Estimates were made of the proved and probable gas reserves underlying the producible shut-in leases of the Louisiana and Texas Outer Continental Shelf. Data from the files of the U.S. Geological Survey and the Federal Power Commission were used. Results are summarized in tabular form.

N75-12392*# Environmental Research Inst. of Michigan, Ann Arbor. Infrared and Optics Div.

MAPPING EXPOSED SILICATE ROCK TYPES AND EXPOSED FERRIC AND FERROUS COMPOUNDS FROM A SPACE PLATFORM Quarterly Report, 8 Jun. - 8 Sep. 1974 Frederick J. Thomson, Principal Investigator 11 Nov. 1974 3 p

(Contract NAS9-13317) (E75-10008; NASA-CR-140658; ERIM-102000-27-L). Avail: NTIS HC \$3.25 CSCL 08B

N75-12416*# Bureau of Mineral Resources, Geology and Geophysics, Canberra (Australia). Div. of Mineral Physics.

A STUDY OF THE USEFULNESS OF SKYLAB EREP DATA FOR EARTH RESOURCES STUDIES IN AUSTRALIA Quarterly Progress Report, Aug. - Oct. 1974
K. L. Burns and J. Shepherd, Principal Investigators Oct. 1974

1 p Sponsored by NASA EREP (E75-10047; NASA-CR-140784) Avail: NTIS HC \$3.25 CSCL-05B

N75-13346*# California Earth Science Corp., Santa Monica.
ANALYSIS OF PSEUDOCOLOR TRANSFORMATIONS OF
ERTS-1 IMAGES OF SOUTHERN CALIFORNIA AREA

P. M. Merifield, Principal Investigator, D. L. Lamar, R. H. Stratton, J. V. Lamar, and Carl Gazley, Jr. Oct. 1974 27 p refs Prepared in cooperation with the RAND Corp., Santa Monica, Calif. Original contains color, imagery. Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue, Sioux Falls, S. D. 57198 ERTS

(Contracts NAS2-7698: DI-14-08-0001-13911)

(E75-10037; NASA-CR-140774; TR-74-2) Avail: NTIS HC \$3.75 CSCL 05B

The author has identified the following significant results. Representative faults and lineaments, natural features on the Mojave Desert, and cultural features of the southern California area were studied on ERTS-1 images. The relative appearances of the features were compared on a band 4 and 5 subtraction image, its pseudocolor transformation, and pseudocolor images of bands 4, 5, and 7. Selected features were also evaluated in a test given students at the University of California, Los Angeles. Observations and the test revealed no significant improvement in the ability to detect and locate faults and lineaments on the pseudocolor transformations. With the exception of dry lake surfaces, no enhancement of the features studied was observed on the bands 4 and 5 subtraction images. Geologic and geographic features characterized by minor tonal differences on relatively flat surfaces were enhanced on some of the pseudocolor images.

N75-13351*# Geological Survey, Denver, Colo.
REMOTE SENSING GEOPHYSICS FROM SKYLAB Monthly
Report, Oct. 1974

Kenneth Watson, Principal Investigator and H. A. Pohn Oct. 1974 2 p EREP

(NASA Order T-6555-B)

(E75-10051; NASA-CR-140953) Avail: NTIS HC \$3.25 CSCL 08G

N75-13354*# Geological Survey, Denver, Colo.

APPLICATION OF SKYLAB EREP PHOTOGRAPHS TO STUDY OF THE MODERN EPISODE OF ACCELERATED EROSION IN SOUTHERN ARIZONA Progress Report, 1 Jan. - 30 Jun. 1974

Roger B. Morrison, Principal Investigator 1 Jul. 1974 12 p

(NASA Order T-4113-B)

(E75-10054; NASA-CR-140962) Avail: NTIS HC \$3.25 CSCL

The author has identified the following significant results. Indexing and analysis of the SL 2, SL 3, and SL 4 photos of the project area has shown that S-190A coverage with less than 30% clouds totals about 123,000 sq km. The 70-mm unenlarged color, color-infrared, B/W red, and B/W green bands from S-190A are of good to excellent quality; the B/W IR bands from SL 2 are excessively grainy and have very low resolution; those from SL 3 are better but nevertheless have low resolution. The 5-inch unenlarged color transparencies from S-190B are generally of excellent photographic quality. However, where cloud cover is extensive, commonly the S-190A and S-190B color and color-IR photos are correctly exposed for the clouds but considerably underexposed for the ground. The 4X enlargements of all bands of S-190A photos taken by SL 2 are much fuzzier than they should be; evidently the enlarger was not focused properly. The 4X enlargements from SL 3 are much superior.

N75-13365*# Geological Survey, Denver, Colo.

APPLICATIONS OF SKYLAB EREP PHOTOGRAPHS TO MAPPING OF LANDFORMS AND ENVIRONMENTAL GEOLOGY IN THE GREAT PLAINS AND MIDWEST Progress Report. 1 Jan. - 30 Jun. 1974

Roger B. Morrison, Principal Investigator 1 Jul. 1974 32 p ERFP

(NASA Order T-4647-B)

(E75-10055; NASA-CR-140963) Avail: NTIS HC \$3.75 CSCL 08B

The author has identified the following significant results. The utility of Skylab 2 and 3 S-190A multispectral photos for environmental-geologic/geomorphic applications is being tested by using them to prepare 1:250,000-scale maps of geomorphic features, surficial geology, geologic linear features, and soil associations of large, representative parts of the Great Plains and Midwest. Parts of Nebraska, Iowa, Missouri, and South Dakota were mapped. The maps were prepared primarily by interpretation of the S-190A photos, supplemented by information from topographic, geologic, and soil maps and reports. The color band provides the greatest information on geology, soils, and geomorphology; its resolution also is the best of all the multispectral bands and permits maximum detail of mapping. The color-IR band shows well the differences in soil drainage and moisture, and vegetative types, but has only moderate resolution. The B/W-red band is superior for topographic detail and stream alinements. The B/W-infrared bands best show differences in soil moisture and drainage but have poor resolution, especially those from SL 2. The B/W-green band generally is so low contrast and degraded by haze as to be nearly useless. Where stereoscopic coverage is provided, interpretation and mapping are done most efficiently using a Kern PG-2 stereoplotter.

N75-13367*# Geological Survey, Denver, Colo.
REMOTE SENSING GEOPHYSICS FROM SKYLAB, Monthly
Report, Jul. 1974

Kenneth Watson, Principal Investigator and H. A. Pohn Jul. 1974 2 p EREP

(NASA Order T-6555-B)

(E75-10057; NASA-CR-140965) Avail: NTIS HC \$3.25 CSCL 08E

N75-13358*# Geological Survey, Denver, Coto.

REMOTE SENSING GEOPHYSICS FROM SKYLAB Monthly Report, Aug. 1974

Kenneth Watson, Principal Investigator and H. A. Pohn Aug. 1974 2 p EREP

(NASA Order T-6555-B)

(E75-10058; NASA-CR-140966) Avail: NTIS HC \$3.25 CSCL ORF

N75-13360*# Jet Propulsion Lab., Calif. Inst. of Tech., Pasadena. PRELIMINARY RESULTS IN THE COMPARISON OF SKYLAB, ERTS AND RB-57 IMAGES FOR THE DETECTION OF LINEAMENTS AND FRACTURES IN PRECAMBRIAN, PALEOZOIC, AND LATE TERTIARY ROCKS ON AND NEAR THE COLORADO PLATEAU, NORTH-CENTRAL ARIZONA Progress Report

A. F. H. Goetz, Principal Investigator, William D. DiPaolo, and Donald P. Elston 12 Dec. 1974 7 p EREP (E75-10080; NASA-CR-141000) Avail: NTIS HC \$3.25 CSCL 086

N75-14188 # Consiglio Nazionale delle Ricerche, Milan (Italy). Lab. per la Geofisica della Litosfera. CONTRIBUTION OF SPACE PLATFORMS TO A GROUND AND AIRBORNE REMOTE SENSING PROGRAMME OVER ACTIVE ITALIAN VOLCANOES

R. Cassinis, Principal Investigator, G. M. Lechi, C. M. Marino, and A. M. Tonelli 5 Nov. 1974 41 p refs Sponsored by NASA Original contains imagery. Original photography may be purchased from the EROS Data Center, Sioux Falls, S. D. 57198 EREP

(E75-10069; NASA-CR-141102) Avail: NTIS HC \$3.75 CSCL 08F

The author has identified the following significant results. A method has been suggested for the forecasting of the lateral eruptions of Mount Etna, through the multispectral analysis of the vegetation behavior. Unknown geological lineaments which seem to be related to deep crustal movements have been discovered using the ERTS-1 imagery. Results in the geological field were obtained in the study of the general structure of the Alpine range. In the field of official vegetation classification. ERTS-1 images were used for a preliminary study of rice fields in northern Italy Very good experimental results have been obtained using the Skylab multispectral photographs. In the field of hydrogeology and soil type discrimination discoveries of unknown paleoriver beds have been made in the northeastern part of the Po Valley using the multispectral imagery of SL3. The superior resolution of Skylab was a fundamental element for the success of this investigation.

N75-14190*# Colorado School of Mines, Golden. Dept. of Geology.

GEOLOGIC AND MINERAL AND WATER RESOURCES INVESTIGATIONS IN WESTERN COLORADO, USING SKYLAB EREP DATA Monthly Progress Report, Nov. 1974 Keenan Lee, Principal Investigator 19 Dec. 1974 5 p EREP (Contract NAS9-13394)

(E75-10071; NASA-CR-141100) Avail: NTIS HC \$3:25 CSCL 108G

N75-14191*# Purdue Univ., Lafayette, Ind. Lab. for Applications of Remote Sensing.

AN INTERDISCIPLINARY ANALYSIS OF MULTISPECTRAL SATELLITE DATA FOR SELECTED COVER TYPES IN THE COLORADO MOUNTAINS, USING AUTOMATIC DATA PROCESSING TECHNIQUES Monthly Progress Report, Nov. 1974

Roger M. Hoffer, Principal Investigator Nov. 1974 4 p EREP (Contract NAS9-13380)

(E75-10072; NASA-CR-141099) Avail: NTIS HC \$3.25 CSCL

The author has identified the following significant results. The utility of the middle infrared portion of the spectrum for snow cover mapping has been demonstrated. Also, the S-192 thermal band seems to have a higher thermal detectivity than previously estimated and predicted.

N75-14195*# Nevada Univ., Reno. Mackay School of Mines

THE GREAT BASIN INVESTIGATION Monthly Progress Report, Nov. 1974

Jack G. Quade, Principal Investigator Nov. 1974 2 p EREP (Contract NAS9-13274)

(E75-10076; NASA-CR-141095) Avail: NTIS HC \$3.25 CSCL 08E

N75-14237 Milan Univ. (Italy). Inst. of Mineralogy.
GEOLOGICAL FEATURES ON ERTS-1 IMAGES OF A TEST
AREA IN THE WEST-CENTRAL ALPS
A. Boriani, C. M. Marino, and R. Sacchi (Turin Univ.) In ESRO

European Earth Resources Satellite Expts. May 1974 p 199-204 refs

A geologic approach was made to the definition of the capability of ERTS-1 to enhance the main geologic characteristics of some typical areas of the Alps. The area examined in the first stage of this research is in the west-central Alps region around some of the main Italian lakes (Orta, Maggiore, Como). A certain number of geologic lineaments were enhanced. Their comparison with previous geologic data shows a very interesting peculiarity: almost all the lineaments discovered are vertical faults. In order to evaluate the findings properly, many ground surveys were performed along the observed lineaments. Some of the most interesting discoveries concern the well-known Insubric Line. Some conclusions can be drawn regarding the ability of ERTS-1 type satellites to assist tectonic or general geologic studies in areas which in the past have only been studied in great detail Author (ESRO) in a traditional way.

N75-14238 Zentralstelle fuer Geo-Photogrammetrie und Fernerkundung, Munich (West Germany).

GEOLOGIC EVALUATION OF CENTRAL ITALY FROM ERTS-1 AND SKYLAB DATA

J. Bodechtel, J. Nithack, and R. Haydn In ESRO European Earth Resources Satellite Expts. May 1974 p 205-215 refs

Utilizing ERTS-1 and Skylab imagery, linear features of the Apennines were enhanced by conventional photogeologic methods and by optical processing techniques using a coherent light source. Horizontal and vertical faults of regional dimensions could be detected. On the basis of fieldwork and aerial photointerpretation, the lineations revealed by ERTS data could be interpreted as ac, bc and shearing faults of an older rectangular system, and of a younger system. A difference in the direction of the b-axis is explained as caused by a clockwise rotation of larger blocks, whereas the causal main stress remained stable in a southwest-northeast direction. Overthrusts in the Umbrian Apennines and the triangular shape of the Po Valley are explained by the clockwise rotation. The transform faults bordering the blocks of the Northern Appennies, the Tuscan-Umbrian Apennines and the Abruzzi Mountains could be shearing faults due to this main stress.

Author (ESRO)

N75-14240 Bureau de Recherches Geologiques et Minieres, Orleans (France).

ERTS IMAGE CONTRIBUTION TO GEOLOGICAL KNOWLEDGE COVERING DIFFERENT TYPES OF CLIMATES (FRANCE, TOGO, DAHOMEY, MADAGASCAR) [CONTRIBUTION DES IMAGES ERTS A LA CONNAISSANCE GEOLOGIQUE SOUS DIFFERENTS TYPES DE CLIMATS (FRANCE, TOGO, DAHOMEY, MADAGASCAR)]

M. Koch, J. Y. Scanvic, and G. Weecksteen In ESRO European Earth Resources Satellite Expts. May 1974 p 231-241 In FRENCH: ENGLISH summary

Possibilities of using ERTS images of several climatological areas were enabled by the choice of three test areas, situated in temperate, tropical and subdesert regions, respectively, with or without a laterite cover. Whereas the subdesert climate favored the observation of outcrop formations and hence facilitated its interpretation, as was expected, a large number of indirect criteria highlight structural elements or some lithological units in temperature region.

Author (ESRO)

N75-14244 Institut Geographique National, Paris (France).
COMPARISON OF VARIOUS NUMERICAL TECHNIQUES
FOR CLASSIFICATION OF MULTISPECTRAL DATA.
APPLICATION TO GEOLOGY [COMPARAISON DE
DIFFERENTES TECHNIQUES NUMERIQUES SERVANT A
LA CLASSIFICATION DES DONNEES MULTISPECTRALES.
APPLICATION A LA GEOLOGIE]

F. Brun, A. Fontanel (Inst. Franc. du Petrole, Paris), C. Lallemand (Inst. Franc. du Petrole, Paris), G. Legendre (Inst. Franc. du Petrole, Paris), J. C. Rivereau (Inst. Franc. du Petrole, Paris), and G. Thomas (Inst. Franc. du Petrole, Paris) In ESRO European

Earth Resources Satellite Expts. May 1974 p 303-316 refs In FRENCH; ENGLISH summary Original contains color illustrations

Numerical computation was used to distinguish and map various terrains, highways, rivers, etc....on ERTS image taken on 6 October 1972 over the Rhone Valley (France). For this purpose three numerical techniques were used: (1) discriminatory analysis (supervised technique) for which it is necessary to define and delineate characteristic zones from which the identification algorithms are calculated: (2) principa! component analysis (unsupervised technique) for which the different classes are obtained directly by computation; (3) adaptive analysis technique which is easier to implement than the two preceding techniques: on the one hand only selected zones are automatically mapped with the computer and on the other hand the delineation of the corresponding test sites (training areas) is not needed. These three techniques were tested on the same area (1500 sq km) and the results compared with vegetation and geological maps.

Author (ESRO)

N75-15103*# Wolf Research and Development Corp., Pocomoke City, Md.

APPLICABILITY OF SKYLAB REMOTE SENSING FOR DETECTION AND MONITORING OF SURFACE MINING ACTIVITIES Quarterly Progress Report, 1 Oct. - 31 Dec. 1974

R. L. Brooks, Principal Investigator and J. D. Pennewell Jan. 1975 8 p EREP (Contract NAS9-13310)

(E75-10085; NASA-CR-141149; QPR-4) Avail: NTIS HC \$3.25 CSCL 08i

N75-15106*# Stanford Univ., Calif. School of Earth Sciences.

FEASIBILITY OF USING S-191 INFRARED SPECTRA FOR GEOLOGICAL STUDIES FROM SPACE Monthly Reports, 1 Jul. - 30 Nov. 1974

J. P. Lyon and F. R. Honey, Principal Investigators 30 Nov. 1974 3 p EREP

(Contract NAS9-13357)

(E75-10088; NASA-CR-141152; MR-12; MR-13; MR-14; MR-15; MR-16) Avail: NTIS HC \$3.25 CSCL 08G

N75-15107*# Stanford Univ., Calif. School of Earth Sciences.

FEASIBILITY OF USING S-191 INFRARED SPECTRA FOR GEOLOGICAL STUDIES FROM SPACE Monthly Report, Dec. 1974

J. P. Lyon and F. R. Honey, Principal Investigators 30 Dec. 1974 3 p EREP

(Contract NAS9-13357)

(E75-10089; NASA-CR-141153; MR-17) Avail: NTIS HC \$3.25 CSCL 08G

N75-15120*# Nevada Univ., Reno. Mackay School of Mines.

THE GREAT BASIN INVESTIGATION Monthly Progress Report, Dec. 1974

Jack G. Quade, Principal Investigator Dec. 1974 2 p EREP (Contract NAS9-13274)

(E75-10102; NASA-CR-141298) Avail: NTIS HC \$3.25 CSCL 08G

N75-15124*# California Earth Science Corp., Santa Monica.
FAULT TECTONICS AND EARTHQUAKE HAZARDS IN THE
PENINSULAR RANGES, SOUTHERN CALIFORNIA Monthly
Progress Report, Dec. 1974
Paul M. Merifield, Principal Investigator 5 Jan. 1975 2 p
EREP
(Contract NAS2-7698)
(E75-10108; NASA-CR-140898; MPR-19) Avail: NTIS
HC.\$3.25 CSCL 08G

N75-15132*# National Aeronautics and Space Administration. Goddard Space Flight Center, Greenbelt, Md.
GEOLOGIC STRUCTURE IN CALIFORNIA: THREE STUDIES WITH ERTS-1 IMAGERY
Paul D. Lowman, Jr. Nov. 1974 23 p refs Submitted for publication
(NASA-TM-X-70799; X-922-74-314) Avail: NTIS HC \$3.25 CSCL 08G

. 1

Results are presented of three early applications of imagery from the NASA Earth Resources Technology Satellite to geologic studies in California. In the Coast Ranges near Monterey Bay, numerous linear drainage features possibly indicating unmapped fracture zones were mapped within one week after launch of the satellite. A similar study of the Sierra Nevada near Lake Tahoe revealed many drainage features probably formed along unmapped joint or faults in granitic rocks. The third study, in the Peninsular Ranges, confirmed existence of several major faults not shown on published maps. One of these, in the Sawtooth Range, crosses in Elsinore fault without lateral offset; associated Mid-Cretaceous structures have also been traced continuously across the fault without offset. It therefore appears that displacement along the Elsinore fault has been primarily of a dip-slip nature, at least in this area, despite evidence for lateral displacement elsewhere.

N75-15133*# National Aeronautics and Space Administration. Goddard Space Flight Center, Greenbelt, Md.

A DEMONSTRATION OF ERTS-1 ANALOG AND DIGITAL TECHNIQUES APPLIED TO STRIP MINING IN MARYLAND AND WEST VIRGINIA

Arthur T. Anderson and Jane Schubert Nov. 1974 19 p refs Submitted for publication

(NASA-TM-X-70798; X-923-74-313) Avail: NTIS HC \$3.25 CSCL 08I

The largest contour strip mining operations in western Maryland and West Virginia are located within the Georges Creek and the Upper Potomac Basins. These two coal basins lie within the Georges Creek (Wellersburg) syncline. The disturbed strip mine areas were delineated with the surrounding geological and vegetation features using ERTS-1 data in both analog (imagery) and digital form. The two digital systems used were: (1) the ERTS-Analysis system, a point-by-point digital analysis of spectral signatures based on known spectral values, and (2) the LARS Automatic Data Processing System. The digital techniques being developed will later be incorporated into a data base for land use planning. These two systems aided in efforts to determine the extent and state of strip mining in this region. Aircraft data, ground verification information, and geological field studies also aided in the application of ERTS-1 imagery to perform an integrated analysis that assessed the adverse effects of strip mining. The results indicated that ERTS can both monitor and map the extent of strip mining to determine immediately the acreage affected and indicate where future reclamation and revegetation may be necessary. Author

Page intentionally left blank

05

OCEANOGRAPHY AND MARINE RESOURCES

Includes sea-surface temperature, ocean bottom surveying imagery, drift rates, sea ice and icebergs, sea state, fish location

A75-10644 * # The detection by ERTS-1 of wind-induced ocean surface features in the lee of the Antilles Islands. R. Cram and K. Hanson (Miami, University, Coral Gables, Fla.). Journal of Physical Oceanography, vol. 4, Oct. 1974, p. 594-600. 8 refs. Contract No. NOAA-04-30-2212. NASA Order S-70246-AG-1.

Photographic data received from the ERTS-1 satellite over the Lesser Antilles Islands show distinct ocean features on the leeward side of each island. Attempts to relate these features to ocean eddy formations with the aid of ground truth data proved unsuccessful. However, surface and upper air wind data indicate a good correlation with the size, shape, and downwind extent of the ocean features. Studies to date indicate strongly that these features result from horizontal differences in a sea surface roughness due to the wind-shadow effect of the islands. The results suggest that horizontal variations in the reflectance of the sea surface will make remote sensing of the ocean mixed layer more difficult than previously anticipated. The surface reflection seems to be large enough to mask the smaller variations in backscattered energy from the mixed layer. Efforts to limit the effect of surface reflectance by photographic differencing of two multi-spectral scanner bands were unsuccessful. (Author)

A75-10645 # A satellite-aircraft thermal study of the upwelled waters off Spanish Sahara: P. E. La Violette (U.S. Department of Defense, Naval Oceanographic Office, Washington, D.C.). Journal of Physical Oceanography, vol. 4, Oct. 1974, p. 676-684. 10 refs. DOD-Navy-sponsored research.

Review of the upwelling along the coast of Spanish Sahara and of several associated thermal features observed in a satellite-aircraft oceanographic survey conducted during the period from the 18th through the 26th of August 1973. Analyses of the aircraft and satellite radiation data show that these features consisted of variable cool and warm eddies which extended more than 100 km northwest from the coastal upwelling zone. These features may represent a periodic occurrence.

M.V.E.

A75-11255 * Satellite photography of eddies in the gulf loop current. G. A. Maul (NOAA, Physical Oceanography Laboratory, Miami, Fla.), D. R. Norris (NASA, Johnson Space Center, Earth Observations Div., Houston, Tex.), and W. R. Johnson (Lockheed Electronics Co., Inc., Houston, Tex.). Geophysical Research Letters, vol. 1, Oct. 1974, p. 256-258. 9 refs.

Cyclonic ocean eddies, approximately 12 to 32 kilometers in diameter, have been photographed in the eastern Gulf of Mexico by SKYLAB. Apparently caused by horizon velocity shear, these features are associated with the Gulf Loop Current, whose position was known a fortnight before and after the observation. The eddies were discovered in sunlight-enhanced patterns of streamlines on the surface, and appear to be embedded in the flow. (Author)

A75-12034 * # Skylab and ERTS-1 investigations of coastal land-use and water properties in Delaware Bay. V. Klemas, D. Bartlett (Delaware, University, Newark, Del.), and R. Rogers (Bendix Corp., Aerospace Systems Div., Ann Arbor, Mich.). American Institute of Aeronautics and Astronautics and American Geophysical Union, Conference on Scientific Experiments of Skylab, Huntsville,

Ala., Oct. 30-Nov. 1, 1974, AIAA Paper 74-1220. 10 p. 10 refs. Contracts No. NAS5-21837; No. NAS1-12304; No. N00014-69-A-0407.

Study of coastal land use and water properties of Delaware Bay using digital and visual analysis of enhanced imagery from NASA's Earth Resources Technology Satellite (ERTS) and from the Skylab Earth Resources Experimental Package (EREP). ERTS is shown to have the advantage of repetitive coverage of the test site which facilitates change detection experiments by gathering data under a variety of tidal, seasonal, and atmospheric conditions. Skylab-EREP data, on the other hand, are superior in both spatial and spectral resolution.

M.V.E.

A75-12719 # Monitoring and predicting El Niño invasions. W. H. Quinn (Oregon State University, Corvallis, Ore.). Journal of Applied Meteorology, vol. 13, Oct. 1974, p. 825-830. 12 refs. Navy-supported research; NSF Grants No. GA-1571; No. GA-27205.

El Niño refers to the anomalously warm surface water invasion along the southern Ecuadorian and Peruvian coastal regions which are usually under the influence of cooler waters from coastal upwelling and the northward flowing Peru Current. Southern Oscillation indices (differences in sea level atmospheric pressure between Easter Island and Darwin, Australia, and between Juan Fernandez Island and Darwin) were treated so as to emphasize interannual changes, and considered for monitoring unusual equatorial Pacific ocean atmosphere developments and certain of their consequences (e.g., El Niño invasions). It now appears that their trends can be used to predict activity of El Niño intensity several months in advance.

A75-15179 # The wind vector in a layer over water (O vektore vetra v privodnom sloe). E. K. Korchagin, R. N. Semenov, and I. A. Mandurovskii. *Meteorologiia i Gidrologiia*, Oct. 1974, p. 41-48, 8 refs. In Russian.

The horizontal component of the wind vector over the sea was determined from an aircraft at various altitudes from 100 to 1000 m. A Doppler radar for measuring ground speed and drift angle, an airspeed sensor, and a course selector were used. The surface-wind parameters were determined from aerial photographs of the sea surface and from calculations with respect to the atmospheric pressure field. The statistical relationship found between the surface-wind vector and the vector determined on board the aircraft supports the possibility of finding surface-wind and water-surface parameters from aircraft flying at altitudes up to 1000 m, and of extrapolating ship-borne observations of the wind vector to altitudes in the friction layer.

A.T.S.

A75-17120 # A study of microwave emission properties of sea ice - AIDJEX 1972. D. C. Meeks (Aerojet ElectroSystems Co., Azusa, Calif.), R. O. Ramseier (Department of the Environment, Ottawa, Canada), and W. J. Campbell (U.S. Geological Survey, Tacoma, Wash.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 307-322. 8 refs.

Results derived from a comparative study of surface-based 13.4 GHz passive microwave measurements with detailed surface-truth measurements concerning the physical, chemical and structural properties of Arctic sea ice illustrate distinct decreasing microwave emissions for first-year, transitional and multi-year sea ice types. For transitional and multi-year ice, a structural relationship for microwave emissions by sea ice exists. Both vertically and horizontally polarized measured brightness temperatures decrease linearly with increasing average ice porosity. In the case of first-year ice, however, measured brightness temperatures are comparatively uniform and microwave emission appears to be more strongly influenced by high near-surface salinity concentrations combined with the occurrence of uniform porosity.

(Author)

A75-17121 * # Investigation of radar discrimination of sea ice. S. K. Parashar, A. W. Biggs, A. K. Fung, and R. K. Moore (University of Kansas Center for Research, Inc., Lawrence, Kan.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 323-332. 15 refs. Research supported by the University of Kansas; Contracts No. NAS9-10261; No. N60921-70-C-0221.

The ability of radar to discriminate sea ice types and their thickness was studied. Radar backscatter measurements at 400 MHz (multi-polarization) and 13.3 GHz (VV polarization) obtained from NASA Earth Resources Aircraft Program Mission 126 were analyzed in detail. The scatterometer data were separated into seven categories of sea ice according to age and thickness as interpreted from stereo aerial photographs. The variations of radar backscatter cross-section with sea ice thickness at various angles are presented at the two frequencies. There is a reversal of angular character of radar return from sea ice less than 18 cm thick at the two frequencies. Multi-year ice (sea ice greater than 180 cm thick) gives strongest return at 13.3 GHz. First-year ice (30 cm to 90 cm thick) gives strongest return at 400 MHz. Open water can be differentiated at both the frequencies. Four-polarization 16.5 GHz radar imagery was also obtained. Open water and three categories of sea ice can be identified on the images. The results of the imagery analysis are consistent with the radar scatterometer results. (Author)

A75-17177 * # Computer derived coastal water classifications via spectral signatures. D. K. Clark, J. B. Zaitzeff, L. V. Strees, and W. S. Glidden (NOAA, National Environmental Satellite Service, Washington, D.C.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1213-1239. 9 refs. NASA-NOAA-supported research.

In April 1973, the National Environmental Satellite Service conducted a remote sensing investigation within the coastal waters of the New York Bight. Remote sensor records acquired from the ERTS-1 Multispectral Scanner and the Bendix 24 Channel Multispectral Scanner records flown on the NASA C-130 were used for water mass classification. Computer-derived classifications are discussed and compared. Such features as the Hudson River's turbid discharge plumes, acid waste and shelf water are examined in terms of their distribution of suspended particulates (2-203 microns), transmissivity, diffuse attenuation, incident and returned spectral irradiances. The characteristics of these features and their relationship to the computer derived classifications are presented and discussed with respect to radiative transfer theory. (Author)

A75-17183 * # Ocean internal waves off the North American and African coasts from ERTS-1. J. R. Apel, R. L. Charnell (NOAA, Atlantic Oceanographic and Meteorological Laboratories, Miami, Fla.), and R. J. Blackwell (California Institute of Technology, Jet Propulsion Laboratory, Pasadena, Calif.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1345-1351. ARPA-sponsored research.

A75-17184 # Lake Ontario water mass delineation from ERTS-1. J. C. Munday, Jr. (Erindale College, Clarkson, Ontario, Canada). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1355-1368. 18 refs. Department of Supply and Services Contracts No. 01GRKW111-2-1073; No. 0GR2-0522.

Multiband analysis of ERTS-1 MSS photographic images of western Lake Ontario has been conducted to delineate water masses.

Modification of water hues by thin cloud cover is detected by chromaticity analysis. Density-sliced MSS 5 images indicate suspended sediment plumes near river and harbor mouths and in near-shore zones, and discoloration in scattered other areas. High percentage cloud cover necessitates a long-term ERTS mapping project to obtain a useful satellite picture of Lake Ontario circulation dynamics. (Author)

A75-17203 # Measurement of sea surface currents using airborne Doppler radar and inertial navigation systems. J. F. R. Gower (Environment Canada, Marine Sciences Directorate, Victoria, British Columbia, Canada). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1657-1667.

A75-17233 # Application of ERTS-1 data to the protection and management of New Jersey's coastal environment. R. L. Mairs, R. T. Macomber, D. T. Stanczuk, F. J. Wobber, L. R. Pettinger (Earth Satellite Corp., Washington, D.C.), R. S. Yunghans, E. B. Feinberg, and J. Stitt (New Jersey Department of Environmental Protection, Trenton, N.J.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann. Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 2087-2091.

A75-17479 Pacific range electromagnetic signature studies /Project PRESS/. W. S. Hahn (RCA, Missile and Surface Radar Div., Moorestown, N.J.). RCA Engineer, vol. 20, Oct. Nov. 1974, p. 60-63.

The activities of the Pacific range electromagnetic signature studies (PRESS) program including mission planning, test execution, data reduction and analysis, and the development of radar for sensor systems are reviewed. The paper discusses the roles of the PRESS program which are (1) to provide metric and signature data in test reentry vehicles, and (2) to act as a radar development test bed. Certain features of the sensors and related systems in the PRESS program are summarized.

A75-18124 # The gravity field and the definition of stationary sea surface topography. R. S. Mather (New South Wales, University, Kensington, Australia). In: Geodesy and physics of the earth; Proceedings of the Second International Symposium, Potsdam, East Germany, May 7-11, 1973. Part 2. Potsdam, East Germany, Deutsche Akademie der Wissenschaften, Zentralinstitut für Physik der Erde, 1974, p. 381-414, 21 refs.

The investigation reported shows that for the intended investigation of ocean circulation it is necessary to determine accurately the instantaneous geocentric position of the spacecraft carrying the altimeter used in the study. The criteria required in the definition of the global gravity anomaly field are discussed along with the determination of the features of the ocean geoid.

G.R.

N75-10529*# National Oceanic and Atmospheric Administration. Miami, Fla.

REMOTE SENSING OF OCEAN CURRENT BOUNDARY LAYER Monthly Progress Report. Sep. 1974
George A. Maul, Principal Investigator Sep. 1974 2 p EREP (NASA Order T-4713-B) (E74-10801; NASA-CR-140495) Avail: NTIS HC \$3.25 CSCL 08C

N75-10554* Corps of Engineers, Washington, D.C. ERTS PROGRAM OF THE US ARMY CORPS OF ENGINEERS

Jack W. Jarman In NASA. Goddard Space Flight Center Third

ERTS Symp., Vol. 2 May 1974 p 62-75

CSCL 08E

The Army Corps of Engineers research and development efforts associated with the ERTS Program are confined to applications of investigation, design, construction, operation, and maintenance of water resource projects. Problems investigated covered: (1) resource inventory; (2) environmental impact: (3) pollution monitoring; (4) water circulation; (5) sediment transport; (6) data collection systems; (7) engineering; and (8) model verification. These problem areas were investigated in relation to bays, reservoirs, lakes, rivers, coasts, and regions. ERTS-1 imagery has been extremely valuable in developing techniques and is now being used in everyday applications.

N75-10555* National Oceanic and Atmospheric Administration, Washington, D.C.

A REVIEW OF INITIAL INVESTIGATIONS TO UTILIZE ERTS-1 DATA IN DETERMINING THE AVAILABILITY AND DISTRIBUTION OF LIVING MARINE RESOURCES

William H. Stevenson (NOAA, Bay St. Louis, Mo.), Andrew J. Kemmerer, Buddy H. Atwell (NASA. Mississippi Test Facility), and Paul M. Maughan (Earth Satellite Corp.) In NASA. Goddard Space Flight Center Third ERTS Symp., Vol. 2 May 1974 p 76-86 refs CSCL 08A

The National Marine Fisheries Service has been studying the application of aerospace remote sensing to fisheries management and utilization for many years. The 15-month ERTS study began in July 1972 to: (1) determine the reliability of satellite and high altitude sensors to provide oceanographic parameters in coastal waters; (2) demonstrate the use of remotely-sensed oceanographic information to predict the distribution and abundance of adult menhaden; and (3) demonstrate the potential use of satellites for acquiring information for improving the harvest and management of fisheries resources. The study focused on a coastal area in the north-central portion of the Gulf of Mexico, including parts of Alabama, Mississippi, and Louisiana. The test area used in the final analysis was the Mississippi Sound and the area outside the barrier islands to approximately the 18-meter (10-fathom) curve.

N75-10559* National Aeronautics and Space Administration. Goddard Space Flight Center, Greenbelt, Md.

WATER RESOURCES

Vincent V. Salomonson In its Third ERTS Symp., Vol. 2 May 1974 p 126-137 CSCL 08H

ERTS-1 applications in snow and ice monitoring, surface water monitoring, including monitoring of wetland areas and flood inundated area mapping, and also watershed monitoring for runoff prediction are discussed. Results also indicate that geological features can be noted which relate to ground water. ERTS-1 data can be used successfully in operational situations by water resources management agencies. Author

N75-10566*# Virginia Inst. of Marine Science, Gloucester Point. REMOTE SENSING APPLICATIONS IN MARINE SCIENCE PROGRAMS AT VIMS

Hayden H. Gordon, Michael E. Penney, and Robert J. Byrne [1974] 86 p refs

(Contract NAS6-1902)

(NASA-CR-137475) Avail: NTIS HC \$4.75 CSCL 14B

Scientists at the Virginia Institute of Marine Science (VIMS) utdized remote sensing in three programs: (1) tonal variations in imagery of wetlands; (2) use of the thermal infrared to delineate the discharge cooling water at the Virginia Electric and Power Company (VEPCO) nuclear power station on the James River: and (3) the use of aerial photography to determine the volume storage function for water in the marsh-bay complex fed by Wachapreague Inlet on the Eastern Shore of Virginia. Details of the investigations are given, along with significant results.

N75-10571*# Chesapeake Research Consortium, Inc., Baltimore,

A SURVEY FOR THE USE OF REMOTE SENSING IN THE CHESAPEAKE BAY REGION

Robert E. Ulanowicz (Maryland Univ.) Jun. 1974 93 p (Contract NAS6-2433)

(NASA-CR-137476; CRC-Publ-6) Avail; NTIS HC \$4.75 CSCL

Environmental problem areas concerning the Chesapeake Bay region are reviewed along with ongoing remote sensing programs pertaining to these problems, and recommendations are presented to help fill lacunae in present research and to utilize the remote sensing capabilities of NASA to their fullest. A list of interested organizations and individuals is presented for each category. The development of technologies to monitor dissolved nutrients in bay waters, the initiation of a census of the disappearing rooted acquatic plants in the littoral zones, and the mapping of natural building constraints in the growth regions of the states of Maryland and Virginia are among the recommendations presented. Author

N75-10671*# Battelle Columbus Labs., Ohio. INTERACTION OF MARINE GEODESY, SATELLITE TECH-NOLOGY AND OCEAN PHYSICS

A. G. Mourad and D. M. J. Fubara Jun. 1972 105 p refs (Contract NAS6-2006)

(NASA-CR-137470) Avail: NTIS HC \$5.25 CSCL 08J

The possible applications of satellite technology in marine geodesy and geodetic related ocean physics were investigated. Four major problems were identified in the areas of geodesy and ocean physics: (1) geodetic positioning and control establishment; (2) sea surface topography and geoid determination; (3) geodetic applications to ocean physics; and (4) ground truth establishment. It was found that satellite technology can play a major role in their solution. For solution of the first problem, the use of satellite geodetic techniques, such as Doppler and C-band radar ranging, is demonstrated to fix the three-dimensional coordinates of marine geodetic control if multi-satellite passes are used. The second problem is shown to require the use of satellite altimetry, along with accurate knowledge of oceandynamics parameters such as sea state, ocean tides, and mean sea level. The use of both conventional and advanced satellite techniques appeared to be necessary to solve the third and fourth problems. Author

N75-10673*# Battelle Columbus Labs., Ohio. BISTATIC RADAR SEA STATE MONITORING

George T. Ruck, Donald E. Barrick, and Thaddeus Kaliszewski Jun. 1972 124 p refs

(Contract NAS6-2006)

(NASA-CR-137469) Avail: NTIS HC \$5.25 CSCL 08C

Bistatic radar techniques were examined for remote measurement of the two-dimensional surface wave height spectrum of the ocean. One technique operates at high frequencies (HF), 3-30 MHz, and the other at ultrahigh frequencies (UHF), approximately 1 GHz. Only a preliminary theoretical examination of the UHF technique was performed; however the principle underlying the HF technique was demonstrated experimentally with results indicating that an HF bistatic system using a surface transmitter and an orbital receiver would be capable of measuring the two-dimensional wave height spectrum in the vicinity of the transmitter. An HF bistatic system could also be used with an airborne receiver for ground truth ocean wave spectrum measurements. Preliminary system requirements and hardware configurations are discussed for both an orbital system and an aircraft verification experiment. Author

N75-10674# World Meteorological Organization, Geneva (Switzerland).

REPORT OF THE MEETING ON DRIFTING BUOYS FOR THE FIRST GARP GLOBAL EXPERIMEN?

Mar. 1974 91 p Conf. held at Geneva, Mar. 1974 Prepared jointly with Intern. Council for Sci. Unions

(GARP-Spec-Rept-13) Avail: NTIS HC \$4.75; WMO, Geneva

05 OCEANOGRAPHY AND MARINE RESOURCES

Within the framework of the first GARP global experiment, the status of the development of the buoy observing systems appropriate for the experiment is discussed. The requirements of the minimum capability of buoys for the experiment as well as the requirements for the polar orbiting system for the drifting buoy observing program are outlined. The deployment strategy, steps towards realization of the buoy system, and possible oceanographic programs during the FGGE using expendable buoys are also described.

N75-10681# Florida State Univ., Tallahassee.
THE COHO PROJECT: LIVING RESOURCES PREDICTION
FEASIBILITY STUDY, VOLUME 1 Final Report

James J. Brien, Bruce M. Woodworth, and David J. Wright 1974 37 p refs Prepared in cooperation with Oregon State Univ. (Grants NOAA-043-022-28; NSF GX-33502) (PB-234057/8; NSF/IDOE-74-18) Avail: NTIS HC \$3.25 CSCL 08C

The Coho project demonstrated a system which provides a substantial improvement in fish finding technique. It is the first known system which combines oceanographic and meteorological variables with the knowledge that certain species are temperature dependent, in order to arrive at an accurate forecast of the location of harvestable concentrations of salmon. The area chosen to test the prediction system was off the central coast of Oregon between Cape Lookout and Seal Rock.

N75-10682# Florida State Univ., Tailahassee.
THE COHO PROJECT: LIVING RESOURCES PREDICTION
FFASIRILITY STUDY VOLUME 2: ENVIRONMENTAL

REPORT

James J. OBrien, Bruce M. Woodworth, and David J. Wright 1974 191 p Prepared in cooperation with Oregon State Univ. (Grants NOAA-043-022-28: NSF GX-33502) (PB-234058/6; NSF/IDOE-74-19) Avail: NTIS HC\$5.50 CSCL 06C

The Coho project was a pilot project to study the application of remote sensing techniques for the benefit of the Central Oregon offshore Coho fishery. The system, designed to provide a true daily forecast of environmental factors conductive to concentrations of harvestable stocks of Coho salmon, was operated to include the offshore area between Cape Lookout (45 deg 20.5 min N) and Seal Rock (44 deg 30 min N) along the Central Oregon coast, during the period June 15, 19 1973, to August 16, 1973. This volume contains the data on all relevant environmental factors utilized in the prediction studies. GRA

N75-10683# Florida State Univ., Tallahassee.
THE COHO PROJECT: LIVING RESOURCES PREDICTION FEASIBILITY STUDY. VOLUME 3: SYSTEM EVALUATION REPORT

James J. OBrien, Bruce M. Woodworth, and David J. Wright 1974 64 p Prepared in cooperation with Oregon State Univ. (Grants NOAA-043-022-28; NSF GX-33502) (PB-234059/4; NSF/IDOE-74-20) Avail: NTIS HC\$3.75 CSCL

The Coho salmon is known to be a temperature dependent species preferring the relatively warm waters in the 52 degrees to 56 degrees Fahrenheit range. The upwelling phenomenon, on the other hand, brings cold water into the Coho habitat thus tending to drive the fish away. The cold jwater, however, is laden with nutrients which stimulate the initial phases of the food chain at the top of which exists the salmon. It is to be expected, and was presumed in this study, that Coho would be found on the warmer side of the interface of the upwelled and stabilized waters. This volume is devoted to the evaluation of a pilot prediction system operating off the central Oregon coast. The economic merits along with some unfavorable comments are set forth.

N75-11411*# Massachusetts Univ., Gloucester. Marine Station.

OCEAN COLOR, A THREE COMPONENT SYSTEM?

Charles S. Yentsch and William P. Owen 21 Sep. 1972 18 p refs

(Contract NAS5-21813)

(NASA-CR-139145) Avail: NTIS HC \$3.25 CSCL 08J

This study measures the concentrations of phytoplankton chlorophyll and yellow substance in the coastal waters of the Gulf of Maine. Sea surface observations attempt to delineate the principal biochemical parameters responsible for sea surface color. It is shown that the reddish-brown water changed to a blue-green in the open gulf.

M.C.F.

N75-12172*# National Aeronautics and Space Administration. Langley Research Center, Langley Station, Va.

A SIMULATION OF SYNTHETIC APERTURE RADAR IMAGING OF OCEAN WAVES

Calvin T. Swift 29 Oct. 1974 14 p Presented at the 1974 USNC/URSI Meeting, Boulder, Colorado, 14-17 Oct. 1974 (NASA-TM-X-72629) Avail: NTIS HC \$3.25 CSCL 171

A simulation of radar imaging of ocean waves with synthetic aperture techniques is presented. The modelling is simplistic from the oceanographic and electromagnetic viewpoint in order to minimize the computational problems, yet reveal some of the physical problems associated with the imaging of moving ocean waves. The model assumes: (1) The radar illuminates a one-dimensional, one harmonic ocean wave. (2) The scattering is assumed to be governed by geometrical optics. (3) The radar is assumed to be down-looking, with Doppler processing (range processing is suppressed due to the one-dimensional nature of the problem). (4) The beamwidth of the antenna (or integration time) is assumed to be sufficiently narrow to restrict the specular points of the peaks and troughs of the wave. The results show that conventional processing of the image gives familiar results if the ocean waves are stationary. When the ocean wave dispersion relationship is satisfied, the image is smeared due to the motion of the specular points over the integration time. In effect, the image of the ocean is transferred to the near field of the synthetic aperture. Author

N75-12396*# National Oceanic and Atmospheric Administration, Miami, Fla.

AN ASSESSMENT OF THE POTENTIAL CONTRIBUTIONS TO OCEANOGRAPHY FROM SKYLAB VISUAL OBSERVATIONS AND HAND-HELD PHOTOGRAPHY Monthly Progress Report, Aug. 1974

George A. Maul, Principal Investigator and Michael McCaslin Aug. 1974 18 p EREP (NASA Order T-4713-B)

(E75-10017; NASA-CR-140731) Avail: NTIS HC \$3.25 CSCL

N75-12407*# Wolf Research and Development Corp., Pocomoke City, Md.

APPLICABILITY OF SKYLAB REMOTE SENSING FOR DETECTION AND MONITORING OF SURFACE MINING ACTIVITIES Quarterly Progress Report, 1 Jul. - 30 Sep. 1974

R. L. Brooks, Principal Investigator and J. D. Pennewell Oct. 1974 6 p EREP (Contract NAS9-13310) (E75-10038: NASA-CR-140775; QPR-4) Avail: NTIS

HC \$3.25 CSCL 081

N75-13341*# National Aeronautics and Space Administration. Mississippi Test Facility, Bay Saint Louis.

EVALUATION OF SATELLITE REMOTE SENSING AND AUTOMATIC DATA TECHNIQUES FOR CHARACTERIZATION OF WETLANDS AND COASTAL MARSHLANDS Robert H. Cartmill, Principal Investigator [1974] 1 p ERTS (E75-10025; NASA-TM-X-72062) Avail: NTIS HC\$3.25 CSCL

08B

05 OCEANOGRAPHY AND MARINE RESOURCES

The author has identified the following significant results. ERTS digital data has been used to detect and measure accretion in the large estuarial system of the Atchafalaya River in southern Louisiana. Comparisons of aerial photography taken in October 1970 with a computer printout of ERTS digital data collected on October 1, 1972, show that in a delineated area of 1400 hectares (3460 acres) an accretion of land area in the amount of 112 hectares (277 acres) had occurred. Analysis of band 3 of the ERTS MSS was used to make a land-water map of the area. The accretion test area was marked on this map and the percentage of elements indicated as land was calculated. This percentage was then multiplied by the total test area to obtain the area of land on the date of the ERTS observation for comparison with the aerial photography. Significant improvement in classification accuracy of ERTS MSS data has been achieved by use of a priori probabilities of occurrence in pattern recognition programs. Two classifications were made with accuracies of 76.0% and 81.5%, respectively.

N75-13496*# Rockwell International Corp., Downey, Calif. Science Center.

MEASUREMENTS OF THE DIELECTRIC PROPERTIES OF SEA WATER AT 1.43 GHz Final Report

W. W. Ho, A. W. Love, and M. J. VanMelle Washington NASA Dec. 1974 39 p refs (Contract NAS1-10691)

(NASA-CR-2458) Avail: NTIS HC \$3.75 CSCL 08J

Salinity and temperature of water surfaces of estuaries and bay regions are determined to accuracies of 1 ppt salinity and 0.3 kelvin surface temperature. L-band and S-band radiometers are used in combination as brightness temperature detectors. The determination of the brightness temperature versus salinity, with the water surface temperature as a parameter for 1.4 GHz, is performed with a capillary tube inserted into a resonance cavity. Detailed analysis of the results indicates that the measured values are accurate to better than 0.2 percent in the electric property epsilon' and 0.4 percent in epsilon". The calculated brightness temperature as a function of temperature and salinity is better than 0.2 kelvin. Thus it is possible to reduce the measured data obtained with the two-frequency radiometer system with 1 ppt accuracy to values in the salinity range 5 to 40 ppt.

Author

N75-14186*# National Oceanic and Atmospheric Administration, Atlantic Oceanographic and Meteorological Labs. Miami, Fla. REMOTE SENSING OF OCEAN CURRENT BOUNDARY LAYER Monthly Progress Report, Oct. 1974 George A. Maul, Principal Investigator Oct. 1974 2 p EREP (NASA Order T-4713-B) (E75-10031; NASA-CR-140726) Avail: NTIS HC \$3.25 CSCL 08J

N75-14186 # National Oceanic and Atmospheric Administration, Atlantic Oceanographic and Meteorological Labs. REMOTE SENSING OF OCEAN CURRENT BOUNDARY LAYER Monthly Progress Report, Nov. 1974 George A. Maul, Principal Investigator Nov. 1974 2 p EREP (NASA Order T-4713-B) (E75-10067; NASA-CR-141021) Avail: NTIS CSCL 08J

- :...

N76-14228 Ecole Nationale Superieure du Petrole et des Monteurs, Rueil-Malmaison (France).

RECENT AND PRESENT SEDIMENTATION COMPLEXITY OF THE RHONE DELTA AND THE FRENCH MEDITER-RANEAN COAST (GOLION PROJECT) [COMPLEXITE DE LA SEDIMENTATION RECENTE ET ACTUELLE DU DELTA DU RHONE ET DE LA COTE MEDITERRANEENNE FRAN-CAISE (PROJET GOLION)]

J. P. Bertrand, M. Guy, and A. LHommer (Bureau de Rech. Geol. et Minieres, Orleans, France) In ESRO European

Earth-Resources Satellite Expts. May 1974 p 121-133 In FRENCH; ENGLISH summary Original contains color illustrations

A number of ERTS-1 images covering the geographical zone were studied and compared with cartographic maps, aerial photographs and thermal-IR images. The old and recent sediments leave traces in the landscape which are decoded by interpreting the shapes of the clear zones forming a network against the black background representing water and humid zones (MSS bands 6 and 7). Fieldwork is currently being conducted to study the corresponding sedimentary models. The comparison of ERTS images, certain thermal-IR images and information on the flow of the Rhone provided some clarification of the mechanisms associated with river dynamics regarding current sedimentation Author (ESRO)

N75-14231 Ecole Nationale Superieure des Mines, Paris (France). Centre d'Informatique Geologique.

COLOUR COMPOSITES OF ERTS MULTISPECTRAL IMAGERY FOR APPLICATIONS IN HYDROLOGY AND **OCEANOGRAPHY**

J. M. Monget In ESRO European Earth-Resources Satellite Expts. May 1974 p 151-153 Original contains color illustrations Work was carried out aimed largely at finding the combination of false colors best suited to the monitoring of suspended matter and chlorophyll patterns. The best results were obtained off the coast of Italy, where evidence of water pollution by oil slicks was also clearly detected. On the North Sea coast, the turbid nearshore surface water distribution has been greatly enhanced, allowing good mapping of sediment transport in this area. The color composites also contain useful information regarding in-land hydrology. For example, the difference between humid and dry vegetation is clearly visible all along the North Sea coast, so that an accurate outline of the polders can be obtained.

Author (ESRO)

N75-14234 Deutsches Hydrographisches Institut, Hamburg (West Germany).

USE OF ERTS-1 IN SEA-ICE STUDIES

K. Struebing In ESRO European Earth-Resources Satellite Expts. May 1974 p 173-178 refs

Arctic and subarctic regions are presented to demonstrate the specific characteristics and advantages of ERTS-1 data. The 60-100 m resolution of the images allows very detailed and precise interpretation and mapping of even the smaller sea ice features. Making use of the differences in reflectance in the visible and near-IR bands, different ice types and development stages can be distinguished. Furthermore, a special relationship between the ice cover and its regional occurrence as shown by ERTS images is emphasized. Author (ESRO)

N75-14262# Washington Univ., Seattle. Arctic Ice Dynamics Joint Experiment Office. **AIDJEX BULLETIN NUMBER 25**

Jul. 1974 126 p refs

(Contract NSF C-625)

(PB-235298/7; AIDJEX-74-25) Avail: NTIS HC \$5.75

A variety of articles dealing with field work and data analysis well as pack ice modeling is presented. Articles discuss: suitability of the pack ice surface to vehicle travel; thickness and roughness variations of multiyear sea ice; determining dynamics and morphology of Beaufort Sea ice from data obtained by satellites, aircraft, and drifting stations; a climate simulation model applied to arctic pack ice; and problems of structuring large capacity computer programs. An index to all ERTS-1 photographs that are presently in the AIDJEX data bank is included. GRA

05 OCEANOGRAPHY AND MARINE RESOURCES

N75-14793*# Kansas Univ. Center for Research, Inc., Lawrence. Remote Sensing Lab.

PRELIMINARY ANALYSIS OF SKYLAB RADSCAT RESULTS OVER THE OCEAN

R. K. Moore, J. D. Claassen, J. D. Young, W. J. Pierson, Jr. (City Univ. of New York), and V. J. Cardone (City Univ. of New York) Nov. 1974 9 p refs (Contract NAS9-13642)

(NASA-CR-141454; RSL-TR-254-2; CUNY-43) Avail: NTIS

HC \$3.25 CSCL 22A

Preliminary observations at 13.9 GHz of the radar backscatter and microwave emission from the sea were analyzed using data obtained by the radiometer scatterometer on Skylab. Results indicate approximately a square-law relationship between differential scattering coefficient and windspeed at angles of 40 deg to 50 deg, after correction for directional effect, over a range from about 4 up to about 25 meters/sec. The brightness temperature response was also observed, and considerable success was achieved in correcting it for atmospheric attenuation and emission. Measurements were made in June, 1973, over Hurricane Ava off the west coast of Mexico and over relatively calm conditions in the Gulf of Mexico and Caribbean Sea.

Author

N75-16116*# National Oceanic and Atmospheric Administration, Miami, Fla. Atlantic Oceanographic and Meteorological Labs. REMOTE SENSING OF OCEAN CURRENT BOUNDARY LAYER Monthly Progress Report, Dec. 1974

George A. Maul, Principal Investigator 9 Jan. 1975 2 p

(NASA Order T-4713-B)

(E75-10098; NASA-CR-141294) Avail: NTIS HC \$3.25 CSCL 08J

٠s

· 06 HYDROLOGY AND WATER MANAGEMENT

06 HYDROLOGY AND WATER MANAGEMENT

Includes snow cover and water runoff in rivers and glaciers, saline intrusion, drainage analysis, geomorphology of river basins, land uses, and estuarine studies.

A75-12873 * Water depth penetration film test. H. E. Lockwood, L. Perry, G. E. Sauer (Technicolor Graphics Services, Inc.), and N. T. Lamar (NASA, Johnson Space Center, Houston, Tex.). *Photogrammetric Engineering*, vol. 40, Nov. 1974, p. 1303-1314. Contract No. NAS9-11500. NASA Order HT-73.

As part of the National Aeronautics and Space Administration Earth Resources Program, a comparative and controlled evaluation of nine film-filter combinations was completed to establish the relative effectiveness in recording water subsurface detail if exposed from an aerial platform over a typical water body. The films tested, with one exception, were those which prior was suggested had potential. These included an experimental 2-layer positive color film, a 2-layer (minus blue layer) film, a normal 3-layer color film, a panchromatic black-and-white film, and a black-and-white infrared film. Selective filtration was used with all films.

A75-13578 Can spinoffs from space provide water for a thirsty world. L. Virden (Brevard Community College, Fla.). In: Technology today for tomorrow; Proceedings of the Eleventh Space Congress, Cocoa Beach, Fla., April 17-19, 1974. Volume 1.

Cape Canaveral, Fla., Canaveral Council of Technical Societies, 1974, p. 2-3 to 2-8. 19 refs.

This analysis of the origin of our water supply system illustrates the seriousness and complexity of our present water problems which will require all the technology acquired by space-age scientists and engineers to solve. It is intended to show that present water depends as much on water vapor in the atmosphere as on that below the crust of the earth. Spinoffs from the space program should be put to use to overcome water pollution and develop prototypes for desalting, reclaiming and recycling water. (Author)

A75-13579 Water-management models in Florida from ERTS-1 data. A. L. Higer, E. H. Cordes (U.S. Geological Survey, Miami, Fla.), and A. E. Coker (U.S. Geological Survey, Tampa, Fla.). In: Technology today for tomorrow; Proceedings of the Eleventh Space Congress, Cocoa Beach, Fla., April 17-19, 1974. Volume 1.

Cape Canaveral, Fla., Canaveral Council of Technical Societies, 1974, p. 3-1 to 3-12. 10 refs.

Questions concerning the water supply for southeast Florida are considered along with the ERTS-1 data which can be obtained from this area. The investigation indicates that an operational water-management model for the Everglades basin is feasible. An ecological model is designed to relate, the wildlife in the Shark River Slough to the availability of food and water. A gradual reduction of water levels prior to bird rookery formation, would result in the concentration of fish in the Shark River Slough. Such a concentration of fish could ensure adequate food for successful hatches of several rare and endangered bird species.

G.R.

A75-13580 * Coastal wetlands - Prospects for satellite inventory. R. R. Anderson and V. Carter (American University, Washington, D.C.). In: Technology today for tomorrow; Proceedings of the Eleventh Space Congress, Cocoa Beach, Fla., April 17-19, 1974. Volume 1. Cape Canaveral, Fla., Canaveral Council of Technical Societies, 1974, p. 3-13 to 3-24. Contract No. NAS5-21752.

It is pointed out that coastal wetlands are among the nation's most valuable natural resources. A relatively low cost and moderately accurate method for mapping these areas would, therefore, be very attractive. It appears that such a method could be found by utilizing ERTS-1 data. Two test sites were selected for an intensive study of the feasibility of such an approach. One test area involves a salt marsh complex located at the mouth of the Chincoteague Bay in Virginia. The second area constitutes a near-saline marsh at the mouth of the Nanticoke River, Maryland. The study shows that ERTS-1 digital data provide maximum gray-level resolution for the mapping of wetland species and features.

G.R.

A75-13596 Rainfall estimations from geosynchronous satellite, imagery, C. G. Griffith and W. L. Woodley (NOAA, Experimental Meteorology Laboratory, Coral Gables, Fla.). In: Technology today for tomorrow; Proceedings of the Eleventh Space Congress, Cocoa Beach, Fla., April 17-19, 1974. Volume 2.

Cape Canaveral, Fla., Canaveral Council of Technical Societies, 1974, p. 5-16 to 5-24. Contracts No. NOAA-25-22-0211; No. NOAA-03-3-022-18.

A method to estimate rainfall from visible geosynchronous satellite images is outlined. The two component relationships, derived from ATS-3 and WSR-57 radar data, are discussed. Calculations are made on two days with this method and compared with ground truth rainfall. Satellite estimates on both days are within a factor of two of ground truth. Sources of error in the component relationships are enumerated. Several planned refinements, such as stratification of the data by synoptic condition and origin of convection, are presented. (Author)

A75-16240 * Coastal and estuarine studies with ERTS-1 and Skylab. V. Klemas, D. Bartlett, W. Philpot (Delaware, University, Newark, Del.), R. Rogers, and L. Reed (Bendix Corp., Aerospace Div., Ann Arbor, Mich.). Remote Sensing of Environment, vol. 3, no. 3, 1974, p. 153-174. 13 refs. Contracts No. NAS5-21837; No. NAS1-12304; No. N00014-69-A-0407.

Coastal vegetation, land use, current circulation, water turbidity, and ocean waste dispersion were studied by interpreting ERTS-1 and Skylab imagery with the help of ground truth collected during overpasses. Based on high-contrast targets such as piers and roads, the ERTS-1 multispectral scanner was found to have a resolution of 70-100 m, Skylab's S190A cameras about 20-40 m, and its S190B camera about 10-20 m. Important coastal land-use details can be more readily mapped using Skylab's imagery. On the other hand, the regular 18-day cycle of ERTS-1 allows observation of important man-made and natural changes, and facilitates collection of ground truth. (Author)

A75-17138 # Flood inundation in the southeastern United States from aircraft and satellite imagery. G. K. Moore and G. W. North (U.S. Geological Survey, Bay St. Louis, Miss.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 607-620. 5 refs.

A75-17140 * # A hydrogeomorphic approach to evaluating flood potential in central Texas from orbital and suborbital remote sensing imagery. V. R. Baker, R. K. Holz, and S. D. Hulke (Texas, University, Austin, Tex.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 629-645. 27 refs. Contract No. NAS9-13312.

A75-17160 * # Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites. A. E. Strong, H. G. Stumpf, J. L. Hart, and J. A. Pritchard (NOAA, National Environmental Satellite Service, Hillcrest Heights, Md.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 923-932. 7 refs. NASA Order S-70246-AG.

Two studies are presented that utilize data from the NOAA-2 and ERTS-1 satellites. The studies are concentrated on two summer upwelling episodes in Lake Michigan when considerable contrast was observed in both surface water temperature as observed by NOAA-2 and surface water color as observed by ERTS-1. Physical, biological and chemical processes support the hypothesis that much of the observed 'whitening' is calcium carbonate precipitating as an immediate result of the upwelling. (Author)

A75-17161 # Remote sensing of western Lake Superior. K. Stortz and M. Sydor (Minnesota, University, Duluth, Minn.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 933-937. Grant No. DACW37-74-C-

Correlation of ERTS data with measurements of turbidity and transmittance for water in the Duluth-Superior harbor and the adjacent Lake Superior water is used in production of turbidity maps for the extreme western arm of Lake Superior. Comparison of reflectance of water obtained from ERTS Bands 4, 5, and 6 as a function of suspended solids indicates the possibility of using this data in effluent tracing. Correlation of ground truth with satellite data allows for extension of results obtained from measurements at few key stations to a large area of the lake. The ability to do this is exceedingly important in the environmental studies of large bodies of water, where the dynamic nature of the system cost prohibits sufficient sampling measurements to cover the entire area. (Author)

A75-17163 # The use of remote sensing in limnological studies. C. T. Wezernak (Michigan, Environmental Research Institute, Ann Arbor, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 963-980. 6 refs. NSF Grant No. GI-34809X.

The concept of a remote sensing trophic index for use in lake assessment is discussed and a comparative analysis of field and remote sensing data for a group of lakes is presented. The results demonstrate the potential of remote sensing for use in limnological studies. Results indicate that changes in the optical properties of a lake, as determined by remote sensing, can be used to assess the trophic state of a lake.

(Author)

A75-17164 # Automatic classification of eutrophication of inland lakes from spacecraft data. R. H. Rogers, L. E. Reed, N. J. Shah (Bendix Corp., Aerospace Systems Div., Ann Arbor, Mich.), and V. E. Smith (Cranbrook Institute of Science, Bloomfield Hills, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 981-993.

Spacecraft data and computer techniques can be used to rapidly map and store onto digital tapes watershed land-use information. Software is now available by which this land-use information can be rapidly and economically extracted from the tapes and related to coliform counts and other lake contaminants, e.g., phosphorus. These tools are basic elements for determining those land-use factors and sources of nutrients that accelerate eutrophication in lakes and reservoirs. (Author)

A75-17178 * # Coastal wetlands analysis from ERTS MSS digital data and field spectral measurements. V. Carter (U.S. Geological Survey, Reston, Va.) and J. Schubert (NASA, Goddard Space Flight Center, Greenbelt, Md.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1241-1260. 17 refs.

Classification, delineation and evaluation of coastal wetlands can be made on the basis of major vegetative associations. To produce wetland maps, two vegetation-analysis look-up tables were developed for use in the ERTS ANALYSIS System. These look-up tables are based on Earth Resources Technology Satellite (ERTS) digital values in Multispectral Scanner (MSS) bands 4, 5, and 7 and were developed using seasonal spectral reflectance measurements from field observations. Computer-generated maps at an approximate scale of 1:20,000 were produced for the primary test site, Chincoteague Marsh, Virginia. There is a high degree of accuracy in the identification of wetland features and plant associations. The classification was also tested on other Atlantic coast salt marshes and a brackish marsh in the Chesapeake Bay.

A75-17179 * # California nearshore processes - ERTS 1. D. D. Steller (Geosource International, Inc., Seal Beach, Calif.) and D. M. Pirie (U.S. Army, Corps of Engineers, San Francisco, Calif.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1261-1278. NASA Order S-70257-AG.

The detectability of many nearshore processes from ERTS is made possible due to the suspended sediment present in the coastal waters. From viewing and analyzing the California coastal imagery collected during the last year and a half, the overall current patterns and their changes have become evident. It is now possible to map monthly and seasonal changes that occur throughout the year. The original objectives of detecting currents, sediment transport, estuaries and river discharge have now been expanded to include the use of ERTS information in operational problems of the U.S. Army Corps of Engineers. This incorporates the detected nearshore features into planning and organizing shore protection facilities. (Author)

A75-17180 # Surface currents along the California coast observed on ERTS imagery. P. R. Carlson (U.S. Geological Survey, Menlo Park, Calif.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1279-1288.

A75-17181 * # Correlation of coastal water turbidity and current circulation with ERTS-1 and Skylab imagery. V. Klemas, M. Otley, W. Philpot, C. Wethe (Delaware, University, Newark, Del.), R. Rogers, and N. Shah (Bendix Corp., Aerospace Div., Ann Arbor, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1289-1317. 14 refs. Contracts No. NAS5-21837; No. N00014-69-A-0407.

The article reviews investigations of current circulation patterns, suspended sediment concentration, coastal frontal systems, and waste disposal plumes based on visual interpretation and digital analysis of ERTS-1 and Skylab/EREP imagery. Data on conditions in the Delaware Bay area were obtained from 10 ERTS-1 passes and one Skylab pass, with simultaneous surface and airborne sensing. The current patterns and sediments observed by ERTS-1 correlated well with ground-based observations. Methods are suggested which would make it possible to identify certain pollutants and sediment types from multispectral scanner data.

A.T.S.

A75-17200 # Microwave radiometric characteristics of snow-covered terrain. R. P. Moore and J. O. Hooper (U.S. Naval Weapons Center, China Lake, Calif.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1621-1632. 7 refs.

A very sensitive Ka-band microwave radiometer has been used to map agricultural areas during winter, spring, and summer seasons, and at various altitudes. This radiometer employs a parametric amplifier to achieve its ultrahigh sensitivity. The mapping function is accomplished by rotating parabolic antennas which scan the beam in the cross-track direction. The forward motion of the aircraft provides the other dimension. The area mapped is a strip which has a width four times the altitude of the aircraft. The system is capable of rapid installation and removal. The radiometer will operate automatically over a wide range of speeds and altitudes and is capable of either a Dicke or high-speed mode of operation. The temperature sensitivity and spatial resolution are sufficient to produce detailed images of the terrain. In addition to solving the problem of terrain remote sensing through snow cover, this technique provides a means by which to survey large areas in a short period of time. (Author)

A75-17950 Microwave profiling of snowpack free-water content. W. I. Linlor (NASA, Ames Research Center, Moffett Field, Calif.), M. F. Meier (U.S. Geological Survey, Tacoma, Wash.), and J. L. Smith (U.S. Forest Service, Pacific Southwest Forest and Range Experiment Station, Berkeley, Calif.). In: Advanced concepts and techniques in the study of snow and ice resources. Washington, D.C., National Academy of Sciences, 1974, p. 729-736.

A microwave system is proposed to measure the amount of liquid-phase water in a snowpack operating in the range of 1 to 10 GHz. Attenuation of the beam between source and receivers is produced by the water in the snow. The relationships of frequency, distance, and volume percent of water are calculated for an assumed detector sensitivity, together with the estimated cost of a representative system. A laboratory test is described that shows the attenuation for snow at maximum wetness at 9.35 GHz. A configuration is proposed that involves a vertical tube containing microwave and radioactive sources and another vertical tube containing microwave and gamma-ray detectors, so that density and wetness profiles are obtained simultaneously over essentially the same path. (Author)

A75-18067 Monitoring trophic status of inland lakes - A quantitative application of ERTS imagery. F. L. Scarpace (Wisconsin, University, Madison, Wis.) and R. E. Wade (Wisconsin Department of Natural Resources, Madison, Wis.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings.

Church, Va., American Society of Photogrammetry, 1974, p. 232-239.

N75-10539*# Geological Survey, Tacoma, Wash.
EVALUATE ERTS IMAGERY FOR MAPPING AND DETECTION OF CHANGES OF SNOWCOVER ON LAND AND ON GLACIERS Semiannual Progress Report, 1 Jan. - 30 Jun. 1973.

Mark F. Meier, Principal Investigator 1 Jul. 1973 7 p ERTS (NASA Order S-70243-AG)

(E74-10812; NASA-CR-140569) Avail: NTIS HC \$3.25 CSCL 08L

The author has identified the following significant results. A new procedure to determine snowcovered areas has been devised. Aside from problems in heavily forested areas this method shows promise in predicting snowmelt runoff from mountain areas and will also assist in energy balance modeling of large snowfields. Snowcover results compare favorably with measurements made by high altitude aircraft photography. Changes in snowcover in areas as small as 3 x 5 km can be determined from ERTS-1, images by both optical and electronic methods. Snowcover changes determined by these two methods in the experimental South Cascade Glacier Basin were verified by field mapping.

Image enahancement techniques on ERTS-1 images of large Alaskan glaciers (the Hubbard, Yentna, and Kahiltna) have given new insights into the large-scale structures and flow dynamics of these potentially hazardous glaciers. The Hubbard Glacier, in particular, is one which poses a threat to man and should be monitored for future changes.

N75-10540*# Geological Survey, Miami, Fla.
SURFACE WATER MODELING EVERGLADES WATER
BASIN, FLORIDA Bimonthly Progress Report, 1 May - 30
Jun. 1974

Aaron L. Higer, Principal Investigator, E. T. Wimberly, E. H. Cordes, and A. E. Coker Jul. 1974 4 p ERTS (NASA Order S-70243-AG-7)

(E74-10814; NASA-CR-140616) Avail: NTIS HC \$3.25 CSCL 08H

N75-10541*# Geological Survey, Reston, Va.
DYNAMICS OF SUSPENDED SEDIMENT PLUMES IN LAKE
ONTARIO Semiannual Progress Report, 1 Jan. - 30 Jun.
1974

Edward J. Pluhowski, Principal Investigator 1 Jul. 1974 4 p ERTS

(NASA Order S-70243-AG-2)

(E74-10815: NASA-CR-140617) Avail: NTIS HC \$3.25 CSCL 08H

The author has identified the following significant results. Although turbidity plumes in Lake Ontario are usually not visible during the winter, meteorologic and hydrologic events may combine to ensure their detection. The clearly defined Niagara River plume of January 25, 1974, was the result of turbid water entering the river at its source near the eastern end of Lake Erie. A persistent southwest wind mild temperature resulted in a pile-up of ice free but turbid water at the source of the Niagara River where the highly colored water entered the river. Upon discharge into Lake Ontario, the Niagara River water appears several shades lighter in tone than the ambient lake water. On February 12, 1974, eastward moving ice floes along the Ontario shoreline were forced to move around the hydraulic barrier created by the Niagara River jet. As a result the Niagara River plume was clearly portrayed by a halo-like band of slush ice borne by wind-driven nearshore currents.

N75-10542*# American Univ., Washington, D.C. Dept. of Biology.

SKYLAB-EREP INVESTIGATIONS OF WETLANDS ECOLOGY Bimonthly Progress Report, 6 Aug. - 6 Oct. 1974

Richard R. Anderson, Principal Investigators, Virginia P. Carter, and Linda J. Alsid 6 Oct. 1974 4 p EREP (Contract NAS1-12209)

(E74-10816; NASA-CR-140618) Avail: NTIS HC \$3.25 CSCL

N75-11409*# Department of the Environment, Ottawa (Ontario).
Applied Hydrology Div.

WATER SURVEY OF CANADA: APPLICATION FOR USE OF ERTS-A FOR RETRANSMISSION OF WATER RESOURCES DATA Progress Report, 1 Apr. - 30 Sep. 1974 R. A. Halliday, Principal Investigator and I. A. Reid Oct. 1974 6 p. Sponsored by NASA ERTS

(E74-10805; NASA-CR-140499) Avail: NTIS HC \$3.25 CSCL 08H

The author has identified the following significant results. The fact that water resources data can be retransmitted from remote areas of Canada by polar orbiting spacecraft to users in population centers on a near real time basis reliably, accurately, and at relative low cost continues to be demonstrated. Over 60,000 transmissions from the nine data collection platforms installed at Water Survey of Canada gauging stations have been received. The stage and ice-out data retransmitted via ERTS-1 have been plotted on a chart record produced by a water stage

servo-manometer installed on the Albany River. The stage increased smoothly until shortly after noon on May 19, 1974. During this time the indicator showed that the ice surface was intact. The stage then dropped sharply and the indicator read that the ice was out. The erratic chart trace after that was consistent with the assumption that the ice surface had broken up and that some short duration jams of broken ice were occurring.

N75-12401*# Geological Survey, Bay Saint Louis, Miss.
HYDROLOGIC SIGNIFICANCE OF LINEAMENTS IN
CENTRAL TENNESSEE Quarterly Progress Report, 1 Jun. 31 Aug. 1974

Gerald K. Moore and Este F. Hollyday, Principal Investigators 31 Aug. 1974 6 p EREP

(NASA Order H-2810-B)

(E75-10030; NASA-CR-140741) Avail: NTIS HC:\$3.25 CSCL

N75-12408*# Environmental Research Inst. of Michigan, Ann Arbor.

A SKYLAB PROGRAM FOR THE INTERNATIONAL HYDRO-LOGICAL DECADE (IHD) Quarterly Report, Dec. 1973 -Feb. 1974

Fabian C. Polcyn and Thomas W. Wagner, Principal Investigators 18 Nov. 1974 3 p EREP (Contract NAS9-13275)

(E75-10039; NASA-CR-140776; ERIM-102300-8-L) Avail: NTIS HC \$3.25 CSCL 08H

N75-13342*# Department of the Environment, Ottawa (Ontario).

Applied Hydrology Div.
WATER SURVEY OF CANADA: APPLICATION FOR USE
OF ERTS-A FOR RETRANSMISSION OF WATER RESOURCES DATA Progress Report, Jun. 1973 - Jul. 1974
R. A. Halliday, Principal Investigator and fr. A. Reid Oct. 1974
10 p Sponsored by NASA ERTS
(E75-10026; NASA-CR-140739) Avail: NTIS HC \$3.25 CSCL

The author has identified the following significant results. Water resources data were retransmitted from nine data collection platforms (DCP) located in remote regions of Canada. The DCPs located in the Arctic operated in temperatures lower than -40 C and the DCP antennas have survived wind speeds of greater than 80 kph and snow loads of a depth of one metre. Ice-out indicators were installed at a few DCP sites. The purpose of these indicators was to enable the detection of the movement of ice out of river channel during spring break-up. The suitability of satellite retransmission as a means of obtaining data from remote areas of Canada continues to be demonstrated. A modest expansion of the DCP network is planned.

N75-13344*# National Aeronautics and Space Administration.
Mississippi Test Facility, Bay Saint Louis.

EVALUATION OF SATELLITE REMOTE SENSING AND AUTOMATIC DATA TECHNIQUES FOR CHARACTERIZATION OF WETLANDS AND COASTAL MARSHLANDS Final Report, Jul. 1972 - Jun. 1974

Robert H. Cartmill, Principal Investigator 29 Jul. 1974 139 p refs Original contains color illustrations. Original contains color imagery. Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue, Sioux Falls, S. D. 57198 ERTS

(E75-10028; NASA-TM-X-72063; Rept-128) Avail: NTIS HC \$5.75 CSCL 05B

The author has identified the following significant results. The evaluation was conducted in a humid swamp and marsh area of southern Louisiana. ERTS digital multispectral scanner data was compared with similar data gathered by intermediate altitude aircraft. Automatic data processing was applied to several data sets to produce simulated color infrared images, analysis.

of single bands, thematic maps, and surface classifications. These products were used to determine the effectiveness of satellites to monitor accretion of land, locate aquatic plants, determine water characteristics, and identify marsh and forest species. The results show that to some extent all of these can be done with satellite data. It is most effective for monitoring accretion and least effective in locating aquatic plants. The data sets used show that the ERTS data is superior in mapping quality and accuracy to the aircraft data. However, in some applications requiring high resolution or maximum use of intermittent clear weather conditions, data gathering by aircraft is preferable. Data processing costs for equivalent areas are about three times greater for aircraft data than ERTS data. This is primarily because of the larger volume of data generated by the high resolution aircraft system.

to, the second

N75-13348*# Environmental Research Inst. of Michigan, Ann Arbor.

SKYLAB WATER DEPTH DETERMINATION Quarterly Progress Report, 1 Sep. - 30 Nov. 1973
F. C. Polcyn and D. R. Lyzenga, Principal Investigators 25 Nov.

F. C. Polcyn and D. R. Lyzenga, Principal Investigators 25 Nov. 1974 3 p EREP

(Contract NAS9-13278)

(E75-10048; NASA-CR-140950; ERIM-102100-11-L) Avail: NTIS HC \$3.25 CSCL 08J

N75-13350*# Agricultural Research Service, Weslaco, Tex. IRRIGATION SCHEDULING, FREEZE WARNING AND SOIL SALINITY DETECTING Monthly Progress Report, Nov. 1974

Craig L. Wiegand, Principal Investigator Nov. 1974 4 p EREP

(NASA Order T-4105-B)

(E75-10050; NASA-CR-140952; MPR-11) Avail: NTIS HC \$3.25 CSCL 02C

N75-13353*# Department of the Environment, Ottawa (Ontario). Applied Hydrology Div.

WATER SURVEY OF CANADA: APPLICATION FOR USE OF ERTS-A FOR RETRANSMISSION OF WATER RESOURCES DATA Final Report, Jul. 1972 - Jun. 1974

R. A. Halliday, Principal Investigator and I. A. Reid Aug. 1974 27 p refs Sponsored by NASA ERTS

(E75-10053; NASA-CR-140959) Avail: NTIS HC \$3.75 CSCL 08H

The author has identified the following significant results. Water resources data including water level, water velocity, precipitation, air temperature, ice condition, DCP battery voltage, and water stage recorder clock operation have been transmitted from remote areas in Canada using the ERTS Data Collection System. The Data Collection Platforms have met all requirements. The suitability of satellite retransmission as a means of obtaining data from remote areas has been demonstrated. The present network of 9 Data Collection Platforms will be expanded to 28 to develop a quasi-operational network.

N75-13365*# Corps of Engineers, Waltham, Mass.

NEW ENGLAND RESERVOIR MANAGEMENT Quarterly

Progress Report, 23 Jul. - 23 Oct. 1974

Saul Cooper and Duwayne Anderson, Principal Investigators

(CRREL) 23 Oct. 1974 4 p EREP

(NASA Order T-4646-B)

(E75-10065: NASA-CR-141005) Avail: NTIS HC \$3.25 CSCL

08H

N75-13371*# National Aeronautics and Space Administration. Lyndon B. Johnson Space Center, Houston, Tex.

THE ERTS-1 INVESTIGATION (ER-600). VOLUME 2: ERTS-1 COASTAL/ESTUARINE ANALYSIS Report for period Jul. 1972 - Jun. 1973

R. Bryan Erb Jul. 1974 286 p refs Original contains color illustrations

(NASA-TM-X-58118; JSC-08457) Avail: NTIS HC \$8.75 CSCL ORJ

The Coastal Analysis Team of the Johnson Space Center conducted a 1-year investigation of ERTS-1 MSS data to determine its usefulness in coastal zone management. Galveston Bay, Texas, was the study area for evaluating both conventional image interpretation and computer-aided techniques. There was limited success in detecting, identifying and measuring areal extent of water bodies, turbidity zones, phytoplankton blooms, salt marshes, grasslands, swamps, and low wetlands using image interpretation techniques. Computer-aided techniques were generally successful in identifying these features. Aerial measurement of salt marshes accuracies ranged from 89 to 99 percent. Overall classification accuracy of all study sites was 89 percent for Level 1 and 75 percent for Level 2.

N75-14194*# Kansas Univ., Lawrence.

SKYLAB STUDY OF WATER QUALITY Progress Report, Sep. - Nov. 1974

Harold L. Yarger, Principal Investigator and James R. McCauley Nov. 1974 7 p EREP

(Contract NAS9-13271)

(E75-10075; NASA-CR-141096) Avail: NTIS HC \$3.25 CSCL 08H

The author has identified the following significant results. Radiometric data derived from S-190A photography appears to correlate reasonably well with suspended solids without the need of rationing the radiances of different bands.

N75-14227 Deutsche Forschungs- und Versuchsanstalt fuer Luft- und Raumfahrt, Oberpfaffenhofen (West Germany). Inst. fuer Satellitenelektronik.

MAPPING OF WATER BODIES IN NORTHERN GERMANY FROM ERTS TAPES

G. Kritikos, B. Sahai, and E. Triendl *In* ESRO European Earth-Resources Satellite Expts. May 1974 p 107-119

Computer-compatible magnetic tapes carrying ERTS MSS data pertaining to northern Germany were examined with the aim of making a classification map of the water bodies in the region. A digital picture processing system was used to reconstruct and process the pictures. The water bodies that are not clearly distinguishable in the individual spectral bands because of unwanted background such as clouds and shadows can be demarcated clearly by taking the difference between intensities in different spectral bands. Using ratio mapping and other techniques, an automatic classification map of the water bodies in the region has been prepared.

Author (ESRO)

N75-14230 Institut Geographique National, Paris (France).
INTERPRETATION OF ERTS-1 DATA FOR INVESTIGATING
THE FRENCH WESTERN COASTAL ZONES (FRALIT
PROGRAM) [L'INTERPRETATION DES DONNEES ERTS-1
POUR L'ETUDE DE LA FACADE OUEST DE LA FRANCE
(PROGRAMME FRALIT)]

P. Demathieu and R. Brossier *In* ESRO European Earth-Resources Satellite Expts. May 1974 p 143-150 ref In FRENCH; ENGLISH summary Original contains color illustrations

Application of ERTS 1 data to a program consisting of a study of the western coastline of France covering both the sea (currents, sedimentology) and the wetlands (sea marshes) is described. The first stage of the work involved trying the maximum number of methods, ranging from conventional photointerpretation to numerical processing. In a second phase, the numerical methods

were extensively developed. The ERTS-1 data proved to be a source of completely new knowledge as regards the dynamic sedimentology of coastal zones. ERTS images can also reveal hitherto completely unknown hydrological phenomena.

Author (ESRO)

N75-14232 Zurich Univ. (Switzerland).
METHODOLOGICAL ASPECTS AND REGIONAL EXAMPLES
OF SNOW-COVER MAPPING FROM ERTS-1 AND EREP
IMAGERY OF THE SWISS ALPS

H. Haefner (Swiss Federal Inst. of Technol.) and K. Seidel In ESRO European Earth-Resources Satellite Expts. May 1974 p 155-164 refs Original contains color illustrations

A quick-look method of mapping snow cover and measuring the areal extent of its surface was developed, allowing rapid interpretation of each image. The method combines the advantages of an automatic photo-quantier with high-resolution photographic techniques for a rapid determination of the variation in the snow cover and its boundaries. The method is demonstrated for a test area of about 170 sq km in the southeastern Swiss Alps and with vertical differences of about 2000 m. The results are combined and compared with ground information. The method could also be applied to data from a source other than ERTS.

Author (ESRO)

N75-14233 Norwegian Water Resources and Electricity Board, Oslo.

ERTS-1 IMAGES IN GLACIOLOGY AND HYDROLOGY: NORWEGIAN EXPERIENCE

G. Ostrem In ESRO European Earth-Resources Satellite Expts. [May 1974] (date] p 165-172 Original contains color illustrations

In order to determine glaciological mass balance, use was made of aerial photographs and ERTS images. By projecting enlarged ERTS images onto topographic maps, the height of the transient snow line can be determined (particularly from MSS 6 and 7) and a good estimate can be obtained of the glacier mass balance over large areas. Both color-composite techniques and image enhancement with contour film have been tried. Experiments with photographic density slicing of ERTS images have shown that the areal extent of snow cover can be delineated and areas of very thick snow (with the highest reflectance) can be separated from thin (or wet) snow. Heavily sediment-loaded glacier streams can be distinguished from less sediment-loaded waters with a similar technique. A cost /benefit estimate is presented.

N75-14260# Montana State Univ., Bozeman. Water Resources Research Center.

WATER RESOURCE AND HAZARD PLANNING REPORT FOR THE CLARK FORK RIVER VALLEY ABOVE MISSOULA, MISSOULA COUNTY, MONTANA

Robert J. Wheeler Jul. 1974 78 p refs Sponsored in part by Dept. of Agriculture

(Contract DI-14-31-0001-3826)

(PB-236041 /0: MUJWRRC-51-0WRR-A-085-MONT(1)) Avail: NTIS HC \$4.75 CSCL 08H

A 33-mile reach of the Clark Fork River east of Missoula, Montana, was the subject of a two-year hydrology and flood hazard study. The 100-year flood plain was delineated by a unique method incorporating aerial photographs obtained during a flood of known recurrence interval, and synthesized rating curves developed for several sites in the reach. The maps, which were developed at a cost of \$120 per river mile, are believed to be nearly as accurate as maps obtained by more sophisticated and costly techniques. Water levels in 20 wells were monitored during a one-year period to determine piezometric surfaces and water table fluctuations. Gravity traverses were made to determine bedrock configurations and depths to bedrock.

N75-15108*# Delaware Univ., Newark. Coll. of Marine Studies.

MONITORING PHYSICAL AND CHEMICAL PARAMETERS

OF DELAWARE BAY WATERS WITH AN ERTS-1 DATA COLLECTION PLATFORM Report on Significant Results

V. Klemas, Principal Investigator and C. Wethe 6 Jan. 1975 2 p ERTS

(Contract NAS5-21837)

(E75-10090; NASA-CR-141154) Avail: NTIS HC \$3.25 CSCL 08J

The author has identified the following significant results. Results of the analysis of data collected during the summer of 1974 demonstrate that the ERTS Data Collection Platform (DCP) is quite responsive to changing water parameters and that this information can be successfully transmitted under all weather conditions. The monitoring of on-site probe outputs reveals a rapid response to changing water temperature, salinity, and turbidity conditions on incoming tides as the tidal salt wedge passes the probe location. The changes in water properties were corroborated by simultaneously sampling the water for subsequent laboratory analysis. Fluctuations observed in the values of salinity, conductivity, temperature and water depth over short time intervals were extremely small. Due to the nature of the probe, 10% to 20% fluctuations were observed in the turbidity values. The use of the average of the values observed during an overpass provided acceptable results. Good quality data was obtained from the satellite on each overpass regardless of weather conditions. Continued use of the DCP will help provide an indication of the accuracy of the probes and transmission system during long term use.

N75-15111*# Environmental Research Inst. of Michigan, Ann Arbor

A SKYLAB PROGRAM FOR THE INTERNATIONAL HYDRO-LOGICAL DECADE (IHD) Quarterly Report, Jun. - Aug. 1974

Fabian C. Polcyn, Principal Investigator and Diana L. Rebel 9 Dec. 1974 $\,$ 3 $\,$ p $\,$ EREP

(Contract NAS9-13275)

(E75-10093; NASA-CR-141157; ERIM-102300-13-L) Avail: NTIS HC \$3.25 CSCL 08H

The author has identified the following significant results. The development of the algorithm (using real data) relating red and IR reflectance to surface soil moisture over regions of variable vegetation cover will enable remote sensing to make direct inputs into determination of this important hydrologic parameter.

N75-15112*# Environmental Research Inst. of Michigan, Ann Arbor.

A SKYLAB PROGRAM FOR THE INTERNATIONAL HYDRO-LOGICAL DECADE (IHD) Quarterly Report, Mar. - May 1974

Fabian C. Polcyn, Principal Investigator and Diana L. Rebel 9 Dec. 1974 3 p EREP

(Contract NAS9-13275)

(E75-10094; NASA-CR-141057; ERIM-102300-12-L) Avail:

NTIS HC \$3.25 CSCL 08H

O7 DATA PROCESSING AND DISTRIBUTION SYSTEMS

Includes film processing, computer technology, satellite and aircraft hardware, and imagery.

A75-10435 Automated image processing in forestry. D. V. Smith. (American Society of Photogrammetry, Annual Convention, St. Louis, Mo., Mar. 11, 1974.) Photogrammetric Engineering, vol. 40, Oct. 1974, p. 1187-1190.

Investigated are methods of automated image processing in natural resources. The interpretation phase and the mapping phase are recognized in the developed automated techniques for mapping forest resources. An automated digital mapping system is being developed to form a part of a resource information system in combination with certain automated interpretation methods. Specific information may be extracted from color and color-infrared films rather than using multiband photography. Multiband photographic equipment is expensive, but the system offers flexibility in terms of the results obtained in each of the spectral bands.

T.S.

A75-11274 Satellites and computer in the service of meteorology: WWW - The global weather observation system (Satelliten und Computer im Dienst der Meteorologie: WWW - Das globale Wetterbeobachtungs-System). R. F. Gebhart (World Meteorological Organization, Education Training and Research Dept., Geneva, Switzerland). VDI-Z, vol. 116, no. 14, Oct. 1974, p. 1157-1159. In German.

A plan for a new global weather observation system, the World Weather Watch (WWW), was developed by the World Meteorological Organization. The WWW project is to be stepwise implemented. The global observation system of WWW is to collect the meteorological data from all areas of the earth which are needed for the daily weather forecasts and for research applications. The observation system makes use of the most modern technological developments including novel radar systems and satellites. The processing of the data in an integrated system of meteorological centers utilizing modern computing technology is also discussed.

G.R.

A75-11419

Analytical estimates of precision in determining parameters of motion of artificial earth satellite when analyzing imprecise measurements with correlated errors. L. F. Porfir'ev and V. V. Smirnov. (Kosmicheskie Issledovaniia, vol. 12, Mar.-Apr. 1974, p. 299-301.) Cosmic Research, vol. 12, no. 2, Sept. 1974, p. 270-272. Translation.

A75-11822 • The community satellite. III. Spaceflight, vol. 16, Nov. 1974, p. 423-426, 440.

A number of technology experiments conducted with the satellite are discussed, giving attention to a very high resolution radiometer experiment, a radio frequency interference experiment, a millimeter wave propagation experiment, a COMSAT propagation experiment, a cesium bombardment ion engine experiment, an advanced thermal control flight experiment, a spacecraft attitude experiment, a radio beacon experiment, and environmental measurements experiments. Special investigations considered are related to the use of a spacecraft vibration accelerometer, a quartz crystal microbalance contamination monitor, and a television camera. G.R.

A75-13310 # A generator of high-power current pulses of semisinusoidal shape for an aerial variant of the method of transient processes (Generator moshchnykh tokovykh impul'sov polusinusoidal'noi formy dlia aerovarianta metoda perekhodnykh protessov). A. S. Lutsychin and V. A. Mamaev. Otbor i Peredacha Informatsii. no. 38. 1974. p. 83-90. In Russian.

Description of a generator of high-power alternating semisinusoidal current pulses which is constructed on the basis of a bridge-type thyristor inverter. An analysis is made of the processes occurring in the inverter circuit, and the conditions for obtaining the maximum magnetic dipole moment that can be supplied by this generator are indicated. The proposed generator is shown to have a number of advantages which make it suitable for aerial prospecting by the method of transient processes.

A.B.K.

A75-13526 A camera for photography of atmospheric particles from aircraft. T. W. Cannon (National Center for Atmospheric Research, Boulder, Colo.). Review of Scientific Instruments, vol. 45, Nov. 1974, p. 1448-1455. 12 refs. NSF-sponsored research.

A75-13741 # Geoscientific research with rockets and satellites. J. Bodechtel (Zentralstelle für Geo-Photogrammetrie und Fernerkundung, Munich, West Germany). International Astronautical Federation, International Astronautical Congress, 25th, Amsterdam, Netherlands, Sept. 30-Oct. 5, 1974, Paper 74-116. 10 p.

The Gemini and Apollo photographic programs showed the significant advantages of using spaceborne remote sensing in earth sciences. Satellites offer continuous repetition rate, long-term observation, and large area coverage. Skylab showed the importance of manned, predominantly experimental, space missions and their restricted use for operational monitoring. Spacelab, on experimental missions, will serve as an optimum sensor platform to define operational models. Earth-observation satellites are supported by airborne and ground-based observation and analysis.

A.T.S.

A75-13758 # Orientation of methods of information processing for the control of terrestrial resources (Orientation des méthodes de traitement de l'information pour le contrôle des ressources terrestres). J. L. Bessis (Centre National d'Etudes Spatiales, Toulouse, France). International Astronautical Federation, International Astronautical Congress, 25th, Amsterdam, Netherlands, Sept. 30-Oct. 5, 1974, Paper 74-140. 16 p. 5 refs. In French.

The Mathematical and Processing Division participates in the development of the coherent program put in place by the National Center of Space Studies since 1970 in a new technique named 'teledetection'. In the European context this program is characterized by a plurality of missions of restricted dimension well adapted to the multidisciplinary character of its applications. Whatever the type of detector utilized, it receives the radiation provided by the ground in one or several ranges of wavelength, transforms it into an electric signal, and memorizes the information. The data issued from the scanning systems (radiometer) or on-board cameras are only considered, with the data from the on-board cameras stored on an analog band or in an on-board memory. F.R.L.

A75-13760 # Comparison of existing earth resources data processing facilities. G. P. Fishman (General Electric Co., Philadelphia, Pa.). International Astronautical Federation, International Astronautical Congress, 25th, Amsterdam, Netherlands, Sept. 30-Oct. 5, 1974, Paper 74-144. 16 p.

The paper provides readers potentially interested in establishing their own capability for accessing ERTS data, with an overview of the ground systems currently in operation in the U.S., Canada, and Brazil, and under active development in Italy and the Federal Republic of Germany. The reader is cautioned to look upon these facilities as examples only. There are many ways to implement ERTS

07 DATA PROCESSING AND DISTRIBUTION SYSTEMS

ground systems ranging from small, relatively inexpensive options to large, complex, and costly implementations. Each implementation plan should be carefully analyzed to fit the needs and budget of the prospective user.

F.R.L.

A75-16476 * Ground-based determination of atmospheric radiance for correction of ERTS-1 data. K. Peacock (Bendix Corp., Aerospace Systems Div., Ann Arbor, Mich.). Applied Optics, vol. 13, Dec. 1974, p. 2741, 2742. Contract No. NAS5-21863.

A technique is described for estimating the atmospheric radiance observed by a downward sensor (ERTS) using ground-based measurements. A formula is obtained for the sky radiance at the time of the ERTS overpass from the radiometric measurement of the sky radiance made at a particular solar zenith angle and air mass. A graph illustrates ground-based sky radiance measurements as a function of the scattering angle for a range of solar air masses. Typical values for sky radiance at a solar zenith angle of 48 degrees are given.

T.S.

A75-16569 Transformation of terrestrial survey data to Doppler satellite datum. R. J. Anderle (U.S. Naval Material Command, Naval Weapons Laboratory, Dahlgren, Va.). (International Association of Geodesy, Symposium on Computation Methods in Geometrical Geodesy, Oxford, England, Sept. 1973.) Journal of Geophysical Research, vol. 79, Dec. 10, 1974, p. 5319-5331. 7 refs.

Doppler satellite observations have been used to determine the positions of 37 sites that were part of a network occupied by BC-4 cameras. Analysis of the Doppler observations by fitting orbits to 48-hour spans of data yielded positions for the sites that are consistent with the results of geometric analysis of the camera observations. Analysis of the data for the 37 sites provided shifts bringing 26 local datums to a world geodetic system. The estimated accuracy of the Doppler datum is about 1.5 m in each component of station position with a probable scale bias of 1 ppm. (Author)

A75-16690 Cloud screening from aerial photography applying coherent optical pattern recognition techniques. G. E. Lukes (U.S. Army, Engineering Topographic Laboratories, Fort Belvoir, Va.). In: Coherent optics in mapping; Proceedings of the Tutorial Seminar and Technology Utilization Program, Rochester, N.Y., March 27-29, 1974. Palos Verdes Estates, Calif., Society of Photo-optical Instrumentation Engineers, 1974, p. 265-272. 8 refs.

A75-17110 * # Classification and mapping of coal refuse, vegetative cover types, and forest types by digital processing ERTS-1 data. F. Y. Borden, B. F. Merembeck, D. N. Thompson, B. J. Turner, and D. L. Williams (Pennsylvania State University, University Park, Pa.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 133-152. 9 refs. Research supported by the Hatch Fund; Contract No. NAS5-23133.

A75-17115 * # Feasibility of using multiplexed SLAR imagery for water resources management and mapping vegetation communities. B. Drake and R. A. Shuchman (Michigan, Environmental Research Institute, Ann Arbor, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 219-249. 6 refs. Contract No. NAS10-8333.

A two-wavelength (X band and L band) multiplexed synthetic aperture side-looking airborne radar (SLAR), providing parallel- and cross-polarized images, has been tested for application in mapping vegetation and water resources. Indications of the relative heights, densities, surface roughness and other parameters provided by the

multiplexed radar imagery can be used to differentiate and map various types of vegetation. The multiplexed SLAR is superior to thermal IR imagery and aerial photography for determining heights of vegetation and water-land boundaries.

A.T.S.

A75-17116 * # Four channel simultaneous X-L band imaging SAR radar. R. Rawson and F. Smith (Michigan, Environmental Research Institute, Ann Arbor, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 251-270. Contracts No. NAS9-12967; No. NAS9-11036.

Design considerations, construction, and testing of a fourchannel microwave radar imaging system are described. Various components of the system are described in some detail including the receiver, transmitter, recorders, and the general timing circuitry. Results from flight tests of the system are described. The system is unique in that images from four microwave channels are provided and the data is obtained simultaneously. Energy is radiated at both 3 cm and 25 cm wavelengths; the four receiving channels are 3 cm, like and crossed polarization and 25 cm, like and crossed polarization.

(Author

A75-17117 # Extraction of urban land cover data from multiplexed synthetic aperture radar imagery. M. L. Bryan (Michigan, Environmental Research Institute, Ann Arbor, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 271-288. 20 refs.

The dual-frequency, dual-polarized side-looking airborne radar (SLAR) was used to collect land-use data in two separate areas in metropolitan Detroit. The data used consisted of a set of vertical panchromatic aerial photographs, 1:24,000 topographic maps, and SLAR data. Results showed that (1) radar has the potential to provide morphological and spatial data in urban areas, (2) with the available four images from dual-band, dual-polarized SLAR the exactness of the positive identification of the land-use groupings are, in a decreasing order - residential, industrial, open and other, transportation and utilities, commercial and services, and institutional, and (3) SLAR may be employed in the study of surface roughness changes relative to meteorological disasters.

T.S.

A75-17132 # On the natural limitations of target differentiation by means of spectral discrimination techniques. M. J. Duggin (Commonwealth Scientific and Industrial Research Organization, Minerals Research Laboratories, North Ryde, New South Wales, Australia). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 499-515.

The work described in this paper has been directed towards determining the minimum differences in detected target radiances necessary to exceed noise at the detector. This is caused by atmospheric fluctuations across the scene and by variations in directional reflectance across the target surface. Only when detected radiance differences exceed this noise can terrain classification be unambiguous. To achieve this aim, measurements have been made of global irradiance in the ERTS bandpasses for Sydney over a period of four months, and spectral global irradiance has been measured for Sydney in the range 400-1100 nm over a period of seven months and at Hobart during the winter solstice. (Author)

A75-17134 # Interactive computer processing for land use planning. E. E. Nelson (McDonnell Douglas Corp., Huntington Beach, Calif.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceed-

ings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 537-547.

A thematic mapping project is being performed featuring multispectral remote sensing, interactive computer analysis, and extensive participation by land use planners in Utah. A test site in the Salt Lake City area was flown in July 1973 with a McDonnell Douglas scanner and cameras. The organizations in Utah have organized in study groups for Land Use, Water Resources, Vegetation-Soils, Geology-Geomorphology, and Imagery. Interactive analysis of the data is being conducted with an XDS-930 computer, facilities for digitizing film or analog magnetic tape, black and white and color output display devices, and an alphanumeric terminal for operator communication with the computer. Computer classification results are being overlayed to standard base maps for final evaluation by the land use planners in Utah.

A75-17135 * # Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota. C. J. Frazee, P. H. Rahn, F. C. Westin, and V. I. Myers (South Dakota State University, Brookings; South Dakota School of Mines, Rapid City, S.D.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 549-568. 7 refs. Grant No. NGL-42-003-007.

A75-17151 # Adaptive processing of MSS data using a decision-directed Kalman filter. R. B. Crane (Michigan, Environmental Research Institute, Ann Arbor, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 781-791.

A decision-directed Kalman filter is shown to be applicable to adaptive processing of multispectral data. Improved recognition accuracy results from the use of the Kalman filter, and the improvement is relatively insensitive to filter parameters; e.g., updating rate. Approximately ten percent additional computer time is required.

(Author)

A75-17154 # Modifications in a computer-implemented method for the detection and extraction of objects in aerial photographs. S. Klausner and D. Karmeli (Technion - Israel Institute of Technology, Haifa, Israel). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 825-836

The K-S filter technique of picture segmentation has been used to detect and extract both man-made objects and terrain types from black and white aerial photographs. Several modifications of the filter are proposed, and their effect on filter performance is discussed.

(Author)

A75-17156 * # SICLOPS - A system of computer programs for rectified mapping of airborne scanner imagery. M. M. Spencer, J. M. Wolf, and M. A. Schall (Michigan, Environmental Research Institute, Ann Arbor, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2.

Mich., Environmental Research Institute of Michigan, 1974, p. 853-864. Contract No. NAS9-9784.

This paper describes SICLOPS, a system of computer programs, which has been developed to perform geometrically corrected line-by-line mapping of airborne multispectral scanner data to ground coordinates and to estimate ground areas. As inputs, SICLOPS accepts aircraft attitude and positional information furnished by ancillary aircraft equipment, plus ground truth information giving the location, in map grid coordinates, of a number of recognizable

landmarks on the ground. The geometric correction and mapping procedure locates the scan lines, or the pixels on each line, in terms of map grid coordinates. The area estimation procedure gives ground area for each pixel or for a predesignated parcel specified in map grid coordinates. The results of exercising the system with simulated data showed the uncorrected video and corrected imagery and produced area estimates accurate to better than 99.7%. Real input data, though prepared, have not yet been processed by the system. (Author)

A75-17191 # Photographs from balloons - Their use in agronomy and management of environment. C. M. Girard-Ganneau and M. C. Girard (Institut National Agronomique Paris-Grignon, Thiverval-Grignon, Yvelines, France). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1467-1474. 7 refs.

The authors have studied photographs taken from stratospheric balloons during flights over the south-west of France. The first step made was to set up a glossary containing all the terms used to describe the landscape units. Afterwards the authors use the measures of rank distances to compare the units. At last some applications of interpretation of photographs from balloons, and of the treatment of the data are given. Two fields of application have been selected: agriculture, and thematic maps in different fields of specialization. (Author)

A75-17193 # Evaluation of ERTS data utilization in developing countries. E. J. Greenblat (ECON, Inc., Princeton, N.J.), D. S. Lowe (Michigan, Environmental Research Institute, Ann Arbor, Mich.), and R. A. Summers (System Planning Corp., Arlington, Va.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1509-1515.

This paper summarizes a study performed by the authors for USAID. ERTS imagery may be utilized by less developed countries for a broad range of applications, many of which appear to have significant economic benefits. The realization of the economic potential will depend in large measure upon the capability for organization of an ERTS applications program. These organizational aspects may be as important for a successful program as the technical capability to interpret the imagery. (Author)

A75-17198 # Deteriorating effects on 3 mm wave passive imagery. G. Schaerer and E. Schanda (Bern, Universität, Berne, Switzerland). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1593-1602.

A high resolution (10 arcminutes) microwave scanning radiometer with a real time display on a TV screen, operating at 3.3 mm wavelength has been used for thermal mapping. Radiometric images of characteristic soil signatures (snow and water) are presented and some special features are discussed. Deteriorated thermograms due to various atmospheric effects are presented. It is shown that the state of the cloud cover means a severe limitation of mapping of snow and water surfaces at 3 mm wavelength.

(Author)

A75-17199 # The usefulness of imaging passive microwave for rural and urban terrain analysis. D. N. Brunelle, J. E. Estes, M. R. Mel, R. R. Thaman, F. E. Evanisko (California, University, Santa Barbara, Calif.), R. P. Moore, C. A. Hawthorne, and J. O. Hooper (U.S. Naval Weapons Center, China Lake, Calif.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1603-1620. 5 refs. Contract No. N00123-73-2352.

In-depth ground truth data have been collected in conjunction with four microwave radiometer (MICRAD) flights. Images were evaluated to determine their information content in studies of the cities of Lost Hills, Coalinga, and Bakersfield, California. Results indicate that passive microwave radiometers can yield accurate information for general land use maps. In addition, the agricultural landscapes around Lost Hills and Coalinga were imaged. In all analyses, emphasis has been placed on problems of target recognition and identification in order to build a background of known MICRAD tone signatures. Preliminary analysis indicates that, while most conventional remote sensing systems have shown considerable utility in terrain analyses, the MICRAD system may add information concerning specific terrain phenomena unavailable from conventional sensor systems. (Author)

A75-17202 * # Electrically scanning microwave radiometers.

R. F. Mix (Aerojet Electrosystems Co., Azusa, Calif.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1649-1655. Contracts No. NAS5-21115; No. NAS5-21635.

The electrically scanning microwave radiometer (ESMR) developed for and currently used onboard the Nimbus 5 meteorological satellite is described, along with the ESMR developed for the Nimbus F satellite. They serve for synoptic mapping of microwave emissions from the earth's surface, the instrument on Nimbus 5 measuring these emissions at a wavelength of 1.55 cm (19.35 GHz) and the instrument on Nimbus F, at a wavelength of 0.81 cm (37 GHz). Radiative transfer characteristics measured at these wavelengths are sufficiently different from IR measurements to permit derivation and interpretation of unique meteorological, geomorphological, and oceanographic data.

A75-17218 # Geothermal reconnaissance from quantitative analysis of thermal infrared images. K. Watson (U.S. Geological Survey, Denver, Colo.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1919-1932. 28 refs.

A75-17219 * # Optical data processing analysis of stream patterns exhibited on ERTS-1 imagery. D. Egbert, J. McCauley, F. Ulaby, and J. McNaughton (University of Kansas Center for Research, Inc., Lawrence, Kan.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1933-1951. 11 refs. Contract No. NAS5-21822.

The purpose of the investigation described in this paper has been the analysis of large-scale geologic ground patterns in Kansas using ERTS-1 imagery and optical data processing techniques. Optical spatial frequency data provide a repeatable quantitative means of specifying ground pattern characteristics. (Author)

A75-18052 American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings. Falls Church, Va., American Society of Photogrammetry, 1974, 460 p. Members, \$2.50; nonmembers, \$5.00.

Topics discussed include the use of a photomapper to produce orthophotographs and height information, a two-level resource-oriented land use classification system for use with remote sensor data, the use of ERTS imagery in regional land use planning, application of the four-color process printing technique to the production of aeronautical maps and charts, a new multilayer mapping system for land use decisions, the use of high-altitude photography and ERTS-1 imagery for mapping wildlife habitats over

extensive areas, the use of ERTS imagery in mapping the trophic level of inland lakes, the use of ERTS-1 imagery in the planning and development of natural resources, a multiple-image processing system to facilitate the extraction of terrain information, the use of small-scale color infrared film for the purpose of soil mapping, and the use of thermal infrared imagery in computerized object identification.

A.B.K.

A75-19268 The application of pattern classification techniques to Skylark earth resources rocket imagery. E. S. Owen-Jones, I. R. Galloway, and B. J. Chandler (Bedford College, London, England). In: Conference on Environmental Sensors and Applications, London, England, November 18, 19, 1974, Proceedings.

London, Institution of Electronic and Radio Engineers, 1974, p. 81-89.

A supervised learning method has been applied to an infra-red false-color image of part of Central Argentina which was obtained by means of a Skylark sounding rocket. The principal field types have been classified, and a separation of the main crop-type, sorghum, into two stages of maturity has been attempted. (Author)

N75-10202# Rhode Island Univ., Kingston. Dept. of Chemistry.

LASER RAMAN SPECTROSCOPY OF SOLUTES DISSOLVED IN WATER FROM A REMOTE PLATFORM Completion Report, 1 Jul. 1971 - 31 Jan. 1974

Christopher W. Brown 31 Jan. 1974 19 p refs (Contract DI-14-31-0001-3641)

(PB-233432/4; W74-09255; OWRR-B-036-RI(4)) Avail: NTIS. HC \$3.00 CSCL 07B

An optical system for obtaining remote Raman spectra of ionic solutes dissolved in water was built and tested. At present, the system has been used to detect and identify 150 ppm of NO3(-)at a distance of 21 ft. Furthermore, the same system has been used to detect oil slicks on the surface of water, and dissolved SO4(-2). RO4(-3), and CO3(-2).

N75-10425# Applied Physics Lab., Johns Hopkins Univ., Silver Soring, Md.

A RADAR ALTIMETER FOR SATELLITE APPLICATION
J. L. MacArthur Oct. 1973, 38 p refs
(Contract NO0017-72-G-4401)

(AD-784373; APL-TG-1226) Avail: NTIS CSCL 14/2

A number of satellite radar altimeter designs have been proposed for the purpose of remotely sensing ocean surface parameters. A precision goal of 10 cm is commonly stated to fulfill the major desires of geodesy and oceanography. An altimeter design approach is discussed that requires only a few watts of peak transmit power, a particular advantage in satellite use. A burst programmed, phase-coded waveform is proposed with an effective compression ratio of 4096, which allows a significant power reduction over previous designs. A design capable of implementation with proven devices is achieved, requiring no new developments. At the same time, flexibility is achieved, and several variations of the basic waveform are discussed with digfering levels of performance, complexity, and power consumption.

N75-10526*# Cornell Univ., Ithaca, N.Y.
EVALUATION OF SKYLAB.IMAGERY AS AN INFORMATION
SERVICE FOR INVESTIGATING LAND USE AND NATURAL
RESOURCES Monthly Progress Report, Jul. 1974

Ernest Hardy, Principal Investigator, James Skaley, and Elmer Phillips 31 Jul. 1974 3 p Original contains color imagery. Original photography may be purchased from the EROS Data Center. 10th and Dakota Avenue, Sioux Falls, S. D. 57198 EREP

(Contract NAS9-13364)

(E74-10730; NASA-CR-139576) Avail: NTIS HC \$3.25 CSCL 08B

N75-10527*# Environmental Research Inst. of Michigan, Ann Arbor.

DEVELOPING PROCESSING TECHNIQUES FOR SKYLAB DATA Monthly Progress Report, Sep. 1974

Richard F. Nalepka and William A. Malila, Principal Investigators .

14 Oct. 1974 5 p EREP (Contract NAS9-13280) (E74-10799; NASA-CR-140493; ERIM-101900-40-L) Avail:

NTIS HC \$3.25 CSCL 05B

N75-10531*# Purdue Univ., Lafayette, Ind. Lab. for Applications of Remote Sensing.

AN INTERDISCIPLINARY ANALYSIS OF MULTISPECTRAL SATELLITE DATA FOR SELECTED COVER TYPES IN THE COLORADO MOUNTAINS, USING AUTOMATIC DATA PROCESSING TECHNIQUES Monthly Progress Report, Sep. 1974

Roger M. Hoffer, Principal Investigator Sep. 1974 5 p EREP (Contract NAS9-13380) (E74-10803; NASA-CR-140497) Avail: NTIS HC \$3.25 CSCL

08F

N75-10570*# National Aeronautics and Space Administration.
Lyndon B. Johnson Space Center, Houston, Tex.

JSC EARTH RESOURCES DATA ANALYSIS CAPABILITIES AVAILABLE TO EOD REVISION B

Jun. 1974 55 p Prepared by Lockheed Electron. Co. (Contract NAS9-12200)

(NASA-TM-X-72029; LEC-3949) Avail: NTIS HC \$4.25 CSCL 05B

A list and summary description of all Johnson Space Center electronic laboratory and photographic laboratory capabilities available to earth resources division personnel for processing earth resources data are provided. The electronic capabilities pertain to those facilities and systems that use electronic and/or photographic products as output. The photographic capabilities pertain to equipment that uses photographic images as input and electronic and/or table summarizes processing steps. A general hardware description is presented for each of the data processing systems, and the titles of computer programs are used to identify the capabilities and data flow.

Author

N75-11419*# Scientific Translation Service, Santa Barbara, Calif. SPACE ICONICS

B. N. Radionov, ed., Ya. L. Ziman, ed., A. A. Izotov, ed., V. I. Pavlov, ed., T. P. Popova, ed., and Yu. M. Chesnokov, ed. Washington NASA Sep. 1974 289 p refs Transl. into I ENGLISH of the publ. "Kosmicheskaya Ikonika" Moscow, Nauka Press, 1973 240 p (Contract, NASw-2483)

(NASA-TT-F-798) Avail: NTIS HC \$8.75 CSCL 14C

A broad range of questions of space iconics and iconology are covered, beginning with methodological questions and the definitions of these terms. The papers on iconology examine the questions of determining spacecraft orientation, techniques and equipment for transforming and correlating television and IR pictures of the earth, geological, geophysical, and geographical interpretation of pictures taken from space, questions of machine interpretation of cloud images obtained from space.

N75-11425* Scientific Translation Service, Santa Barbara, Calif. GEOMETRIC DISTORTIONS OF OPTICOMMERCIAL PANORAMIC TELEVISION SYSTEMS

V. M. Govorov In its Space Iconics (NASA-TT-F-798) Sep. 1974 p 51-59 refs Transl. into ENGLISH of the publ. "Kosmicheskaya Ikonika" Moscow, Nauka Press, 1973 p 44-51 CSCL 09F

One of the problems solvable by the spaceborne television system, topographical surveying, makes high demands on image quality, particularly on the geometric distortions introduced by the television camera. It is the geometric distortions which determine measurement accuracy and, consequently, the possibility of creating reliable planetary surface maps. Comparative analysis of the different television systems capable of solving the problem showed that the requirements on quality of the transmitted images are best satisfied by television cameras with opticomechanical scanning. The design of panoramic television systems and the process of image construction by the opticomechanical camera are discussed. Results indicate that panoramic television cameras have the necessary instrumental accuracy and permit determination of the direction to objects with an error practically equal to the resolution. Author

N75-11453# Army Foreign Science and Technology Center, Charlottesville, Va.

USE OF A DIGITAL COMPUTER TO CONSTRUCT ISOLINE MAPS BASED ON A STEREOMODEL OF TERRAIN

V. M. Zaitsev, V. S. Lavrova, and A. A. Chigirev 18 Apr. 1974 12 p refs Transl. into ENGLISH from Geodeziya i Kartografiya (USSR), no. 4, 1973 p 48-53

(AD-784170: FSTC-HT-23-0276-74) Avail: NTIS CSCL 08/5
A new method of constructing isolines in accordance with an altitude matrix given within the nodes of a grid is proposed. The algorithm for this method was applied in a subsystem including an input device, a control digital computer, and an output device. If a Dnepr-1 computer is used, all the isolines on a map measuring 200 x 300 mm with a 0.2 mm stride can be plotted in about 3 hours, but the time can be shortened by several times if more highly advanced equipment is used.

N75-11602 World Meteorological Organization, Geneva (Switzerland).

METHODS FOR THE AUTOMATIC RECORDING OF ATMOSPHERIC DUST CONCENTRATIONS AND SOME OF THE RESULTS OBTAINED

N. N. Aleksandrov, S. A. Konkov, and G. S. Gunija In its Observation and Meas of Atmospheric Pollution 1974 p 257-264

The photometric and radiometric processes of determining the quantities of atmospheric dust deposited on filters are summarized, and the characteristics of various types of filters are noted. A photometric method utilizing a sampling device which draws air through a tape filter is described, and results are given of gravimetric and photometric analyses of the dust.

ESRO

N75-11504 World Meteorological Organization, Geneva (Switzerland).

AIRCRAFT MEASUREMENTS OF THE CONCENTRATION AND SIZE DISTRIBUTION OF ATMOSPHERIC SULFATE PARTICLES

E. Meszaros In its Observation and Meas. of Atmospheric Pollution 1974 p 276-281 refs

A simple set of aircraft collectors designed to capture aerosol particles in three size ranges for subsequent chemical analysis is described. The preliminary results concerning the size distribution of sulfate particles at different levels of the lower troposphere are also presented.

N75-11551* Akademiya Nauk SSSR, Novosibirsk.
SOLUTION OF THE RADIATIVE TRANSFER THEORY
PROBLEMS BY THE MONTE CARLO METHOD

G. I. Marchuk and G. A. Mikhailov *In* Calif. Univ. Proc. of the UCLA Intern. Conf. on Radiation and Remote Probing of the Atmosphere 1974 p 58-73 refs

CSCL 04A

The Monte Carlo method is used for two types of problems. First, there are interpretation problems of optical observations from meteorological satellites in the short wave part of the spectrum. The sphericity of the atmosphere, the propagation function, and light polarization are considered. Second, problems dealt with the theory of spreading narrow light beams. Direct simulation of light scattering and the mathematical form of medium radiation model representation are discussed, and general integral transfer equations are calculated. The dependent tests method, derivative estimates, and solution to the inverse problem are also considered.

J.A.M.

N75-11571# National Environmental Satellite Service, Washington, D.C.

ENVIRONMENTAL SATELLITE IMAGERY, SEPTEMBER 1974 Key to Meteorological Records Documentation no. 5.4

Sep. 1974 95 p refs Avail: NTIS HC \$4.75

Commence on the NOAA 2 satellite.

Commence on the NOAA 2 satellite.

N75-12279*# ~ckheed Electronics Co., Houston, Tex. Aerospace Systems ...
CORRELATION OF MISSIONS 191, 51M AND HELICOPTER PHOTOGRAPHY

B. A. Bryan and S. H. Tunnel Aug. 1972 43 p (Contract NAS9-12200)

(NASA-CR-134289; EOD2876; TWP-72-004) Avail: NTIS HC \$3.75 CSCL 14E

The data obtained during aerial photography flights using a helicopter are presented. The areas photographed are identified as three areas within Test Site 175: (1) Rosenberg, (2) Houston Ship Channel/Trinity Bay, and (3) the Somerville Dam. Data are presented in the form of charts. Reproductions of the aerial photographs are included.

Author

N75-12409*# Environmental Research Inst. of Michigan, Ann

DEVELOPING PROCESSING TECHNIQUES FOR SKYLAB DATA Monthly Progress Report, Oct. 1974

Richard F. Nalepka and William A. Malila, Principal Investigators 14 Nov. 1974 4 p EREP

(Contract NAS9-13280)

,

(E75-10040; NASA-CR-140777; ERIM-101900-42-L) Avail: NTIS HC \$3.25 CSCL 05B

N75-12412*# Colorado School of Mines, Golden. Dept. of Geology.

GEOLÓGIC AND MINERAL AND WATER RESOURCES INVESTIGATIONS IN WESTERN COLORADO, USING SKYLAB EREP DATA Monthly Progress Report, Oct. 1974 Keenan Lee, Principal Investigator 15 Nov. 1974 4 p EREP (Contract NAS9-13394)

(E75-10043; NASA-CR-140780) Avail: NTIS HC \$3.25 CSCL

N75-12417*# Environmental Research Inst. of Michigan, Ann Arbor. Infrared and Optics Div.

NEW THEORETICAL MODELS AND RATIO IMAGING TECHNIQUES ASSOCIATED WITH THE NASA EARTH RESOURCES SPECTRAL INFORMATION SYSTEM Technical Report, 1 Feb. 1973 - 31 Jan. 1974

R. K. Vincent Jul. 1974 116 p refs

(Contract NAS9-9784)

(NASA-CR-140357; ERIM-190100-30-T) Avail: NTIS HC \$5.25 CSCL 05B

Four independent investigations are reported; in general these are concerned with improving and utilizing the correlation between the physical properties of natural materials as evidenced in laboratory spectra and spectral data collected by multispectral scanners. In one investigation, two theoretical models were devised that permit the calculation of spectral emittance spectra for rock and mineral surfaces of various particle sizes. The simpler of the two models can be used to qualitatively predict the effect of texture on the spectral emittance of rocks and minerals; it is also potentially useful as an aid in predicting the identification of natural atmospheric aerosol constituents. The second investigation determined, via an infrared ratio imaging technique, the best pair of infrared filters for silicate rock-type discrimination. In a third investigation, laboratory spectra of natural materials were compressed into 11-digit ratio codes for use in feature selection, in searches for false alarm candidates, and eventually for use as training sets in completely automatic data processors. In the fourth investigation, general outlines of a ratio preprocessor and an automatic recognition map processor are developed for on-board data processing in the space shuttle era.

N75-12420*# Control Data Corp., Minneapolis, Minn.
CORRELATION AND REGISTRATION OF ERTS MULTISPECTRAL IMAGERY

L. O. Bonrud and P. J. Henrikson Apr. 1974 127 p refs (Contract NAS9-13114)

(NASA-CR-134294) Avail: NTIS HC \$5.25 CSCL 05B

Examples of automatic digital processing demonstrate the feasibility of registering one ERTS multispectral scanner (MSS) image with another obtained on a subsequent orbit, and automatic matching, correlation, and registration of MSS imagery with aerial photography (multisensor correlation) is demonstrated. Excellent correlation was obtained with patch sizes exceeding 16 pixels square. Qualities which lead to effective control point selection are distinctive features, good contrast, and constant feature characteristics. Results of the study indicate that more than 300 degrees of freedom are required to register two standard ERTS-1 MSS frames covering 100 by 100 nautical miles to an accuracy of 0.6 pixel mean radial displacement error. An automatic strip processing technique demonstrates 600 to 1200 degrees of freedom over a quater frame of ERTS imagery. Registration accuracies in the range of 0.3 pixel to 0.5 pixel mean radial error were confirmed by independent error analysis. Accuracies in the range of 0.5 pixel to 1.4 pixel mean radial error were demonstrated by semi-automatic registration over small geographic areas. Author

N75-12422*# Electromagnetic Systems Labs., Sunnyvale, Calif. IMAGE SELECTION SYSTEM

M. A. Knutson, Donald Hurd, Larry Hubble, and Richard M. Kroeck Washington NASA Nov. 1974 98 p ref (Contract NAS2-7064)

(NASA-CR-2475; ESL-TM-456) Avail: NTIS HC \$4.75 CSCL 05B

An image selection (ISS) was developed for the NASA-Ames Research Center Earth Resources Aircraft Project. The ISS is an interactive, graphics oriented, computer retrieval system for aerial imagery. An analysis of user coverage requests and retrieval strategies is presented, followed by a complete system description. Data base structure, retrieval processors, command language, interactive display options, file structures, and the system's capability to manage sets of selected imagery are described. A detailed example of an area coverage request is graphically presented.

N75-13335*# Kansas Univ. Center for Research, Inc., Lawrence. Remote Sensing Lab.

USE OF FEATURE EXTRACTION TECHNIQUES FOR THE TEXTURE AND CONTEXT INFORMATION IN ERTS IMAGERY: SPECTRAL AND TEXTURAL PROCESSING OF ERTS IMAGERY Final Report, 1 Aug. 1972 - 17 Mar. 1974

Robert H. Haralick, Principal Investigator and Robert J. Bosley 17 Mar. 1974 99273 p refs Original contains imagery. Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue, Sioux Falls, S. D. 57198 ERTS (Contract NASS-21822)

(E75-10011; NASA-CR-140655) Avail: NTIS HC \$8.50 CSCL 05B

The author has identified the following significant results. A procedure was developed to extract cross-band textural features from ERTS MSS imagery. Evolving from a single image texture extraction procedure which uses spatial dependence matrices to measure relative co-occurrence of nearest neighbor grey tones, the cross-band texture procedure uses the distribution of neighboring grey tone N-tuple differences to measure the spatial interrelationships, or co-occurrences, of the grey tone N-tuples present in a texture pattern. In both procedures, texture is characterized in such a way as to be invariant under linear grey tone transformations. However, the cross-band procedure complements the single image procedure by extracting texture information and spectral information contained in ERTS multiimages. Classification experiments show that when used alone, without spectral processing, the cross-band texture procedure extracts more information than the single image texture analysis. Results show an improvement in average correct classification from 86.2% to 88.8% for ERTS image no. 1021-16333 with the cross-band texture procedure. However, when used together with spectral features, the single image texture plus spectral features perform better than the cross-band texture plus spectral features, with an average correct classification of 93.8% and 91.6%, respectively.

N75-13336*# Cornell Univ., Ithaca, N.Y. Dept. of Natural

EVALUATION OF SKYLAB IMAGERY AS AN INFORMATION SERVICE FOR INVESTIGATING LAND USE AND NATURAL RESOURCES Progress Report, Oct. 1974

Ernest E. Hardy, Principal Investigator 31 Oct. 1974 12 p EREP

(Contract NAS9-13364)

(E75-10013; NASA-CR-140651) Avail: NTIS HC \$3.25 CSCL 08B

N75-13347*# Nebraska Univ., Lincoln.

USE OF ERTS-1 IMAGERY TO INTERPRET WIND EROSION HAZARD IN THE SANDHILLS OF NEBRASKA

James V. Drew, Principal Investigator; Paul M. Seevers, and David T. Lewis 22 Nov. 1974 16 p refs Original contains imagery. Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue, Sioux Falls, S. D. 57198 ERTS

(Contract NAS5-21756)

(E75-10044; NASA-CR-140781) Avail: NTIS HC \$3.25 CSCL 08F

N75-13367*# Scientific Translation Service, Santa Barbara, Calif. FROM AEROSPACE PHOTOGRAPHS TO FORECASTING AND CALCULATING FLOWS

G. P. Kalinin Washington NASA Dec. 1974 63 p refs Transl. into ENGLISH from the book "Ot Aerokosmicheskikh Snimkov k Prognozam: Raschetam Stoka" Leningrad, Hydrometeorological Press, 1974 p 1-41

(Contract NASw-2483) (NASA-TT-F-16006) Avail: NTIS HC \$4.25 CSCL 08E

A theoretical basis is provided for using hydrological data obtained by remote measurements (from spacecraft and aircraft).

Methods of calculating flow and the characteristics of the snow cover are suggested, based on photographs that characterize the dynamics of the hydrological processes that occur on the surfaces of river basins. Problems are posed of experimental investigations leading to new possibilities of analysis and calculation of a number of hydrological elements.

N75-14187*# Cornell Univ., Ithaca, N.Y. Dept. of Natural Resources.

EVALUATION OF SKYLAB IMAGERY AS AN INFORMATION SERVICE FOR INVESTIGATING LAND USE AND NATURAL RESOURCES Progress Report, Nov. 1974

Ernest E. Hardy, Principal Investigator 30 Nov. 1974 1 p

(Contract NAS9-13364)

(E75-10068; NASA-CR-141022) Avail: NTIS HC \$3.25 CSCL

N75-14189 *# Kansas Univ. Center for Research, Inc., Lawrence. Remote Sensing Lab.

DESIGN DATA COLLECTION WITH SKYLAB/EREP MI-CROWAVE INSTRUMENT S-193 Monthly Letter Progress Report No. 12, Sep. 1974

Arun Sobti, Richard K. Moore, Fawwaz T. Ulaby, John C. Barr, Evan Davison, Chia Sung, Saad Ulaby, and Tom Burton Sep. 1974 4 p EREP

(Contract NAS9-13331)

(E75-10070; NASA-CR-141101) Avail: NTIS HC \$3.25 CSCL 05B

N75-14217 Bochum Observatory (West Germany). Inst., fo Space Research.

ENVIRONMENTAL SATELLITES AND ERTS-1 IMAGERY H. Kaminski and Anne-Marie Martin In ESRO European Earth-Resources Satellite Expts. May 1974 p 1-19 refs Original contains color illustrations

The application of environmental satellite data to complement ERTS-1 imagery is discussed. The first Earth Resources Technology Satellite was conceived as a non-meteorological system with a long period of repetitive coverage (18 days) and sensor bandwidths in the range of visible light and photographic infrared. Since the start of ERTS-1, it has been found necessary to complement this concept with synoptic visible and infrared imagery from environmental satellites, such as Nimbus-3 and NOAA-2. The use of this high resolution imagery is demonstrated for the recognition of thermal phenomena such as warm and cold currents in coastal seas, definition of differing layers in the atmosphere, and delineation of urban hot spots. Available examples of ERTS imagery are compared with the above to illustrate the compatibility of these sensor systems for geoscientific research. **ESRO**

N75-14218 Zentralstelle fuer Geo-Photogrammetrie und Fernerkundung, Munich (West Germany).

EVALUATION OF ERTS-1 DATA BY ANALOGUE AND DIGITAL TECHNIQUES

J. Bodechtel, R. H. Dittel, and R. Haydn *In* ESRO European Earth Resources Satellite Expts. May 1974 p 21-35 Original contains color illustrations

Digital and analog image evaluation was performed on ERTS-1 data representing a Central European summer scene and a North Italian winter scene. A statistical analysis of the data, as first step in the evaluation process, showed strong correlation between both infrared channels for these types of landscape and these seasonal periods. For representative areas south of the Starnberg Lake near Munich, different features such as water, conifer forests, agricultural areas and particular soil category were identified by applying factor analysis. Feature identification was also performed by a digital quick-look method which considers the slope between the intensity levels for the channels of the visible and infrared spectrum caused by different reflection properties of surfaces.

The results obtained digitally are compared with those obtained by analog density slicing. A hardware concept for an interactive hybrid image evaluation system is presented. Author (ESRO)

N75-14219 Research Inst. of National Defence, Stockholm (Sweden).
PROCESSING OF ERTS CCT IMAGERY OF SWEDEN. A
PILOT STUDY

T. Orhaug and S. I. Aekersten In ESRO European Earth-Resources Satellite Expts. May 1974 p 37-45 Original contains color illustrations

Data handling, processing, and presentation problems are discussed in connection with a Swedish remote sensing program aimed at natural resources inventories and environmental monitoring. For want of large quantities of remotely sensed national data suitably recorded for automatic processing and analysis, ERTS-1 CCT's of Swedish scenes were acquired as a means of making an inventory of forest-rejuvenation areas. In addition to standard methods of computer imagery generation, a technique developed in Sweden is used. An ink-jet plotter yields a 3 color image on ordinary paper. Examples of imagery from ERTS-1 recordings are presented.

N75-14220 Sheffield Univ. (England). Dept. of Geography. AN EVALUATION OF STEREOSCOPIC VIEWING OF ERTS AND SKYLAB IMAGES

J. L. VanGenderen *In* ESRO European Earth-Resources Satellite Expts. May 1974 p 47-55 refs Original contains color illustrations

It is shown that stereoscopic examination of orbital data can contribute to studies of spatial, spectral, and temporal variations on the imagery. The combination of true stereo parallax plus shadow parallax offers many possibilities to human interpreters for making meaningful analyses of orbital imagery. Using the procedures of stereoscopic image interpretation described, many earth scientists who have neglected the important information source that ERTS and Skylab data can be, because of their lack of expensive data-analysis equipment, should be encouraged to use the data with simple stereoscopic instruments and the analysis techniques defined.

Author (ESRO)

N75-14221 Hunting Surveys, Ltd., Boreham*Wood (England). THE POTENTIAL OF SMALL-SCALE MAPPING FROM ERTS-1 IMAGERY

H. Chisman In ESRO European Earth-Resources Satellite Expts. May 1974 p 57-59

Comparison tests are to be made between existing mapping, the accuracy and completeness of which can be relied upon, and that obtained directly from ERTS imagery with a minimum of control. By comparing satellite coverage and existing mapping the completeness of detail that can be interpreted, and the locational accuracy of interpreted detail can be examined. The effectiveness of the different spectral band images for the interpretation of different features will be evaluated and also photogrammetric and image enhancement techniques applied to the imagery that may be useful in map printing. If satisfactory results are obtained from these studies, the techniques will be applied to mapping in those parts of the world where present maps are inadequate.

N75-14222 International Training Centre for Aerial Survey, Delft (Netherlands)

THE ITC 8-IMAGE COMPARATOR AND ITS POSSIBLE USE IN ERTS AND EREP INVESTIGATIONS

S. A. Hempenius In ESRO European Earth-Resources Satellite Expts. May 1974 p 61-70 refs

A multi-image projector is described for ERTS and EREP imagery. The single-lamp viewer projects up to eight, three times enlarged, 60 mm by 60 mm transparencies through one objective onto an almost horizontal table-top screen. Used as a color additive viewer, any subset of the eight multispectral images can be presented; the intensity and color of the individual images can be selected independently (red, green, blue, or white). In the flickering mode, the series of single images, or of pairs.

triplets, etc. can be thrown onto the screen in slow or rapid succession (frequency adjustable up to 30 Hz), the time constant of the shutters being about 5 msec. The number of images per cycle (1 to 8), the presentation time per image, the temporal overlap of the images, the color, and the intensity can be set independently. Visual faculties to be explored for discrimination and identification on multispectral and multi-temporal ERTS images include color viewing, detection of flickering fields, and disappearance of fluctuations when passing the critical flicker frequency.

Author (ESRO)

N75-14223 Institut de Geographie, Aix en Provence (France). Centre de Telèdetection.

OPTICAL PROCESSING OF ERTS-1 (MSS) IMAGES: EXAMPLES FOR THE ARCACHON BASIN AND RHONE DELTA

J. Pouquet, J. O. Palgen (Earth Satellite Corp., Washington, D.C.), and P. A. Rey (Toulouse Univ.) In ESRO European Earth-Resources Satellite Expts. May 1974 p 71-74 Original contains color illustrations

Conventional examples showing the possibilities of density slicing or color-composite techniques are presented. It is emphasized, however, that these old procedures should only be used as a first approach; only correct digitization (reflectance, radiance temperature values according to wavelength) is likely to yield valuable scientific results.

Author (ESRO)

N75-14225 Messerschmitt-Boelkow-Blohm G.m.b.H., Munich (West Germany).

REDUCTION OF INFORMATION REDUNDANCY IN ERTS-1 AND EREP DATA. A PROPOSAL FOR AN EXPERIMENTAL STUDY

D. Davidts and A. Loeffler (Ges. fuer Weltraumforsch.) In ESRO European Earth-Resources Satellite Expts. May 1974 p 81-91 refs

An experiment is discussed for measuring the information density of remotely sensed data collected from different altitudes (low-flying aircraft, high-altitude aircraft, Spacelab, ERTS, EREP) and for determining the redundancy with respect to the final output of the interpreter and his information needs and processing capabilities.

Author (ESRO)

N75-14229 Swiss Inst. of Meteorology, Zurich.

METEOROLOGICAL INTERPRETATION OF HIGH RESOLU-TION IMAGES FOR REMOTE SENSING OF EARTH RESOURCES [INTERPRETATION METEOROLOGIQUE DES IMAGES A HAUTE RESOLUTION DESTINEES A LA TELEDETECTION DES RESSOURCES TERRESTRES]

A. Piaget In ESRO European Earth-Resources Satellite Expts.
May 1974 p 135-141 refs in FRENCH; ENGLISH summary

It is shown that remote sensing of earth resources is very valuable to meteorology, because the very high image resolution required for remote sensing (less than 100 m) enables a study to be made of the microstructure of clouds over a relatively wide area. Another application makes use of the fine detail of the temperature distribution field. It is often possible to study the penetration of air from the plain into the valleys, or the stagnation of cold air in the dips in the ground surface. This is useful information for the protection of the environment and for the planting of new crops, to give only two examples. The extent of the snow cover and its effect on the neighboring air can only be appreciated by studying both ordinary and very high resolution images in combination.

Author (ESRO)

N75-14242 Louvain Univ. (Belgium). Inst. of Earth-Sciences. PRELIMINARY EVALUATION OF ERTS-1 IMAGERY OF CENTRAL AFRICA

J. Sterckx and J. DePloey *In ESRO* European Earth Resources Satellite Expts. May 1974 p 289-295 refs

The investigation was conducted using black and white ERTS-1 imagery, both transparencies and contact paper prints. The Mweru Wantipa swamp, situated between Lake Tanganyika and Lake Mweru, appears on the ERTS imagery as an important take, bordered on all sides by fault scarps. Only a few features of human presence could be recognized. The parallelism of the rivers on the Kwango plateau is disturbed by a system of grabens, north of Feshi. These are not marked on the geological map. Remains of a forest on the divides permit an evaluation of human density. A large area around the Itimbiri-Zaire confluence was examined on the ERTS imagery in order to propose a geographical division based on the coincidence and possible relationships of particular physical and human features.

Author (ESRO)

N75-14243 Louvain Univ. (Belgium). PEDOLOGICAL INTERPRETATION OF ERTS-1 IMAGERY OF THE RIMA-SOKOTO RIVER BASIN (N. NIGERIA) R. Gombeer and J. dHoore /n ESRO European Earth Resources Satellite Expts. May 1974 p 297-301 refs

Ground truth data were compared with two adjacent ERTS-1 MSS scenes, covering an area roughly located between 4 deg and 7 deg E and 12 deg and 13 deg 30 min N. These 100% cloudless scenes were recorded one month after the monthly rainfall had dropped sharply. Some variation in plant available ground water and in plant cover could therefore be expected and the reflections of the vegetation, together with those characteristic of soil surfaces under different physical conditions, should give rise to recognizable features. By projection of positive transparencies, meaningful enlargements up to a scale of 1:1000,000 could be obtained. MSS bands 5, 6 and 7 in particularly yielded good pictures of drainage patterns and other geomorphologic features, and excellent correlation was observed with the 1:5000,000 scale geomorphological map of the area by Sombroek and Zonneveld. Author (ESRO)

N75-14249 Overseas Development Administration, Surrey (England). Land Resources Div.

A COMPARISON OF ERTS IMAGERY WITH CONVEN-TIONAL AERIAL PHOTOGRAPHY FOR LAND-RESOURCE SURVEYS IN LESS DEVELOPED COUNTRIES. EXAMPLES FROM THE RIFT VALLEY LAKES BASIN, ETHIOPIA

R. B. King and A. BlairRains In ESRO European Earth Resources Satellite Expts. May 1974 p 371-379 refs

The extent to which ERTS imagery can aid land resource surveys in less developed countries was analyzed, with particular regard to: (1) multiband imagery; (2) recent and repetitive imagery; (3) mapping; (4) relief estimation; (5) detail; (6) synoptic viewing; and (7) color composites. These aspects are compared with the data revealed by panchromatic aerial photography. The analysis was carried out within the framework of a conventional land resource survey on a limited budget, and examples are mainly taken from one such survey within the Rift Valley Lakes Basin, Ethiopia. The main conclusions are the ERTS imagery is a significant aid to map production in areas where the present map coverage is poor, but its scale is too small for any detailed analysis. The synoptic aspect of ERTS imagery, however, is useful, particularly as a false-color composite. The ability to monitor changes in hydrology and agriculture is also advantageous.

Author (ESRO)

N75-14265# Army Foreign Science and Technology Center, Charlottesville Va.

INVESTIGATION OF ACCURACY OF DETERMINATION OF PLANE COORDINATES BY MEANS OF THE CENTRAL SCIENTIFIC RESEARCH INSTITUTE OF GEODESY, AERIAL PHOTOGRAPHY AND CARTOGRAPHY AIRBORNE RANGE-FINDER

I. L. Gill, K. N. Gertsenova, and B. A. Batamanovskii 19 Nov. 1973 13 p refs Transl. into ENGLISH from Geod. Kartografiya (USSR), no. 3, 1972 p 37-44 (AD-785886; FSTC-HT-23-1270-73) Avail: NTIS CSCL 08/5

To answer the question of the possibility of using an airborne range finder (ARR) in topographic surveys on a 1:10,000 scale, and to choose the optimum technology of operations, experimental investigations were carried out on the accuracy of creating a radio-geodetic plane base by a method using the ARR.

N75-14257# Army Foreign Science and Technology Center,

WORK DEALING WITH DENSIFICATION OF ALTIMETRIC-PLANOMETRIC DATA POINTS EMPLOYING UNIVERSAL INSTRUMENTS AND RADIO ALTIMETER AND STATO-**SCOPIC INDICATIONS**

N. L. Borodin 25 Jul. 1974 13 p Transl. into ENGLISH from Geodez, i Kartograf, (Moscow), no. 11, 1969 p 46-52 (AD-785877; FSTC-HT-23-780-72) Avail: NTIS CSCL 08/2

Experimental work is reported on the photogrammetric densification of altimetric-planometric data. The photography studied was on a scale of 1:10,000 with a relief profile of 2.5 meters. The construction of photogrammetric grids, the geodetic orientation, and grid linking are discussed. Evaluation results of the study are tabulated. FOS

N75-15101 *# Purdue Univ., Lafayette, Ind. Lab. for Applications of Remote Sensing.

REFORMATTING SKYLAB AND MSS DATA OVER NORTH-

ERN INDIANA Monthly Report, Nov. 1974 LeRoy F. Silva, Principal Investigator Nov. 1974 1 p EREP (Contract NAS9-13301)

(E75-10016; NASA-CR-140647) Avail: NTIS HC \$3.25 CSCL 05B

N75-15102*# Environmental Research Inst. of Michigan, Ann

DEVELOPING PROCESSING TECHNIQUES FOR SKYLAB DATA Monthly Progress Report, Nov. 1974

Richard F. Nalepka and William A. Malila, Principal Investigators 18 Dec. 1974 5 p ref EREP

(Contract NAS9-13280)

(E75-10082; NASA-CR-141106; ERIM-101900-44-L) NTIS HC \$3.25 CSCL 05B

N75-15127*# Louisiana State Univ., Baton Rouge. Div. of Engineering Research.

THE USE OF COLOR INFRARED IMAGERY FOR THE STUDY OF MARSH BUGGY TRACKS

Charles A. Whitehurst and Linda N. Doiron [1974] 13 p refs Original contains color illustrations

(Grant NGL-19-001-105)

Avail: NTIS HC \$3.25 CSCL 08H

Color infrared imagery is used to determine the location of buggy routes and to quantify the extent of tracks in a selected area where the marsh is seriously dissected. The imagery is used to show successive stages of destruction. It is recommended that alternate routes be identified in the opprating area to eliminate continuous use of the same route and facilitate faster revegetation. Author

Page intentionally left blank

08

INSTRUMENTATION AND SENSORS

Includes data acquisition and camera systems and remote sensors.

A75-10075 Performance comparison of techniques for obtaining stereo radar images. G. L. Bair and G. E. Carlson (Missouri, University, Rolla, Mo.). *IEEE Transactions on Geoscience Electronics*, vol. GE-12, Oct. 1974, p. 114-122. Contract No. N00014-69-A-0141-0008. NR Project 387-069.

Three stereo radar imaging techniques namely, (1) an improved single flight technique, (2) a previously proposed single flight technique, and (3) a two-flight technique, are compared with respect to both theoretical performance and performance in the presence of errors. The theoretical data for the comparison of performance-affecting image parameters have been generated by computer simulation of the imaging geometries involved. The comparison shows that the improved single flight technique yields the best error performance for the assumed set of system errors.

M.V.E.

A75-10434 A technology to renovate the search for new mineral deposits. J. T. Awald (Denver Mineral Exploration Corp., Denver, Colo.). (American Society of Photogrammetry, Annual Convention, St. Louis, Mo., Mar. 11, 1974.) Photogrammetric Engineering, vol. 40, Oct. 1974, p. 1173-1186.

Two commercial applications of remote sensing to the search for mineral deposits are presented. The application to the discovery of aggregate limestone and porphyry copper deposits is illustrated. It is shown that the use of satellite and high-altitude remote sensing data through a wide range of the energy spectrum, coupled with electronic processing, offers new prospects in the exploration of lead, zinc, tungsten, molybdenum, uranium, oil, gas, and water resources.

T.S.

A75-10790 # A satellite view of weather systems over the North American Arctic. N. A. Streten (Alaska, University, Fairbanks, Alaska). Weather, vol. 29, Oct. 1974, p. 369-380. 8 refs. Contract No. NOAA-3-35400.

The polar orbiting NOAA-2 satellite launched in October 1972 carries the Very High Resolution Radiometer (VHRR). The VHRR scans simultaneously in both the visible and the thermal infrared with a resolution at nadir of one kilometer. The VHRR imagery possesses considerable potential in polar meteorology. A number of pictures showing VHRR imagery are presented. The value of complementary visible and infrared imagery at high resolution in a synoptic analysis in the arctic is illustrated with the aid of a few examples.

G.R.

A75-11443 # Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation (Optiko-akusticheskoe detektirovanie malykh kontsentratsii molekul ftoristogo vodoroda, okisi azota i dvuokisi ugleroda v gazakh izlucheniem impul'snogo lazera na ftoristom vodorode). A. S. Gomeniuk, V. P. Zharov, D. D. Ogurok, E. A. Riabov, O. A. Tumanov, and V. O. Shaidurov (Akademiia Nauk SSSR, Institut Spektroskopii, Moscow, USSR). Kvantovaia Elektronika (Moscow), vol. 1, Aug. 1974, p. 1805-1811, 11 refs. In Russian.

A75-11720 The 1973 program of measurement of the minor constituents of the stratosphere using the Concorde 001 (Campagne 1973 de mesures des constituents mineurs dans la stratosphère par Concorde 001). M. Ackerman (Institut d'Aéronomie

Spatiale de Belgique, Brussels, Belgium) and R. Joatton (Société Nationale Industrielle Aérospatiale, Paris, France). L'Aéronautique et l'Astronautique, no. 47, 1974, p. 15-22. 10 refs. In French.

Review of the genesis, definition, and performance of the 1973 program of measurement of the rare (NO, NO2, SO2, CH4, HCHO, and HNO3) constituents of the stratosphere using infrared spectroscopy on 12 specially planned flights of the Concorde 001. The obtained data have made possible the validation of some models predicting the condition of the stratosphere after a few decades of contamination by pollutants.

M.V.E.

A75-11730 More information and less data from space. H. M. Gurk (RCA, Astro-Electronics Div., Princeton, N.J.). *RCA Engineer*, vol. 20, Aug.-Sept. 1974, p. 38-41. 9 refs.

Higher resolution, more spectral bands, and greater frequency of coverage are desires of most users of remote sensor data. However, considering only current programs like the Earth Resources Technology Satellite, environmental satellites, and aircraft-based sensor systems, the volume of data offers a substantial processing problem to NASA. Potential data users are experiencing similar difficulties. Increased facilities and manpower may not be the best solution. Current analyses are revealing how much of the data is redundant. If the important information were extracted from the large volume of data, it would be possible simultaneously to reduce the volume and permit desired improvements in resolution or frequency of observation. Techniques such as data compression, statistical sampling, and mixed-highs multispectral sensing can reduce the data greatly. Data volume reduction, problems of utilization, and usefulness are discussed for each technique. (Author)

A75-11732 A microwave correlation spectrometer for air pollution monitoring. A. K. Ghosh and H. J. Moody (RCA, Ltd., Montreal, Canada). *RCA Engineer*, vol. 20, Aug.-Sept. 1974, p. 46-48. 11 refs.

The new instrument reported makes use of the adaptation of a correlation technique to microwave spectroscopy. Questions concerning the employment of microwave spectroscopy for pollutant analysis are considered. The principle of correlation spectroscopy is based on the fact that the cross-correlation between spectra of different gases is very small. In the new instrument the basic principles are somewhat modified to ensure better stability and sensitivity.

G.R.

A75-12031 * # The progeny of Skylab - The pioneering phase of our expansion into space. H. H. Schmitt (NASA, Office of Energy Programs, Washington, D.C.). American Institute of Aeronautics and American Geophysical Union, Conference on Scientific Experiments of Skylab, Huntsville, Ala., Oct. 30-Nov. 1, 1974, AIAA Paper 74-1216. 4 p.

The opportunities provided by such near-earth space resources as the continuous view of the earth, sun, and stars, the presence of ultrahigh vacuum, and the absence of gravitational stress are briefly reviewed, along with the significance of manned observation and experimentation in near-earth space. The stimulation triggered in young minds by a week or summer in space, as the Apollo and Skylab experiences show, simply defies the imagination. The plea is made not to miss these opportunities.

M.V.E.

A75-12035 * # Skylab altimeter applications and scientific results. J. T. McGoogan, C. D. Leitao (NASA, Wallops Flight Center, Wallops Island, Va.), W. T. Wells (Wolf Research and Development Corp., Riverdale, Md.), L. S. Miller, and G. S. Brown (Applied Science Associates, Inc., Apex, N.C.). American Institute of Aeronautics and Astronautics and American Geophysical Union, Conference on Scientific Experiments of Skylab, Huntsville, Ala., Oct. 30-Nov. 1, 1974, AIAA Paper 74-1221. 14 p. 7 refs.

Analysis of the Skylab S-193 Altimeter Experiment data has produced many significant results. The purpose of this paper is to present some of these results which illustrate the programmatic altimeter technology gained and show the scientific potential of altimetry for various applications, including correlation with existing topographic features, geoids and physical models. It will be shown that there exists a very strong correlation between subsurface topography in the broad ocean area and the shape of the ocean surface as determined from the altimeter. Modeled scattering waveforms will be correlated with Skylab Altimeter waveforms. The agreement of a variety of existing geoid models with the altimeter data will be shown. In addition the calibration, accuracy and stability of the S-193 altimeter instrumentation will be discussed. (Author)

A75-12036 * # Automatic land use classification using Skylab S-192 multispectral data. L. Kirvida and M. Cheung (Honeywell Systems and Research Center, Minneapolis, Minn.). American Institute of Aeronautics and Astronautics and American Geophysical Union, Conference on Scientific Experiments of Skylab, Huntsville, Ala., Oct. 30-Nov. 1, 1974, AIAA Paper 74-1224. 7 p. Contract No. NAS9-13386.

Investigation of the accuracy attainable in automatic land use classification using 13 bands of multispectral data from the Skylab S-192 scanner. Classification to levels containing seven urban classes, five agricultural, and three water classes is shown to be achievable. With 17 classes, a classification accuracy of 72% was obtained. A wide spectral range, including the thermal band, appears to be most useful for distinguishing urban classes. Agricultural and water classes can be separated using spectral bands covering the visible to far IR.

A75-12109 * Very high resolution radiometer. A. I. Aronson (RCA, Astro-Electronics Div., Princeton, N.J.). RCA Engineer, vol. 20, June-July 1974, p. 20-25. 6 refs. Contract No. NAS5-10306.

A primary sensor used in environmental and earth-resource observation, the Very High Resolution Radiometer (VHRR) was designed for use on the ITOS D series spacecraft. The VHRR provides a 0.47 mile resolution made possible with a mercury-cadmium-telluride detector cooled to approximately 105 K by a passive radiator cooler. The components of this system are described. The optical subsystem of the VHRR consists of a scanning mirror, a Dall-Kirkham telescope, a dichroic beam splitter, relay lenses, spectral filters, and an IR detector. Signal electronics amplify and condition the signals from the infrared and visible light detector. Sync generator electronics provides the necessary time signals. Scan-drive electronics is used for commutation of the motor winding, velocity, and phase control. A table lists the performance parameters of the VHRR.

A75-12317 # Detection of synchronous changes in fluxes of energetic electrons and gamma-quanta during a magnetic disturbance (Obnaruzhenie sinkhronnykh izmenenii potokov energichnykh elektronov i gamma-kvantov vo vremia magnitnogo vozmushcheniia). S. A. Voronov, A. M. Gal'per, V. G. Kirillov-Ugriumov, A. V. Kurochkin, N. G. Leikov, B. I. Luchkov, A. I. Fesenko, and Iu, T. Iurkin. (Vsesoiuznaia Konferentsiia po Fizike Kosmicheskikh Luchei, Kharkov, Ukrainian SSR, Sept. 25-28, 1973.; Akademiia Nauk SSSR, Izvestiia, Seriia Fizicheskaia, vol. 38, Sept. 1974, p. 1966-1969. 5 refs. In Russian.

A75-12874 Special color analysis techniques. P. G. Smith (USAF, Rome Air Development Center, Griffiss AFB, N.Y.), K. R. Piech, and J. E. Walker (Calspan Corp., Buffalo, N.Y.). Photogrammetric Engineering, vol. 40, Nov. 1974, p. 1315-1319, 1321, 1322. 11 refs.

Special color analysis techniques offer an approach to improve ithe U.S. Air Force's capability for extracting additional information

from reconnaissance sensor records through the use of the photometric properties of the sensor records. A program concerned with the development of such techniques has provided a console for the performance of photometric analysis. Nonmilitary applications of the new techniques are related to crop analysis, mapping of flooding conditions, monitoring of water quality, and analyses of ERTS and Skylab imagery.

G.R.

A75-12947 # Automated remote measurement system in agricultural meteorology (Avtomatizirovannaia sistema distantsionnykh izmerenii v agrometeorologii). V. A. Kovalenko, A. D. Kleshchenko, and O. V. Virchenko (Glavnoe Upravlenie Gidrometeorologicheskoi Sluzhby SSSR, Institut Eksperimental'noi Meteorologii, Obninsk, USSR). Meteorologiia i Gidrologiia, Sept. 1974, p. 93-97. 15 refs. In Russian.

An automated system for the acquisition and processing of aerophotometry data on young agricultural crops is examined. The system's principles of operation are outlined, and its block diagram and principal elements are discussed.

A75-12980 * # Satellite techniques for the study of secular variations in position. J. W. Siry (NASA, Washington, D.C.). In: Symposium on Earth's Gravitational Field and Secular Variations in Position, Sydney, Australia, November 26-30, 1973, Proceedings. Kensington, Australia, University of New South Wales, 1974, p. 360-379, 30 refs.

Some crustal motions which are precursory to earthquakes are discussed and a system for monitoring such motions is proposed. Studies have shown that the scale of a dilatant region is an indicator both of the magnitude of subsequent earthquake and of the time interval between the earthquake and the precursory phenomena. Monitoring may be accomplished by the use of corner reflectors at sites situated in an array over a given region, such as California, for example, the sites being tracked by means of a laser system operating in the Geopause satellite.

P.T.H.

A75-12985 * # Earth parameters from global satellite triangulation and trilateration. I. I. Mueller (Ohio State University, Columbus, Ohio). In: Symposium on Earth's Gravitational Field and Secular Variations in Position, Sydney, Australia, November 26-30, 1973, Proceedings. Kensington, Australia, University of New South Wales, 1974, p. 529-552; Discussion, p. 553. 14 refs. Grant No. NGL-36-008-093.

Results obtained from 159-station global satellite triangulation and trilateration (including Baker-Nunn, BC-4, PC-1000 camera observations, SECOR, C-Band radar and EDM distance measurements) indicate differences in the semidiameter and orientation of the earth compared to results obtained from dynamic satellite solutions. Geoidal undulations obtained can be made consistent with dynamically determined ones at the expense of slight changes in the currently accepted parameters defining the gravity field of the level ellipsoid. (Author)

A75-13148 Trace gas analysis of power plant plumes via aircraft measurement · O3, NOx, and SO2 chemistry. D. D. Davis, G. Smith, and G. Klauber (Maryland, University, College Park, Md.). Science, vol. 186, Nov. 22, 1974, p. 733-736. 7 refs. NSF Grant No. GI-36338X.

A75-13589 Apollo Telescope Mount experiments technology. W. C. Keathley (NASA, Marshall Space Flight Center, Huntsville, Ala.). In: Technology today for tomorrow; Proceedings of the Eleventh Space Congress, Cocoa Beach, Fla., April 17-19, 1974. Volume 2.1

Cape Canaveral, Fla., Canaveral Council of Technical Societies, 1974, p. 1-1 to 1-45. 10 refs.

The Skylab Apollo Telescope Mount (ATM) experiments, consisting of a white light coronagraph, four ultraviolet instruments,

two X-ray telescopes and two hydrogen-alpha telescopes, observed the sun daily for nine solar rotations. The ATM instruments were individually larger, more complex, and provided better spatial resolution than previous solar satellite instruments. An additional major advantage of the telescope complex was its ability to simultaneously collect multispectral data of specific solar phenomena. To maximize the scientific benefits of the orbiting observatory, a coordinated observing program involving worldwide ground-based observatories was conducted. A description of the Skylab, the ATM, and the ATM experiments is given, and the daily process of flight planning and execution is described. (Author)

A75-13765 Spectrophotometric observations of earth from 'Soyuz-13'. A. A. Buznikov, K. Ia. Kondrat'ev, V. V. Lebedev, and V. M. Orlov (Leningradskii Gosudarstvennyi Universitet, Leningrad, USSR). International Astronautical Federation, International Astronautical Congress, 25th, Amsterdam, Netherlands, Sept. 30-Oct. 5, 1974, Paper 74-156. 20 p. 15 refs.

The use of different geometry of measurements from a satellite and of different conditions of solar illumination of the atmosphere and the underlying surface permits the obtaining of the information both of the gaseous and aerosol composition and of the state of different natural formations on the earth's surface. Therefore it is necessary to point out that the results of measurements on Soyuz 13 of solar radiation reflected and scattered by the surface-atmosphere system open up the possibility of a further development of the methods for the solution of inverse problems of atmospheric space optics.

F.R.L.

A75-13868 Investigations of the ionosphere by space techniques. S. A. Bowhill (Illinois, University, Urbana, Ill.). Journal of Atmospheric and Terrestrial Physics, vol. 36, Dec. 1974, p. 2235-2243. 56 refs.

Key developments in ionosphere research from new types of measurement using space vehicles are outlined. The usefulness of radio-propagation techniques in space research is discussed. Ground-based techniques such as incoherent scatter may be used to duplicate measurements made from space vehicles. The remote sensing of atmospheric minor constituents by optical techniques is reviewed.

T.S

A75-13963 Skylab. II - The three month vigil. D. Baker. Spaceflight, vol. 16, Dec. 1974, p. 456-461.

The Skylab 3 flight is traced chronologically from Day 7 to Day 15. Continuing problems were created by the failure of Control Moment Gyro (CMG) No. 1 on Day 8 and abnormal operation of CMG-2. Extra-vehicular activities (EVA), Kohoutek photography, the Black Brant IV rocket launching, earth resources experiments, medical experiments, and solar flare observations are considered.

A.T.S

A75-14726 Electronics in meteorological instrumentation. C. V. Else, D. J. Painting, and K. J. T. Sands (Meteorological Office, Bracknell, Berks., England). *IEE Reviews*, vol. 121, Nov. 1974, p. 1201-1223, 57 refs.

The evolution of electronic circuit applications in meteorological instrumentation is surveyed by discussing methods and equipment used for satellite, atmospheric-vehicle, ground, and sea based measurements. Attention is given to the principles of radiosondes, high-altitude wind measurement, ozone studies, and determination of such parameters as cloud height, visibility, surface winds, and insolation.

A75-14792 Performance of a hologram matrix radar. K. lizuka, H. Ogura, J. L. Yen, V. K. Nguyen (Toronto, University, Toronto, Canada), and J. Weedmark (Bell Northern Research, Ltd., Ottawa, Canada). *IEEE, Proceedings*, vol. 62, Dec. 1974, p. 1718, 1719. Research supported by the Department of Energy, Mines, and Resources of Canada.

A novel radar system, which had been designed as a holographic ice surveying system for the mapping of ice profiles, was developed and successfully tested. The principles of operation of the new radar are based on the concept of the hologram matrix. The radar can be considered either as a new application of holography or as a synthetic phased array system. The results of an experimental investigation show that the novel radar can be used for mapping the profile of thin ice.

G.R.

A75-14803 * Estimation of sea surface temperature from remote sensing in the 11- to 13-micron window region. C. Prabhakara, V. G. Kunde (NASA, Goddard Space Flight Center, Greenbelt, Md.), and G. Dalu (CNR, Istituto di Fisica dell'Atmosfera, Rome, Italy). Journal of Geophysical Research, vol. 79, Nov. 20, 1974, p. 5039-5044. 26 refs.

The Nimbus 3 and 4 Iris spectral data in the 11- to 13-micron water vapor window region are analyzed to determine the sea surface temperature (SST). The high spectral resolution data of Iris are averaged over approximately 1-micron-wide intervals to simulate channels of a radiometer to measure the SST. In the present exploratory study, three such channels in the 775- to 960-per cm (12.9-10.5 micron) region are utilized to measure the SST over cloud-free oceans. However, two of these channels are sufficient in routine SST determination. The differential absorption properties of water vapor in the two channels make it possible to determine the water vapor absorption correction without detailed knowledge of the vertical profiles of temperature and water vapor. The feasibility of determining the SST is demonstrated globally with Nimbus 3 data, where cloud-free areas can be selected with the help of albedo data from the medium-resolution infrared radiometer experiment on board the same satellite. The SST derived from this technique agrees with the measurements made by ships to about 1 C. (Author)

A75-15434 Hologrammetry - Concepts and applications. E. M. Mikhail (Purdue University, West Lafayette, Ind.). *Photogrammetric Engineering*, vol. 40, Dec. 1974, p. 1407-1422, 12 refs. Army-sponsored research.

Research in holographic applications to mapping during the past several years has led to the introduction of the term hologrammetry. It is an interdisciplinary field combining holographic techniques for purposes of interpretation, mensuration, mapping, and display. A concise account of Fresnel and focused-image holograms, both transmission and reflection types, includes their characteristics and suitable applications. Mensuration and mapping techniques of holograms and results obtained follow. This includes targeting and digital mapping as well as graphical plotting of contours, cross-sections, etc. The use of direct holograms has been practically limited to small close-range objects. To extend the capability to topographic terrain applications from aerial photography, one must apply the concept of the holographic stereomodel. Both Fresnel and focused-image holographic stereomodels possess related advantages. (Author)

A75-15953 # Mathematical modeling for the detection of fish by an airborne laser. D. L. Murphree, C. D. Taylor (Mississippi State University, Mississippi State, Miss.), and R. W. McClendon. AIAA Journal, vol. 12, Dec. 1974, p. 1686-1692. 12 refs. Research supported by the National Marine Fisheries Service.

To assess the feasibility of an airborne remote sensing laser system for fish detection, a mathematical model was developed and solved which incorporated the physical interactions involved in the process of laser transmission through the air/sea interface and ocean environment. For a circular laser beam incident normal to the ocean surface, the power density was determined beneath the ocean surface as a function of surface roughness (wind velocity) and depth. Using the principle of electromagnetic reciprocity, the power reflected from submerged targets and detected at the airborne receiver was determined. The results from the developed mathematical model,

08 INSTRUMENTATION AND SENSORS

using input parameters of presently available equipment and estimates of fish school density and reflectivity, reveal that the power received at an airborne detector from fish reflected incident laser radiation and the signal-to-noise ratio (S/N) are of sufficient magnitude to locate fish schools with an airborne remote sensing laser system. (Author)

A75-16096 * Skylab radar altimeter - Short-wavelength perturbations detected in ocean surface profiles. C. D. Leitao and J. T. McGoogan (NASA, Wallops Flight Center, Wallops Island, Va.). Science, vol. 186, Dec. 27, 1974, p. 1208, 1209. 7 refs.

Short-wavelength anomalies in sea surface topography, caused by the gravitational effects of major ocean bottom topographic features, have been detected by the radar altimeter aboard Skylab. Some features, such as deep ocean trenches, seamounts, and escarpments, displace the ocean surface by as much as 15 meters over 100-kilometer wavelengths. This experiment demonstrates the potential of satellite altimetry for determining the ocean gooid and for mapping major features of the ocean bottom. (Author)

A75-16188 # Images from the NOAA-3 very high resolution radiometer over the North Sea and adjoining countries. P. K. Rao and E. P. McClain (NOAA, National Environmental Satellite Service, Washington, D.C.). Weather, vol. 29, Dec. 1974, p. 436-442.

A75-16191 A radio picture of the earth. W. J. Webster, Jr., T. T. Wilheit, T. C. Chang, P. Gloersen, and T. J. Schmugge (NASA, Goddard Space Flight Center, Atmospheric and Hydrospheric Applications Div., Greenbelt, Md.). Sky and Telescope, vol. 49, Jan. 1975, p. 14-16.

A technique called passive microwave remote sensing can be used to obtain a new view of the planet earth by means of radio telescopes carried aboard artificial satellites. An important relationship between the observed radio brightness temperature and the surface conditions provides the basis for the new technique. A radio image is presented of the entire earth on the basis of Nimbus microwave-image data taken January 12-16, 1973.

G.R.

A75-16485 * Remote atmospheric sensing with an airborne laser absorption spectrometer. R. T. Menzies and M. T. Chahine (California Institute of Technology, Jet Propulsion Laboratory, Pasadena, Calif.). Applied Optics, vol. 13, Dec. 1974, p. 2840-2849. 25 refs. Contract No. NAS7-100.

A laser absorption spectrometer, using an IR laser transmitter and a heterodyne radiometer, can be used from an aircraft or spacecraft to measure altitude profiles of air pollutants and other atmospheric constituents. The technique involves measurement of differential absorption at several wavelengths, using the diffusely reflecting earth's surface to provide a return signal. The pressure broadening of absorption lines allows one to discriminate between high and low altitude absorbers. Application of the technique to measurements of ozone, nitric oxide, and water vapor are presented. CO2 and CO lasers are considered as transmitters. The discussion includes altitude resolution limitations, atmospheric temperature dependence, and frequency stability requirements of the instrument. (Author)

A75-17101 * International Symposium on Remote Sensing of Environment, 9th, University of Michigan, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volumes 1, 2 & 3. Symposium sponsored by the Environmental Research Institute of Michigan, University of Michigan, NASA, et al. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974. Vol. 1, 808 p.; vol. 2, 797 p.; vol. 3, 684 p. Price of three volumes, \$35.

The present work gathers together numerous papers describing the use of remote sensing technology for mapping, monitoring, and management of earth resources and man's environment. Studies using various types of sensing equipment are described, including multispectral scanners, radar imagery, spectrometers, lidar, and aerial photography, and both manual and computer-aided data processing techniques are described. Some of the topics covered include: estimation of population density in Tokyo districts from ERTS-1 data, a clustering algorithm for unsupervised crop classification, passive microwave sensing of moist soils, interactive computer processing for land use planning, the use of remote sensing to delineate floodplains, moisture detection from Skylab, scanning thermal plumes, electrically scanning microwave radiometers, oil slick detection by X-band synthetic aperture radar, and the use of space photos for search of oil and as fields.

Individual items are announced in this issue.

P.T.H.

A75-17111 * # Transference of ERTS-1 spectral signatures in time and space, B. F. Merembeck, F. Y. Borden, and D. N. Applegate (Pennsylvania State University, University Park, Pa.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 153-161. Research supported by the Hatch Funds; Contract No. NAS5-23133.

Some earth resources maps obtained from ERTS-1 multispectral scanning data in 1972-73 were found to contain modulo six banding. A method was found to recalibrate nonconformable data from the sensors in order to eliminate banding in the map output. This investigation led to developing a recalibration method for transferring data signatures from one scene to another scene separated from the first either temporally or spatially. The method was generally successful, although difficulties were encountered in transferring signatures from one season to another for the same location.

A.T.S.

A75-17119 * # Signatures of various earth surfaces measured by the Nimbus-5 microwave spectrometer. K. F. Kunzi, R. L. Pettyjohn, D. H. Staelin (MIT, Cambridge, Mass.), and J. W. Waters (California Institute of Technology, Jet Propulsion Laboratory, Pasadena, Calif.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 299-305. 6 refs. Contract No. NAS7-100.

The Nimbus-5 Meteorological Satellite is equipped with a 5-channel microwave spectrometer. The two lowest channels (22.2 and 31.4 GHz) provide information on surface brightness temperature. Distinctive microwave signatures can be observed for snow, land ice, and sea ice in both polar regions. Some information on subsurface temperatures can be gained for desert areas. (Author)

A75-17124 # Passive microwave sensing of moist soils. A. E. Basharinov, L. F. Borodin, and A. M. Shutko (Akademiia Nauk SSSR, Moscow, USSR). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 363-367. 8 refs.

Passive microwave observations of moist soils were performed at wavelengths of 0.8 and 3.4 cm from an aircraft. Data obtained are compared with the results of direct measurements of soil moisture content. The dependence of brightness temperature on the value of moisture content is observed, as well as the influence of vegetation cover.

(Author)

A75-17141 • # Application of remote sensing to the location of hydrologically active /source/ areas. A. M. Ishaq and D. D. Huff (Wisconsin, University, Madison, Wis.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April

15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 653-666, 21 refs. Grant No. NGL-50-002-127.

Recent hydrologic research has provided theoretical support for a hypothesis generally known as the variable source area concept. However, experimental study of variable source areas has been hampered by lack of suitable methods for identifying and mapping them on a large scale basis. Field studies combined with color infrared photography and direct measurement of soil moisture and other hydrologic variables have shown that it is definitely possible to differentiate high and low soil moisture zones in a watershed. Furthermore, mapping of high moisture zones for the 1839 hectare Lowery Creek, Wisconsin drainage basin has revealed a pattern that is fully consistent with hypotheses concerning source areas. Examination of transmittance curves for high moisture zones indicates that source areas may be delineated on the basis of the ratio of their transmittance at 550 nm to that at 675 nm.

(Author)

A75-17142 * # The use of remote sensing and natural indicators to delineate floodplains - Preliminary findings. S. C. Sollers, G. W. Petersen, D. L. Henninger, and F. Y. Borden (Pennsylvania State University, University Park, Pa.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich, April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 667-681. 9 refs. NASA-supported research.

Feasibility studies to delineate floodplains using aerial photography have been conducted by the Army Corps of Engineers. The objective of this project is to establish floodplain boundaries using natural indicators such as vegetation types, soil types, moisture differences, and geologic variations. Using aircraft multispectral data, an automated mapping routine should enable the Corps of Engineers to reduce costly field surveys and improve the accuracy of floodplain delineation in pristine areas. The purpose of this paper is to document the methodology utilized in this investigation while using one of a series of available data sets and test sites and to report preliminary findings. (Author)

A75-17147 # On the feasibility of remote monitoring of soil moisture with microwave sensors. R. W. Newton, S. L. Lee, J. W. Rouse, Jr. (Texas A & M University, College Station, Tex.), and J. F. Paris (Lockheed Electronics Co., Inc., Houston, Tex.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 725-738, 16 refs.

Recent airborne measurements of radiometric temperature at monitored sites located near Chickasha, Oklahoma, and Weslaco, Texas show that vegetation cover has the effect of masking the soil moisture dependence of the microwave data. Peake's model is inappropriate for describing emission in the presence of a vegetative cover; therefore, a new model was developed which computes emissivity directly from the material properties of the subsurface media and the transmission coefficients of the surface. This work was used to develop a model for the apparent microwave temperature and radar backscatter coefficient of vegetated terrain to illustrate the effects of vegetation on the sensitivity of these parameters to variations of soil moisture. A ground measurement program was established in order to obtain data to compare the predictions of the models. (Author)

A75-17148 • # Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models. L. R. Tinney, J. E. Estes, K. H. Tharman, and R. R. Thaman (California, University, Santa Barbara, Calif.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 739-757. NASA-supported research.

A75-17192 # The interpretation and use of false-colour infra-red and true colour photography of part of Argentina obtained by Skylark earth resources rockets. D. S. H. Drennan, C. J. Bray, J. R. Hardy, C. O. Justice, R. A. G. Savigear, J. R. G. Townshend (Reading University, Reading, England), I. R. Galloway, and E. S. Owen-Jones (Bedford College, London, England). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1475-1496. 16 refs. Ministry of Defense (Procurement Executive) Grants No. AT/2035/015/SP; No. AT/2035/025/ASA.

A75-17194 # A preliminary feasibility study on a synthetic aperture radar /SARSAT/ for earth resources surveys. C. Skenderoff, J. P. Guignard (Thomson - CSF, Meudon, Hauts-de-Seine, France), J. R. Coldrick, H. Mooney (British Aircraft Corp., Bristol, England), and J. Noel (Société d'Etudes Techniques et d'Entreprises Générales, le Plessis-Robinson, Hauts-de-Seine, France). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1517-1540. Research supported by the European Space Research Organization.

A75-17195 * # Investigation of microwave hologram techniques for application to earth resources. R. W. Larson, R. W. Bayma, M. B. Evans, J. S. Zelenka, H. W. Doss (Michigan, Environmental Research Institute, Ann Arbor, Mich.), and J. E. Ferris (Michigan, Environmental Research Institute; Michigan, University, Ann Arbor, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1541-1569. 20 refs. Contract No. NAS1-10810.

An investigation of microwave hologram techniques for application to earth resources was conducted during the period from June 1971 to November 1972. The objective of this investigation has been to verify the feasibility of an orbital microwave holographic radar experiment. The primary advantage of microwave hologram radar (MHR) over the side-looking airborne radar (SLAR) is that of aspect or viewing angle; the MHR has a viewing angle identical with that of photography and IR systems. The combination of these systems can thus extend the multispectral analysis concept to span optical through microwave wavelengths. Another advantage is the capacity of the MHR system to generate range contours by operating in a two-frequency mode. It should be clear that along-track resolution of an MHR can be comparable with SLAR systems, but cross-track resolution will be approximately an order of magnitude coarser than the range resolution achievable with an arbitrary SLAR system. An advantage of the MHR over the SLAR is that less average transmitter power is required. This reduction in power results from the much larger receiving apertures associated with MHR systems. (Author)

A75-17197 # Emissivities and forward scattering of natural and man-made material at three millimeter wavelength. E. Schanda and R. Hofer (Bern, Universität, Berne, Switzerland). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1585-1592.

For the interpretation of the thermal images produced by a high-resolution 3-mm scanning radiometer, an investigation of the emissivities of natural and man-made materials has been performed. In order to exclude the atmospheric contributions to the measured radiation and to study the angular dispersion of a reflected wave due to surface structuring, the forward scatter properties were mainly investigated. In general, it may be stated that the emissivities of all

materials are closer to unity or equal to those at 3-cm wavelength. The emissivity of water varies from roughly 0.7 to 0.6 for a temperature change from a few degrees centigrade up to about 40 C. An oil film of 0.15 mm increases the emissivity by 0.05. (Author)

A75-17206 # Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills. J. F. Fantasia and H. C. Ingrao (U.S. Department of Transportation, Transportation Systems Center, Cambridge, Mass.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1711-1745. 12 refs. Research supported by the U.S. Coast Guard and U.S. Navy.

Results of a study and measurements program to determine the feasibility of using laser-excited oil fluorescence as a means of detecting, classifying, and quantifying oil spills in the marine environment. The study consisted of an analysis of the fluorescence properties of oils and oil slicks on the sea surface, and a theoretical analysis of the remote fluorometry of oil spills. Laboratory measurements were made of 29 crude and refined oils commonly transported in the marine environment. These measurements included API gravity, fluorescence and reflectance spectra, fluorescence coefficient, and fluorescence lifetimes. Similar measurements were made with a laboratory model of an N2 laser oil spill remote sensor. Results of these measurements clearly showed that, under certain conditions, oil spill detection and classification can be made in the ocean environment. Furthermore, remote sensing of oil spills using laser-excited oil fluorescence was found to be technically feasible.

(Author)

A75-17208 # Multi-frequency radiometric measurements of foam and a mono-molecular slick. B. Au, J. Kenney, L. U. Martin (U.S. Navy, Naval Research Laboratory, Washington, D.C.), and D. Ross (NOAA, Miami, Fla.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1763-1773.

Microwave radiometric measurements have been made of both a surf-zone and an ocean region where small-scale roughness was suppressed by an artificial monomolecular slick. The foam measurements show near identical foam temperatures at 8.35 and 14.5 GHz, but large variations at 1.4 GHz. The resultant maximum foam emissivities at nadir range from 0.57 at 1.4 GHz to 0.84 at 14.5 GHz. The presence of the monomolecular slick on the ocean surface had the same effect as a decrease in surface roughness. For horizontal polarization, the emission decreased below that of the surrounding ocean for all viewing angles. At vertical polarization, the emission decreased below and increased above a viewing angle of approximately 60 deg. (Author)

A75-17209 # Oil slick detection by X-band synthetic aperture radar. J. R. Kotlarski and H. R. Anderson (Hughes Aircraft Co., Culver City, Calif.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1775-1790.

A75-17210 # The application of spectrometric and polarization techniques for remote sensing of oil on sea water. K. Ia. Kondrat'ev, A. A. Buznikov, D. V. Pozdiakov, G. A. Ivanian, G. A. Lakhtanov, V. M. Orlov, and S. M. Mikhailova (Leningradskii Gosudarstvennyi Universitet, Leningrad, USSR). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1793-1802. 18 refs.

A75-17212 # An orbiting visible/infrared spectrometer for terrestrial, atmospheric and oceanographic applications. G. R. Pruitt (Block Engineering, Inc., Cambridge, Mass.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1847-1855.

This paper describes the S191 spectrometer developed for use on NASA's Skylab. Design and key features of the instrument are discussed. Performance characteristics, as they relate to earth's resources measurements requirements and capabilities, are presented. Orbital data from the first manned Skylab mission are discussed. Conclusions and recommendations for future usage of this instrument are made. (Author)

A75-17213 # A motion-compensated spatial scanner. K. S. Gordon and J. R. Miller (York University, Toronto, Canada). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1857-1863. 5 refs. Research supported by the National Research Council of Canada.

A high resolution multispectral scanner has been built and tested. An interference filter photometer is employed and the way in which this instrument complicates the design of a line scanner is discussed. Preliminary flight test results are presented and plans for future high resolution water studies of phytoplankton pigment concentrations are described.

(Author)

A75-17214 # The remote Raman spectrometer is a viable instrument for remote sensing of the environment. S. M. Klainer, W. Arden, and T. Hirschfeld (Block Engineering, Inc., Cambridge, Mass.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Yolume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1865-1876.

The major components of the remote Raman spectrometer system, using a pulsed laser as the monochromatic source, are illustrated and discussed. The laser beam illuminates the sample, scattering being caused by all of the constituents. This is observed through a telescope that is through-put matched to a spectral sorter. The Raman lines are sorted into distinct frequencies. The output of the sorter is detected by photomultiplier tubes, is processed, and is displayed on a chart recorder. Use of a pulsed laser and time gating permits measurement of the distance between the scattering molecules and the light source and collector.

V.P.

A75-17217 * # Multi-aspect techniques in remote sensing. W. A. Malila (Michigan, Environmental Research Institute, Ann Arbor, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1905-1918. 12 refs. Contract No. NAS9-9784.

Multi-aspect techniques, which use multispectral scanner information collected from two or more different view angles for each spatial element, are discussed. Empirical recognition results obtained with various combinations of information channels are presented. Theoretical calculations of multi-aspect reflectances of corn fields also are analyzed and compared with empirical data. Also presented is a brief comparison between recognition processing and scene attribute estimation techniques for extracting information from multispectral scanner data. One method for improving information extraction by employing a post-recognition attribute estimation procedure is suggested. (Author)

A75-17221 # Prediction of the Fraunhofer line detectivity of luminescent materials. R. D. Watson, T. D. Hessin, R. C. Bigelow (U.S. Geological Survey, Denver, Colo.), and W. R. Hemphill (U.S. Geological Survey, Reston, 'Va.). In: International Symposium on

Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1959-1980. 11 refs.

A program is described in which the luminescence of materials at six Fraunhofer lines was quantified. Measurements were made with a laboratory fluorescent spectrometer on selected materials, including crude and refined oils, crude-oil films, oil shales, phosphate, effluents, and in-vivo chlorophyll of geochemically stressed and nonstressed vegetation. Results indicate that the luminescence of the petroleum and phosphate effluent samples studied is equivalent to, or exceeds by several orders of magnitude the luminescence of Rhodamine WT dye at the minimum concentration detectable with the Fraunhofer line discriminator (FLD). Geochemically stressed and nonstressed vegetation luminesce at the 656.3-nm Fraunhofer line at intensities that vary both diurnally and seasonally from levels just within the sensitivity limits of an FLD to one order of magnitude above this level. An improved FLD has been built and preliminary tests were performed on some of the materials noted above. This instrument was found to respond to Rhodamine WT dye concentrations as small as microgram/l. (Author)

A75-17223 # Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data. R. Cassinis, C. M. Marino (Milano, Università, Milan, Italy), and A. M. Tonelli (CNR, Laboratorio per la Geofisica della Litosfera, Milan, Italy). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1989-2004. 6 refs.

A75-17230 * # The remote-sensing program of the Geological Survey of Alabama. J. A. Drahovzal, J. L. G. Emplaincourt, and C. C. Wielchowsky (Alabama, University, University, Ala.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 2059-2072. 19 refs. U.S. Geological Survey Contract No. 14-08-001-13377; Contract No. NASS-21876.

Some areas of remote-sensing research in Alabama are reviewed, showing how remote sensing technology can provide data that cannot be obtained by other methods. Applications of this technology to structural and stratigraphic research and to the search for water, mineral, and energy sources are discussed, and some results of these applications are illustrated.

V.P.

A75-17231 * # Resource analysis applications in Michigan. S. W. Schar, W. R. Enslin (Michigan State University, East Lansing, Mich.), I. J. Sattinger (Michigan, Environmental Research Institute, Ann Arbor, Mich.), J. G. Robinson, K. R. Hosford (Michigan Department of Natural Resources, Lansing, Mich.), R. S. Fellows (Soil Conservation Service, Lansing, Mich.), and J. H. Raad (U.S. Department of State Highways and Transportation, Bureau of Transportation Planning, Lansing, Mich.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 2073-2081. Research supported by the Environmental Research Institute of Michigan and NASA.

During the past two years, available NASA imagery has been applied to a broad spectrum of problems of concern to Michigan-based agencies. These demonstrations include the testing of remote sensing for the purposes of (1) highway corridor planning and impact assessments, (2) game management-area information bases, (3) multi-agency river basin planning, (4) timber resource management information systems, (5) agricultural land reservation policies, and (6) shoreline flooding damage assessment. In addition, cost accounting procedures have been developed for evaluating the relative costs of utilizing remote sensing in land cover and land use analysis data collection procedures. (Author)

A75-17232 # Applications of remote sensing by the state of Nebraska. M. Carlson, R. Petersen, R. Hoffman, J. Drew, D. Edwards, G. Hergenrader, N. Rosenberg, L. Sheffield (Nebraska, University, Lincoln, Neb.), M. Ball (Nebraska State Department of Water Resources, Lincoln, Neb.), and J. Barr (Nebraska State Office of Planning and Programming, Lincoln, Neb.), In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 2083-2086.

Some past, present, and future remote sensing projects undertaken by the State of Nebraska are briefly described. A photomosaic and Level I land use map of Nebraska, both derived from ERTS-1 imagery, have been published at a scale of 1:1,000,000. In two counties, over 7000 fields were outlined from ERTS-1 imagery and classified as alfalfa, wheat, wetland pasture, range and pasture, nonirrigated row crops, and irrigated corn. An evapotranspiration project is determining the feasibility of using thermal imagery to obtain temperatures for use in evapotranspiration models. The possibility of measuring selected aspects of water quality in Nebraska reservoirs by remote sensing is also being investigated. The concentrations of chlorophyl and the degree of inorganic turbidity will be quantified by remote sensing.

A75-18071 The Skylab S-190B Earth Terrain Camera, J. D. McLaurin (U.S. Geological Survey, National Center, Reston, Va.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings. Falls Church, Va., American Society of Photogrammetry, 1974, p. 268-278.

The Skylab missions, completed in February 1974, included a wide variety of experiments in the fields of engineering, medicine, astronomy, and earth resources. The sensor of particular interest to the mapping and earth resources community was the S-190B Earth Terrain Camera (ETC). The ETC is a 460-mm focal-length frame camera with a 115- by 115-mm format. It includes an image motion compensation system of an f/4 color-corrected lens. The predicted low contrast (1.6:1) ground resolution for color photographs is 14 m/1p or 70 1p/mm; the expected resolution for black-and-white photographs is 10 m/1p or 100 1p/mm. Considerable cloud-free imagery using color, color-infrared, and black-and-white films was obtained during the three Skylab missions. A map showing the ETC coverage is included in the paper. Preliminary analyses indicate that the photographs have the necessary resolution and geometric fidelity for preparing photomaps at 1:100,000 scale and possibly larger scales. Other investigators are evaluating the photographs for geologic, mineral resources, and land-use mapping, and for archeological, water resources, and recreational studies. (Author)

A75-18072 Early results from the experimental Automated Image Data Extraction System. L. P. Murphy (U.S. Army, Geographic Sciences Laboratory, Fort Belvoir, Va.). In: American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings. Falls Church, Va., American Society of Photogrammetry, 1974, p. 300-321.

The USAETL experimental Automated Image Data Extraction System (AIDES-I) was fabricated to enable experimentation leading to the development of a man/machine interactive, multiple-image processing system to facilitate the extraction of terrain information (e.g. vegetation, soils, etc.) from a variety of imagery such as conventional aerial photographs, multiband images, and radar images. Examples of early data extraction experiments are shown and described which used multiband input images to spectrally separate features at analog processing rates. (Author)

A75-18693 Earth sciences in the age of the satellite. J. Pouquet (Aix-Marseille, Université, Aix-en-Provence, Bouches-du-Rhône, France). (Translation of Les sciences de la terre à l'heure des satellites /Télédétection/, Paris, Presses Universitaires de France, 1971.) Dordrecht, D. Reidel Publishing Co., 1974. 178 p. 146 refs. \$23.40.

Following a brief review of the beginnings of remote sensing, the basic principles are outlined. These make use of various bands of the spectrum, including the visible and near infrared range, the invisible range, and emitted and reflected radiations. It is considered that remote sensing techniques will occupy first place in earth science studies. Types of satellites are described. The problem of the real hour is discussed, followed by description of satellite instrumentation. Satellites intended for earth science studies are NERO, SCMR and ESMR, RBV, and MSS (ERTS A and B). Treatment of information received from satellites is studied, and comparative evaluation is made of aircraft vs satellites. The use of manned satellites, and general principles of interpretation are considered. Some geopedological and nongeopedological examples are discussed, as well as prospects for the future.

A75-19265 Conference on Environmental Sensors and Applications, London, England, November 18, 19, 1974, Proceedings. Conference sponsored by the Institution of Electronic and Radio Engineers. London, Institution of Electronic and Radio Engineers (IERE Conference Proceedings, No. 29), 1974. 239 p. \$20.75.

Recent advances in air pollution monitoring equipment and techniques, meteorological sensors, and remote sensing technology are reported. Some of the topics covered include: the point visibility meter - a forward scatter instrument for the measurement of aerosol extinction coefficient, a continuously recording aerosol photometer for monitoring respirable airborne dust, instrumentation for boundary layer research in meteorology, a new generation of sensors for automatic weather stations, measurement of rainfall and microwave attenuation by radar, the role of lidar in environmental sensing, and a submersible water quality station.

P.T.H.

A75-19266 The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient. J. V. Winstanley (Plessey Radar, Ltd., Cowes, Isle of Wight, England). In: Conference on Environmental Sensors and Applications, London, England, November 18, 19, 1974, Proceedings.

London, Institution of Electronic and Radio Engineers, 19/4, p.

A75-19274 The role of lidar in environmental sensing. J. V. Winstanley, M. J. Adams (Plessey Radar, Ltd., Cowes, Isle of Wight, England), and A. Whiteley (Plessey Radar, Ltd., South Leigh, Hants., England). In: Conference on Environmental Sensors and Applications, London, England, November 18, 19, 1974, Proceedings. . London, Institution of Electronic and

Radio Engineers, 1974, p. 163-173 (3 ff.). 29 refs.

19-31, 14 refs.

The basic lidar concept is explained with reference to a general form of the lidar equation. Six different extensions of the lidar technique are discussed, and their relative merits are reviewed. Two practical lidar systems are described in some detail - a laser ceilometer and a meteorological lidar for measurement of slant visual range at airfields under conditions of poor visibility. Results obtained with these instruments are presented, and their respective degrees of accuracy are determined. (Author)

A75-19547 Measurements of the extraterrestrial solar radiant flux from 2981 to 4000 A and its transmission through the earth's atmosphere as it is affected by dust and ozone. J. J. DeLuisi (National Center for Atmospheric Research, Boulder, Colo.). Journal of Geophysical Research, vol. 80, Jan. 20, 1975, p. 345-354. 25 refs.

N75-10532* National Oceanic and Atmospheric Administration, Rockville, Md.

SKYLAB A PROPOSAL AEROTRIANGULATION WITH VERY SMALL SCALE PHOTOGRAPHY Quarterly Report, 15 Jul. 15 Oct. 1974

Morton Keller, Principal Investigator 15 Oct. 1974 1 p EREP

(NASA Order T-4110-B)

(E74-10804: NASA-CR-140498) Avail: NTIS HC \$3.25 CSCL 08B

N75-10564*# Environmental Research Inst. of Michigan, Ann Arbor. Infrared and Optics Div.

THE APPLICATION OF REMOTE SENSING TECHNIQUES: TECHNICAL AND METHODOLOGICAL ISSUES

Fabian C. Polcyn and Thomas W. Wagner Oct. 1974 72 p refs Supported in part by NASA

(Grant NSF GI-34809X1)

(NASA-CR-140674; ERIM-193500-5-X) Avail: NTIS HC \$4.25 CSCL 14B

Capabilities and limitations of modern imaging electromagnetic sensor systems are outlined, and the products of such systems are compared with those of the traditional aerial photographic system. Focus is given to the interface between the rapidly developing remote sensing technology and the information needs of operational agencies, and communication gaps are shown to retard early adoption of the technology by these agencies. An assessment is made of the current status of imaging remote sensors and their potential for the future. Public sources of remote sensor data and several cost comparisons are included.

N75-10573# Physics Lab. RVO-TNO, The Hague (Netherlands). PATTERN RECOGNITION RELEVANT TO RECONNAIS-SANCE AND REMOTE SENSING

W. J. P. A. Verbeek Aug. 1974 48 p refs (PhL-1974-23; TDCK-65200) Avail: NTIS

An introduction to the nature of problem statements and solution techniques, especially concerning problems in automated aerial reconnaissance and remote sensing is presented. Stimulation of the awareness that classification and decision making involve more than intuition and that the literature offers an abundance of ready solutions in this field is proposed.

N75-10964*# Scientific Translation Service, Santa Barbara, Calif. THE EARTH IN THE LENS OF SALYUT-3

V. Ivanov Washington NASA Nov. 1974 13 p Transl. into ENGLISH from Aviat. Kosmonavt. (Moscow), no. 9, Sep. 1974 p 38-40

(Contract NASw-2483)

(NASA-TT-F-16028) Avail: NTIS HC \$3.25 CSCL 22C

Experiments performed by the cosmonauts on Salyut-3 and Soyuz-14 are described. The geological-morphological objects on the earth, as well as atmospheric formations and phenomena, were studied. It is concluded that this work represents one of the first steps in the conquest of space.

Author

N75-11310*# West Virginia Univ., Morgantown. Dept. of Electrical Engineering.

APPLICATION AND SENSITIVITY INVESTIGATION OF FOURIER TRANSFORMS FOR MICROWAVE RADIOMETRIC INVERSIONS

John J. Holmes and Constantine A. Balanis 15 Nov. 1974 99 p refs

(Grant NGR-49-001-056)

(NASA-CR-140754) Avail: NTIS HC \$4.75 CSCL 14B

Existing microwave radiometer technology now provides a suitable method for remote determination of the ocean surface's absolute brightness temperature. To extract the brightness temperature of the water from the antenna temperature equation, an unstable Fredholm integral equation of the first kind was solved. Fast Fourier Transform techniques were used to invert the integral after it is placed into a cross-correlation form. Application and verification of the methods to a two-dimensional modeling of a laboratory wave tank system were included. The instability of the Fredholm equation was then demonstrated and a restoration procedure was included which smooths the resulting oscillations. With the recent availability and advances of Fast Fourier Transform techniques, the method presented becomes very attractive in the evaluation of large quantities of data. Actual radiometric measurements of sea water are inverted using the restoration method, incorporating the advantages of the Fast Fourier Transform algorithm for computations.

N75-11311# California Univ., Livermore. Lawrence Livermore Lab.

FEASIBILITY STUDY OF A GUNN-DIODE MICROWAVE CAVITY SPECTROMETER AS A FORMALDEHYDE MONITOR Final Report

L. W. Hrubesh Feb. 1974 100 p refs (Contract W-7405-eng-48)

(UCID-16488) Avail: NTIS HC \$4.75

A prototype of a fixed frequency microwave spectrometer to detect a pure rorational absorption transition of the H2(1) C12016 molecule is described. The instrument was laboratory tested for use as a continuous monitor of formaldehyde in air. Design considerations, principles of operation, and calibration procedures are discussed.

F.O.S.

N75-11432* Scientific Translation Service, Santa Barbara, Calif. GEOGRAPHIC CORRELATION OF TELEVISION PICTURES OBTAINED FROM WEATHER SATELLITES

A. V. Bushuyev In its Space Iconics (NASA-TT-F-798) Sep. 1974 p 143-145 refs Transl into ENGLISH of the publ. "Kosmicheskaya Ikonika" Moscow, Nauka Press, 1973 p 123-124 CSCL 08B

The geographical control of satellite pictures, using the terminology of aerial photography, can be treated as the problem of analysis of a single picture with the objective of obtaining a ground contour map. Studies have shown the possibility and capability of the method of composing photographic maps from transformed television pictures. Optico-mechanical transformation solves the problem of geographical correlation for operational purposes. However, this technique does not compensate for electronic distortion, and accounts for earth sphericity only approximately. However, for certain purposes (studying ice drift), the maximum possible accuracy is required. Analytic geographical correlation methods using digital computers should be considered promising.

N75-11433* Scientific Translation Service, Santa Barbara, Calif. GEOGRAPHICAL CORRELATION OF TV AND IR IMAGES OBTAINED FROM WEATHER SATELLITES

Ya. L. Ziman, B. N. Nepoklonov, and B. N. Rodionov In its Space Iconics (NASA-TT-F-798) Sep. 1974 p 146-155 refs Transl. into ENGLISH of the publ. "Kosmicheskaya Ikonika" Moscow, Nauka Press, 1973 p 126-133

CSCL 08B

The requirements on the accuracy of geographical correlation of TV and IR cloud cover images are determined by the objective of the analysis of the weather information contained in these images. In the operational analysis case, the correlation accuracy need not be high. Errors of several tens of kilometers in determining the location of the cloud formation contours are considered acceptable in this case. Such correlation must be

provided in real image reception time. Scientific studies require accurate correlation of the cloud formation contours. The errors in determining their position should be commensurate with the imaging system resolution. The geometric aspects of developing methods and equipment for geographical correlation of television and infrared images of cloud cover taken from Meteor satellites are discussed.

N75-11434* Scientific Translation Service, Santa Barbara, Calif. SYSTEM FOR ELECTRONIC TRANSFORMATION AND GEOGRAPHIC CORRELATION OF SATELLITE TELEVISION INFORMATION

V. P. Dubenskiy, B. L. Nemkovskiy, and B. N. Rodionov In its Space Iconics (NASA-TT-F-798) Sep. 1974 p 155-164 refs Transl. into ENGLISH of the publ. "Kosmicheskaya Ikonika" Moscow, Nauka Press, 1973 p 133-140

CSCL 14E

An electronic transformation and correlation system has been developed for the Meteor space weather system which provides transformation and scaling of the original picture, accounts for satellite flight altitude and inclinations of the optical axes of the transmitting devices, and simultaneously superposes the geographical coordinate grid on the transformed picture.

Author

N76-11435* Scientific Translation Service, Santa Barbara, Calif. GEOGRAPHICAL CORRELATION OF EARTH CLOUD COVER PICTURES OBTAINED FROM UNORIENTED SATELLITES

V. I. Solovyev In its Space Iconics (NASA-TT-F-798) Sep. 1974 p 165-173 refs Transl into ENGLISH of the publ. "Kosmicheskaya Ikonika" Moscow, Nauka Press, 1973 p 141-146 (CSCL 048)

Mathematical models have been developed for the determination of the geographic coordinates of cloud photographs taken from earth satellites. The proposed methods were realized on a Minsk-22 computer, and were used to process the earth cloud pictures obtained from the communications satellite Molyniya 1 and the automatic stations Zond 5 and 7. Experience using these methods showed that the accuracy of orientation angular element determination is on the order of 1 deg.

Author

N75-11447*# Rice Univ., Houston, Tex. Inst. for Computer Services and Applications.

AN ANALYSIS OF APPLICATIONS DEVELOPMENT SYSTEMS FOR REMOTELY SENSED, MULTISPECTRAL DATA FOR THE EARTH OBSERVATIONS DIVISION OF THE NASA LYNDON B. JOHNSON SPACE CENTER

D. L. VanRooy, R. M. Smith, and M. S. Lynn Feb. 1974 127 p

(Contract NAS9-12776)

(NASA-CR-140345; Rept-275-025-018) Avail: NTIS HC \$5.75 CSCL 05B

An application development system (ADS) is examined for remotely sensed, multispectral data at the Earth Observations Division (EOD) at Johnson Space Center. Design goals are detailed, along with design objectives that an ideal system should contain. The design objectives were arranged according to the priorities of EOD's program objectives. Four systems available to EOD were then measured against the ideal ADS as defined by the design objectives and their associated priorities. This was accomplished by rating each of the systems on each of the design objectives. Utilizing the established priorities, it was determined how each system stood up as an ADS. Recommendations were made as to possible courses of action for EOD to pursue to obtain a more efficient ADS.

N75-11450# Army Foreign Science and Technology Center, Charlottesville, Va.

USE OF A PANORAMIC AERIAL CAMERA IN AEROSUR-VEYING

M. D. Konshin 25 Apr. 1974 9 p Transl. into ENGLISH from Geod. Kartografiya (USSR), no. 1, 1973 p 33-38 (AD-784162; FSTC-HT-23-0160-74) Avail: NTIS CSCL 08/2

The use of a panoramic aerial camera when carrying out aerial surveys is discussed. The photographic quality of the image is better due to the use of an exposure time of 1/3000 sec. The amount of field interpretation will be reduced and labor productivity will be increased by using the method suggested.

N75-11454# Texas Instruments, Inc., Dallas. Equipment Group.

OIL-ON-WATER SENSOR Final Engineering Report Kent McCormack, Gerald Fournier, and William Knight Feb.

1974 206 p refs (Contract DOT-CG-31888-A)

(AD-784104; TI-UI-998200-01-F; USCG-D-87-74) Avail:

NTIS CSCL 17/5

The Oil-on-Water Sensor system detects the presence of oil on water by emitting a wideband IR pulse and receiving reflections at two narrow spectral bands. Theory of operation is based on differences in the reflectance of water and oil at these narrow spectral bands. Ratio detection minimizes effects of surface wave action. This final report explains the principles of sensing, describes and explains the prototype equipment, and describes construction details of the Oil-on-Water Sensor. Finally, the report outlines the testing program that was used and provides data from these tests. Author (GRA)

N75-11562* Oxford Univ. (England). Clarendon Lab. INFRARED REMOTE SOUNDING

C. D. Rodgers In Calif. Univ. Proc. of the UCLA Intern. Conf. on Radiation and Remote Probing of the Atmosphere 1974 p 471-506 refs CSCL 04A

Atmospheric radiation was measured as a function of wavelength, position, angle, and time. Radiation was determined by temperature, cloud (aerosol), composition, and surface emissivity measurements. The theory of radiative transfer is reviewed, along with retrieval methods. Instruments onboard space probes for infrared remote sounding of planetary atmospheres are also described. J.A.M.

N75-12283# Research Inst. of National Defence, Stockholm

RECORDINGS OF THE SPECTRAL-RADIO-METER SIGNE DURING THE SUMMER OF 1972 [REGISTRERINGAR MED SPEKTRORADIOMETERN 'SIGNE' UNDER SOMMAREN 1972]

Carl Hugo Agren Nov. 1972 22 p ref in SWEDISH (FOA-2-C-2577-E1) Avail: NTIS HC \$3.25

A technical description of the spectral radiometer is given that is intended for measuring at a distance. An account of its first recordings is presented together with its construction, calibration and operation. Practical problems in measuring are dealt with as well as different factors which effect measurements. In conclusion, a number of recordings are given of the spectral radiance and spectral reflections, mainly of vegetation Author and water.

N75-12410*# Colorado School of Mines, Golden. Dept. of INDEX OF SKYLAB DATA AVAILABLE AT COLORADO

SCHOOL OF MINES

Keenan Lee, Principal Investigator and Gary L. Prost 31 Oct. 1974 32 p Original contains imagery. Original photography may be purchased from the EROS Data Center, 10th and Dakota Avenue, Sioux Falls, S. D. 57198 EREP (Contract NAS9-13394)

(E75-10041; NASA-CR-140778; Rept-74-8) Avail: NTIS HC \$3.75 CSCL 05B

N75-12411*# Long Island Univ., Greenvale, N.Y. Engineering Research Group. IN SITU SPECTRORADIOMETRIC CALIBRATION OF EREP IMAGERY AND OCEANOGRAPHY OF BLOCK ISLAND SOUND Progress Report, Apr. - Nov. 1974 E. Yost, Principal Investigator 15 Nov. 1974 6 p EREP (Contract NAS9-13308) (E75-10042; NASA-CR-140779) Avail: NTIS HC \$3.25 CSCL

N75-12418*# ESL, Inc., Sunnyvale, Calif. EARTH RESOURCES SENSOR DATA HANDLING SYSTEM: NASA JSC VERSION Final Design Report 29 Apr. 1974 23 p (Contract NAS9-14077)

(NASA-CR-140334; ESL-SR-132) Avail: NTIS HC \$3.25 CSCL

The design of the NASA JSC data handling system is presented. Data acquisition parameters and computer display formats and the flow of image data through the system, with recommendations for improving system efficiency are discussed along with modifications to existing data handling procedures which will allow utilization of data duplication techniques and the accurate identification of imagery.

N75-13366*# Purdue Univ., Lafayette, Ind. Lab. for Application of Remote Sensing. SPECTRORADIOMETRIC AND MULTISPECTRAL SCANNER OVER LAKE MONROE Monthly Report, Oct. 1974 LeRoy F. Silva, Principal Investigator Oct. 1974 2 p EREP (Contract NAS9-13301) (E75-10066; NASA-CR-141006) Avail: NTIS HC \$3.25 CSCL

N75-13455# Environmental Data Service, Washington, D.C. CATALOG OF EARTHQUAKE PHOTOGRAPHS Key to Geophysical Records, Documentation No. 3 Jerry L. Coffman, comp. Nov. 1974 46 p. Avail: NTIS HC \$3.75

This catalog represents a beginning effort to systematically collect photographs and maps from many sources and to make them generally available. This publication lists approximately 300 photographs and 200 maps on the subject of earthquakes and contains descriptions and examples of the collection. Many of the photographs show views of earthquake-damaged buildings. bridges, dams, and other man-made structures; several show spectacular landslides, rifts, and ground cracks resulting from destructive shocks. The maps depict intensity distribution of important earthquakes and the seismicity of areas throughout the world. Author

N75-14199 *# Kansas Univ. Center for Research, Inc., Lawrence. Remote Sensing Lab.

DESIGN DATA COLLECTION WITH SKYLAB/EREP MI-CROWAVE INSTRUMENT S-193 Monthly Letter Progress

Report No. 13, Oct. 1974 Arun Sobti Oct. 1974 30 p EREP (Contract NAS9-13331)

(E75-10080; NASA-CR-141104) Avail: NTIS HC \$3.75 CSCL

The author has identified the following significant results. The Skylab S-193 radiometer/scatterometer collected thousands of measurements of scattering coefficient and brightness temperature over various parts of the United States during the summer of 1973 at angles of incidence between vertical and about 45 deg. These measurements have been combined to produce histograms of the response at each of several angles within this range, and to establish average scattering coefficient vs angle curves with 10% and 90% exceedance levels as well. The variation of the radiometeric measurements is primarily in the region from 255 K to 285 K, with very few measurements giving values down to and even below 200 K. The scattering coefficient varies, for the mean, from about 0 db at 1 deg off vertical to a low in the neighborhood of -10 db at 45 deg. The variability of the scattering coefficient measurements with this coarse resolution sensor is surprisingly small. The number of distinguishable levels is slightly more for the scatterometer than for the radiometer, but the amount of variation in brightness temperature caused by the physical temperature of the ground is enough so that the scatterometer can be used to distinguish significantly more meaningful levels than the radiometer.

N75-14200 # Environmental Research and Technology, Inc., Lexington Mass.

STUDY TO DEVELOP IMPROVED SPACECRAFT SNOW SURVEY METHODS USING SKYLAB/EREP DATA Quarterly Progress Report, 15 Sep. - 15 Dec. 1974

James C. Barnes, Principal Investigator Dec. 1974 5 p EREP (Contract NAS9-13305)

(E75-10081; NASA-CR-141105; P-412-10; QPR-7) Avail: NTIS HC \$3.25 CSCL 08L

N75-14236 Consiglio Nazionale delle Ricerche, Milan (Italy). Lab. per la Geofisica della Litosfera.

CONTRIBUTION OF SPACE PLATFORMS TO A GROUND AND AIRBORNE REMOTE-SENSING PROGRAMME OVER ACTIVE ITALIAN VOLCANOES

R. Cassinis, G. M. Lechi, and A. M. Tonelli In ESRO European Earth-Resources Satellite Expts. May 1974 p 185-197 Sponsored by NASA

ERTS-1 imagery of the volcanic areas of southern Italy was used primarily for the evaluation of space platform capabilties in the domains of regional geology, soil and rock-type classification and, more generally, to study the environment of active volcanoes. The test sites were selected and equipped primarily to monitor thermal emission, but ground truth data was also collected in other domains (reflectance of rocks, soils and vegetation). The test areas were overflown with a two channel thermal scanner. while a thermo camera was used on the ground to monitor the hot spots. The primary goal of this survey was to plot the changes in thermal emission with time in the framework of a research program for the surveillance of active volcanoes. However, another task was an evaluation of emissivity changes by comparing the outputs of the two thermal channels. These results were compared with the reflectance changes observed on multispectral ERTS-1 Author (ESRO)

N75-14870# Army Material Command, Texarkana, Tex. Intern Training Center.

LIFE CYCLE COST STUDY OF ARMY SPECTROMETRIC OIL ANALYSIS PROGRAM (ASOAP) Final Report

Henry L. Martin Apr. 1974 59 p refs

(AD-786501; USAMC-ITC-02-08-73-018) Avail: NTIS CSCL 14/2

The scope of this paper is to determine the present life cycle cost (LCC) of ASOAP (Army Spectrometric Oil Analysis Program) based on certain assumptions. A cost model capable of determining the life cycle cost of ASOAP for both the atomic absorption spectrometry and the atomic emission spectrometry is developed. Next, using this cost model, the feasibility of the oil analysis program is determined. Finally, interest is focused on the benefits that the program has brought about in the past, and future benefits will be projected. A brief history of the Army's intervention into oil analysis, LCC, and cost models is presented. A brief discussion on the two methods of analysis that the Army uses and some estimations of various costs that are associated with ASOAP are included. A general cost model and a cost model for ASOAP is developed with a discussion of the various components. (Modified author abstract)

N75-15113*# Purdue Univ., Lafayette, Ind. Lab. for Applications of Remote Sensing.

ANALYSIS OF SKYLAB MSS DATA, AND CALIBRATION OF THERMAL DATA OVER LAKE MONROE, INDIANA Monthly Report, Dec. 1974

LeRoy F. Silva, Principal Investigator Dec. 1974 2 p EREP (Contract NAS9-13301)

(E75-10095; NASA-CR-141291) Avail: NTIS HC \$3.25 CSCL 08H

The author has identified the following significant results. The SL/4-S192 data over northern Indiana looks good. The thermal band looks very encouraging. The preliminary analysis indicates so far that it will be difficult to distinguish some land use classes because of the lack of vegetation. Preliminary analysis of the thermal band in particular indicates that there appears to be a correlation of the hot spots in the data over the Ft. Wayne, Indiana area and its industrial centers.

N75-15119*# National Oceanic and Atmospheric Administration, Rockville, Md. Coastal Mapping Div.

SKYLAB: A PROPOSAL AEROTRIANGULATION WITH VERY SMALL SCALE PHOTOGRAPHY Quarterly Report, 15 Oct. 1974 - 15 Jan. 1975

Morton Keller, Principal Investigator 15 Jan. 1975 1 p

(NASA Order T-4110-B)

(E75-10101; NASA-CR-141297) Avail: NTIS HC \$3.25 CSCL 08B

N75-15122*# Purdue Univ., Lafayette, Ind. Lab. for Applications of Remote Sensing.

AN INTERDISCIPLINARY ANALYSIS OF MULTISPECTRAL SATELLITE DATA FOR SELECTED COVER TYPES IN THE COLORADO MOUNTAINS, USING AUTOMATIC DATA PROCESSING TECHNIQUES Monthly Progress Report, Dec. 1974

Roger M. Hoffer, Principal Investigator Dec. 1974 11 p

(Contract NAS9-13380)

(E75-10104: NASA-CR-141305) Avail: NTIS HC \$3.25 CSCL 08F

The author has identified the following significant results. In the San Juan Mountains there exists a group of structural and topographic features which suggest a common origin. The relative positions of the San Juan Depression with the Silverton and Lake City Calderas indicate they are possibly on the edge of a large circular feature. Comparison with the 1:250,000 scale Durango Geologic Map reveals a series of radial and arcuate faults concentric with the major circular feature. Such a fracture pattern implies that the circular feature may represent the extent of a major domal uplift in the area. The uplift was very likely due to the sub-crustal emplacement of the parent magma of the San Juan volcanics and intrusives. The broad doming led to the formation of a zone of tensional ring fractures which acted as conduits of release for the magma and the eventual development of the various calderas. The ring fracturing was probably incomplete with hinging occurring in the southern and southwestern portions of the dome accounting for the lack of development of caldera structures in that portion of the area.

N75-15131*# Environmental Research Inst. of Michigan, Ann Arbor. Infrared and Optics Div.

INVESTIGATION OF TRANSIENT EARTH RESOURCES PHENOMENA: CONTINUATION STUDY Final Report, 31 Jan. - 30 Sep. 1974

Gary C. Goldman et al Nov. 1974 125 p refs

đ

(Contract NAS5-20021)

(NASA-CR-139163: ERIM-107400-2-F) Avail: NTIS HC \$5.25 CSCL 08E

Calculated sensitivity requirements for an earth resource satellite in a geostationary orbit are reported. Radiance levels at the satellite sensor were computed for twenty top-priority Synchronous Earth Observatory Satellite (SEOS) applications. The observation requirements were reviewed and re-evaluated in terms of spectral band definition, spectral signatures of targets and backgrounds, observation time, and site location. With these data and an atmospheric attenuation and scattering model, the

08: INSTRUMENTATION AND SENSORS

total radiances observed by the SEOS sensor were calculated as were the individual components contributed by the target, target variations, and the atmosphere.

N75-15136*# Scientific Translation Service, Santa Barbara, Calif. FUNDAMENTALS OF THE SPACE INDICATION OF EARTH RESOURCES

V. V. Andrevanov Washington NASA Jan. 1975 62 p Transl. into ENGLISH from the Russian report D-171 (Contract NASw-2483)

(NASA-TT-F-16108; D-171) Avail: NTIS HC \$4.25 CSCL 08E

The methods and instrumentation used in satellite reconnaissance of earth features are reviewed. The electromagnetic spectrum of solar radiation is analyzed along with the atmospheric effects on the transmission of visible and infrared radiation. Statistical methods are given for the identification of natural features and the determination of their characteristics. Remote sensing and data transmission requirements for satellites and the affect of orbit parameters on the results and condition of measurements is given. The principles of operation and the characteristics of photographic and television cameras, scanning radiometers, and active radar methods for the investigation of earth's resources are discussed. Author

N75-15138*# Long Island Univ., Greenvale, N.Y. Science Engineering Research Group.

MULTISPECTRAL PHOTOGRAPHY FOR EARTH SOURCES

Sondra Wenderoth, Edward Yost, Rajender Kalia, and Robert Anderson 1972 279 p refs Original contains color illustra-

(Contract NAS9-11188)

(NASA-CR-141283) Avail: NTIS HC \$8.75 CSCL 14E

A guide for producing accurate multispectral results for earth resource applications is presented along with theoretical and analytical concepts of color and multispectral photography. Topics discussed include: capabilities and limitations of color and color infrared films; image color measurements; methods of relating ground phenomena to film density and color measurement; sensitometry; cosiderations in the selection of multispectral cameras and components; and mission planning.

N75-15333*# Rice Univ., Houston, Tex. Inst. for Computer Services and Applications.

EARTH RESOURCES DATA ANALYSIS PROGRAM. PHASE 2 Final Report, Jun. 1973 - May 1974 Jun. 1974 38 p ref

(Contract NAS9-12776)

(NASA-CR-141477; Rept-275-025-FP2) NTIS

HC \$3.75 CSCL 09B

The efforts and findings of the Earth Resources Data Analysis Program are summarized. Results of a detailed study of the needs of EOD with respect to an applications development system (ADS) for the analysis of remotely sensed data, including an evaluation of four existing systems with respect to these needs are described. Recommendations as to possible courses for EOD to follow to obtain a viable ADS are presented. Algorithmic development comprised of several subtasks is discussed. These subtasks include the following: (1) two algorithms for multivariate density estimation; (2) a data smoothing algorithm; (3) a method for optimally estimating prior probabilities of unclassified data; and (4) further applications of the modified Cholesky decomposition in various calculations. Little effort was expended on task 3, however, two reports were reviewed. Author

09 GENERAL

Includes economic analysis.

A75-12030 * # Planning and management of science programs on Skylab. R. A. R. Parker and J. R. Sevier (NASA, Johnson Space Center, Houston, Tex.). American Institute of Aeronautics and Astronautics and American Geophysical Union, Conference on Scientific Experiments of Skylab, Huntsville, Ala., Oct. 30-Nov. 1, 1974, AIAA Paper 74-1215. 6 p.

Discussion of the experience gained in experiment operation planning during the Skylab mission. The Skylab flight planning activity allowed the experimenters to interact with the system and provided the flexibility to respond to contingencies both major and minor. Both these aspects contributed to make efficient use of crew time thus helping to increase the science return from the mission. Examples of the need for real time scheduling response and of the tradeoffs considered between conflicting experiment requirements are presented. General management principles derived from this experience are developed. The Skylab mission experiences, together with previous Apollo mission experiences, are shown to provide a good background for Shuttle flight planning.

A75-12986 Energy development; Proceedings of the Energy Sources Conference, Anaheim, Calif., July 14-19, 1974. Conference sponsored by the Institute of Electrical and Electronics Engineers. New York, Institute of Electrical and Electronics Engineers, Inc., 1974. 104 p. Members, \$7.50; nonmembers, \$10.00.

The impact of advanced batteries on electric power generation, tidal power and its integration into the electric system, and utilization of municipal refuse for the recovery of energy and other resources are among the topics covered in papers concerned with the latest developments in power engineering research. Other topics covered include current expectations for fusion power from toroidal machines, progress in coal gasification, and pumped air storage for electric power generation.

M.V.E.

A75-12999 U.S. energy resources - Outlook for the future. R. A. Budenholzer (Illinois Institute of Technology, Chicago, III.). In: Energy development; Proceedings of the Energy Sources Conference, Anaheim, Calif., July 14-19, 1974. New York, Institute of Electrical and Electronics Engineers, Inc., 1974, p. 98-102. 5 refs.

Assessment of the extent and nature of the present energy crisis, and search for courses of action that could alleviate the situation on both a short-term and long-term basis. Reviewed estimates of the number of years of supply remaining for each of the various fuel resources in the United States show that coal and uranium are much more abundant than crude oil and natural gas. For the immediate future, this suggests a gradual shift of all power generation to coal and nuclear energy, thus conserving oil and gas for higher priority uses. For the long range future, it is hoped that fusion power can be achieved commercially.

M.V.E.

A75-13581 ERTS-1 and data for national land use planning. J. R. Anderson (U.S. Geological Survey, Reston, Va.). In: Technology today for tomorrow; Proceedings of the Eleventh Space Congress, Cocoa Beach, Fla., April 17-19, 1974. Volume 1.

Cape Canaveral, Fla., Canaveral Council of Technical Societies, 1974, p. 3-25 to 3-28. 8 refs.

An Earth Resources Technology Satellite can provide information about existing land use at a relatively generalized level of presentation on a timely repetitive basis. A land use classification system has been developed to meet the needs for an up-to-date overview of land use throughout the country. The ERTS-1 capabilities for obtaining required land use data are discussed. It has been found that land use maps of a large scale can be produced by the machine processing of ERTS-1 scanner data.

G.R.

A75-13592 * NASA-KSC's earth resources benefits from space exploration. J. M. Ragusa and E. J. Hecker (NASA, Kennedy Space Center, Earth Resources Branch, Cape Canaveral, Fla.). In: Technology today for tomorrow; Proceedings of the Eleventh Space Congress, Cocoa Beach, Fla., April 17-19, 1974. Volume 2.

Cape Canaveral, Fla., Canaveral Council of Technical Societies, 1974, p. 2-9 to 2-14.

The purpose of this paper is to identify the nature and scope of earth resources activities at the Kennedy Space Center (KSC). Because of recent developments from space exploration, NASA and KSC have evolved an earth resources program which focuses on applied R&D activities of direct benefit to a variety of federal, state, and local users. The historical development of this program is traced, and several projects are identified with special emphasis on the use of sidelooking airborne radar in several Brevard County, Florida test areas. (Author)

A75-13618 * Skylab contributions to the future. W. C. Schneider and J. H. Disher (NASA, Office of Manned Space Flight, Washington, D.C.). International Astronautical Federation, International Astronautical Congress, 25th, Amsterdam, Netherlands, Sept. 30-Oct. 5, 1974, Paper. 57 p.

Skylab showed that a multidisciplinary manned space station is practical and can be highly productive. Quality data was produced concerning earth resources (mineral resources, agriculture, climatology, etc.), solar astronomy (solar activity, magnetic fields, and solar winds), materials sciences (e.g., zero-gravity crystal growth), and astronomy and astrophysics (including observations of comet Kohoutek). Two types of astronaut maneuvering unit were tested. Medical data are applicable to adaptational environmental physiology.

A.T.S.

A75-13761 The potential benefits for development derived from space applications. V. Skatchkov (United Nations, Outer Space Affairs Div., New York, N.Y.). International Astronautical Federation, International Astronautical Congress, 25th, Amsterdam, Netherlands, Sept. 30-Oct. 5, 1974, Paper 74-147. 16 p. 12 refs.

Potential benefits from using space systems for communications, navigation, meteorology, and remote sensing of earth resources are considered. Developing countries will derive the greatest benefit from remote sensing from space as applied to agriculture, fisheries, pollution and erosion monitoring, exploration for minerals, etc. Great cost effectiveness of such programs is predicted. Centralized international ground segments for the systems is recommended. System costs and possible sources of investment are examined.

A.T.S.

A75-13770 Economic effectiveness of using the space systems of distant indication of natural resources. S. A. Sarkisian, E. S. Minaev, S.-S. Korunov, and O. U. Kurdinovskii. International Astronautical Federation, International Astronautical Congress, 25th, Amsterdam, Netherlands, Sept. 30-Oct. 5, 1974, Paper 74-165. 26 p.

Remote sensing of natural resources from space can be used in various areas of interest to the national economy, such as geology, forestry, agriculture, water economy, fisheries, etc. Users' economy requirements are most important in devising such space systems. Equations are derived for calculating the total economic effect of substituting space-based systems for tradiational means. A technique

for estimating economic effectiveness is outlined. The example of the fishing industry indicates that reduction of the search fleet, increased catches, exploitation of additional ocean areas, and decreased searching time are possible through use of remote sensing from space.

A75-15319 # The radio emission of the earth as a planet (Radioizluchenie zemli kak planety). A. E. Basharinov, A. S. Gurvich, and S. T. Egorov. Moscow, Izdatel'stvo Nauka, 1974. 188 p. 185 refs. In Russian.

The present work sets forth the basic principles of measurements of the earth's thermal microwave radiation, with special emphasis on satellite techniques. The characteristics of the formation of microwave radiation in the system comprised of the atmosphere and the earth's surface are studied. Aspects of this kind of investigation having practical importance are covered, such as the determination of air humidity, cloud water content, glacier compactness, and the humidity and temperature of land masses. P.T.H.

A75-17103 # Estimation of population density in Tokyo districts from ERTS-1 data. S. Murai (Tokyo, University, Tokyo, Japan). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 13-22.

Study of the feasibility of estimating population density in Tokyo districts by using the multispectral characteristics of urbanized areas, low populated areas, such as forest or crop field, and unpopulated areas, such as seas, lakes, or mountains, in the two-dimensional signature space of MSS 5 and MSS 7. Twodimensional histograms have been acquired for four different classes of population density, and the correlation between density signals of 9.5 in, transparencies of MSS 5 and MSS 7, and population density data taken from the census in 1965, has been analyzed for Tokyo districts consisting of 180 by 160 grids of 1 kilometer interval. From this correlation, seven decision regions have been established in the two-dimensional signature space of MSS 5 and MSS 7. The accuracy of estimation of population density is approximately 90%, in spite of the existence of misregistration and low resolution due to the use of a 1-mm aperture in the density measurement for the 9.5 in. transparencies. Misregistered points and unresolved points near boundaries of water bodies have been eliminated. The accuracy will be improved if video signals stored in the MSS digital tape are available.

A75-17118 # Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data. H. W. Smedes (U.S. Geological Survey, Denver, Colo.), G. Nez (Federation of Rocky Mountain States, Denver, Colo.), A. K. Turner (Colorado School of Mines, Golden, Colo.), J. C. Reed, Jr. (U.S. Geological Survey, Reston, Va.), and E. E. Lutzen. In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 289-298.

A75-17123 # ERTS applications in Thailand - A progress report. P. Angsuwatana (Ministry of Industry, Dept. of Mineral Resources), C. Boonyobhas, B. Klankamsorn, M. Omakupt, P. Piyapongse (Ministry of Agriculture and Cooperatives), J. Morgan (U.S. Geological Survey, Washington, D.C.), K. Suvarnasudhi, and S. Vibulsresth (Applied Scientific Research Corporation of Thailand,

Bangkok, Thailand). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 1. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 341-361.

Analyses of ERTS-1 imagery conducted by four Royal Thai Government agencies responsible for survey, inventory, and development of natural resources are described. Visual interpretation of single band and color composite images was augmented and verified by ground checks and aerial photographs. Preliminary computer printouts of one scene have been found accurate and useful. Agriculture, forestry, land use, and geological mapping are illustrated. The work will be reported in more detail to NASA (SR No. 9620). (Author)

A75-17166 * # The economic impact of remote sensing data as the source of nonpoint pollution monitoring and control. W. L. Miller (Purdue University, West Lafayette, Ind.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 2.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1001-1011. 13 refs. Grant No. NGL-15-005-112.

Nonpoint pollution of streams with sediment as a result of runoff from alternative uses of land has become a socially unacceptable product of economic activity. This report describes a research approach to economically achieve correction of the nonpoint pollution problem. The research approach integrates the economic model with those data which may be obtainable from remotely sensed sources. The economic problem involves measurement of the direct benefits and costs associated with the changes in land management activities necessary to reduce the level of nonpoint pollution. Remotely sensed data from ERTS-1 may provide some of the information required for the economic model which indicates efficient solutions to the nonpoint pollution problem. Three classes of data (i.e., soil categories, vegetative cover, and water turbidity) have the potential to be measured by ERTS-1 systems. There is substantial research which indicates the ability of ERTS-1 to measure these classes of data under selected conditions. (Author)

A75-17211 * # Summary of flight performance of the Skylab Earth Resources Experiment Package /EREP/. A. E. Potter, C. K. Williams, A. L. Grandfield, K. J. Demel, M. C. Trichel, T. L. Barnett, R. D. Juday, W. E. Hensley, N. M. Hatcher, and W. E. McAllum (NASA, Johnson Space Center, Houston, Tex.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 1803-1846. 17 refs.

A group of six remote sensor systems (sensing visible, infrared, and microwave radiation) known as the Earth Resources Experiment Package (EREP) was flown on the NASA Skylab spacecraft to furnish data to numerous investigators in the earth sciences and in technology assessment. Inflight sensor performance in three categories (functional, geometric, and radiometric) was evaluated using: (1) ground measurements of surface, atmospheric, and illumination parameters; (2) ground deployment and operation of microwave receivers and transponders to monitor and excite the active EREP sensors; (3) measurement of surface and atmospheric parameters by instrumented aircraft on underflights of Skylab passes; and (4) analysis of the actual flight data. This resulted in identification and correction of anomalous sensor operation, quantization of geometric distortions or aberrations, improvement or confirmation of calibrations, and determination of sensitivity, accuracy, and stability of the sensors. (Author)

A75-17235 # An overview of Texas activities in remote sensing. R. N. Neece (General Land Office, Austin, Tex.), M. Ellis (Texas Water Development Board, Austin, Tex.), and J. Wells

(Interagency Council on Natural Resources and the Environment, Austin, Tex.). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3.

Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 2107-2112.

A75-17236 * # A summary of ERTS data applications in Alaska. J. M. Miller and A. E. Belon (Alaska, University, Fairbanks, Alaska). In: International Symposium on Remote Sensing of Environment, 9th, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volume 3. Ann Arbor, Mich., Environmental Research Institute of Michigan, 1974, p. 2113-2138. 14 refs. NASA-supported research.

ERTS has proven to be an exceedingly useful tool for the preparation of urgently needed resource surveys in Alaska. For this reason the wide utilization of ERTS data by federal, state and industrial agencies in Alaska is increasingly directed toward the solution of operational problems in resource inventories, environmental surveys, and land use planning. Examples of some applications are discussed in connection with surveys of potential agricultural lands; mapping of predicted archaeological sites; permafrost terrain and aufeis mapping; snow melt enhancement from Prudhoe Bay roads; geologic interpretations correlated with possible new petroleum fields, with earthquake activity, and with plate tectonic motion along the Denali fault system; hydrology in monitoring surging glaciers and the break-up characteristics of the Chena River watershed; sea-ice morphology correlated with marine mammal distribution; and coastal sediment plume circulation patterns.

(Author)

N75-10533*# National Research Council, Bangkok (Thailand).
THAILAND NATIONAL PROGRAMME OF THE EARTH
RESOURCES TECHNOLOGY SATELLITE

Pradisth Cheosakul, Principal Investigator Aug. 1973 14 p Sponsored by NASA ERTS (E74-10806; NASA-CR-140500) Avail: NTIS HC \$3.25 CSCL 08F

N75-10534*# National Research Council, Bangkok (Thailand). THE THAILAND PROGRAMME OF THE EARTH RESOURCES TECHNOLOGY SATELLITE Progress Report Pradisth Cheosakul, Principal Investigator [1974] 3 p Sponsored by NASA ERTS (E74-10807; NASA-CR-140501; PR-1) Avail: NTIS HC

N75-10546*# General Electric Co., Philadelphia, Pa. Space Div.

ERTS 1 FLIGHT EVALUATION REPORT, 23 OCTOBER 1973 - 23 JANUARY 1974

26 Feb. 1974 123 p (Contract NAS5-21808)

\$3.25 CSCL 08F

(NASA-CR-140712; Doc-74SD4205). Avail: NTIS HC \$5.25 CSCL 05A

Detailed analyses are given of flight performance of ERTS-1 systems and subsystems during orbits 6371 to 7651. A.L.

N75-10547*# General Electric Co., Philadelphia, Pa. Space

ERTS 1 FLIGHT EVALUATION REPORT, 23 JANUARY -23 APRIL 1974

18 May 1974 141 p

(Contract NAS5-21808)

(NASA-CR-140810; DOC-74SD4217) Avail: NTIS HC \$5.75 CSCL 05A

Detailed analyses are given of flight performance of ERTS-1 systems and subsystems during orbits 7652 to 8907. A.L.

N75-10548*# General Electric Co., Philadelphia, Pa. Space

ERTS 1 FLIGHT EVALUATION REPORT, JANUARY 23, 1973 TO APRIL 23, 1973

29 May 1973 196 p

(Contract NAS5-21808)

(NASA-CR-140809; Doc-73SD4249) Avail: NTIS HC \$7.00 CSCL 22C

Detailed analyses are given of the performance of systems and subsystems on board ERTS-1 during orbits 2600-3810.

A.L

N75-10549*# National Aeronautics and Space Administration.
Goddard Space Flight Center, Greenbelt, Md.

THIRD EARTH RESOURCES TECHNOLOGY SATELLITE SYMPOSIUM. VOLUME 2: SUMMARY OF RESULTS

Stanley C. Freden, ed., Enrico P. Mercanti, ed., and David B. Friedman, ed. Washington May 1974 184 p refs Symp. held at Washington, D. C., 10-14 Dec. 1973 Original contains color illustrations

(NASA-SP-356) Avail: NTIS MF \$2.25; SOD HC \$3.00 CSCL 05B

Summaries are provided of significant results taken from presentations at the symposium along with some typical examples of the applications of ERTS-1 data for solving resources management problems at the national, state, and local levels.

N75-10550* Canada Centre for Remote Sensing, Ottawa (Ontario).

THE CANADIAN ERTS PROGRAM

Lawrence W. Morley In NASA. Goddard Space Flight Center Third ERTS Symp., Vol. 2 May 1974 p 7-11

CSCL 05B

Under a four-year collaborative agreement between the United States and Canada, the Canada Centre for Remote Sensing (CCRS) reads out and distributes the ERTS data of Canada. The Canadian receiving station is at Prince Albert, Saskatchewan, and covers most of Canada except for the East Coast. Production statistics on Canadian ERTS imagery, a summary of several cost benefit case histories, and recommendations for the future of the international aspects of ERTS are discussed and evaluated.

Author

N75-10568*# General Electric Co., Philadelphia, Pa. Space Div.

ERTS 1 FLIGHT EVALUATION REPORT, 23 APRIL 1974-23 JULY 1974

15 Aug. 1974 219 p

(Contract NAS5-21808)

(NASA-CR-140711; Doc-74SD4236) Avail: NTIS HC \$7.25 CSCL 05A

Detailed analyses are given of flight performance of ERTS-1 systems and subsystems during orbits 8908 to 10182. A.L.

N75-10617# Committee on Interior and Insular Affairs (U. S. Senate).

NATIONAL ENVIRONMENTAL POLICY ACT OF 1989. ENVIRONMENTAL INDICES, STATUS OF DEVELOPMENT PURSUANT TO SECTIONS 102(2)(B) AND 204 OF THE ACT

James W. Curlin Washington GPO 1973 54 p refs Presented to Comm. on Interior and Insular Affairs, 93d Congr., 1st Sess., Dec. 1973 Prepared by the Library of Congr., Environ. Policy Div.

(GPO-25-314) Avail: SOD HC \$0.55

Developments in the design of environmental indices for monitoring the environment are reported. Topics discussed include: indicators and indices, environmental components, obstacles to the development of indices and indicators, air quality index, land use index, and noise pollution index.

F.O.S.

N75-10967*# National Aeronautics and Space Administration. Marshall Space Flight Center, Huntsville, Ala.

MSFC INTEGRATED EXPERIMENTS PRELIMINARY REPORT

Nov. 1974 146 p refs

(NASA-TM-X-64881) Avail: NTIS HC \$5.75 CSCL 22C

Skylab experiments are described and their preliminary results are reported. The types of experiments described include medical, earth resources, space physics, space manufacturing, and spacecraft design.

Author

N75-11406*# Catholic Univ. of America, Washington, D.C. Dept. of Anthropology.

EARTH RESOURCES TECHNOLOGY SATELLITE-1 (ERTS-1)
DATA AND ANTHROPOLOGY: USE OF THESE DATA IN
CARRYING CAPACITY ESTIMATES FOR SITES IN UPPER
VOLTA AND NIGER

Priscilla Reining, Principal Investigator 24 Nov. 1974 10 p Presented at the Amer. Anthropological Assoc. Meeting, Mexico City, 24 Nov. 1974 ERTS

(Contract NAS5-21970)

(E75-10001; NASA-CR-140687) Avail: NTIS HC \$3.25 CSCL

The author has identified the following significant results. Repetitively derived multispectral band imagery from ERTS-1 is now available for many parts of the earth's land surface and represents major new data sources for anthropological work in habitat, land use, and settlement patterns. A completed first step test of ERTS-1 data is available in carrying capacity estimates for Mossi, Hausa, and Sonrai sites derived from: (1) field work; (2) aerial photography; and (3) ERTS-1. Data can test more than one carrying capacity formula.

N75-11414*# Mississippi State Univ., State College. APPLICATION OF REMOTE SENSING TO STATE AND REGIONAL PROBLEMS Semiannual Progress Report, 1 May - 31 Oct. 1974

W. Frank Miller, C. W. Bouchillon, J. C. Harris, Brad Carter, Frank Whisler, and Randell Robinette 31 Oct. 1974 24 p refs

(Grant NGL-25-001-054)

(NASA-CR-140805; SAPR-2) Avail: NTIS HC \$3.25 CSCL 08B

The primary purpose of the remote sensing applications program is for various members of the university community to participate in activities that improve the effective communication between the scientific community engaged in remote sensing research and development and the potential users of modern remote sensing technology. Activities of this program are assisting the State of Mississippi in recognizing and solving its environmental, resource and socio-economic problems through inventory, analysis, and monitoring by appropriate remote sensing systems. Objectives, accomplishments, and current status of the following individual projects are reported: (1) bark beetle project; (2) state park location planning; and (3) waste source location and stream channel geometry monitoring.

N75-11549*# California Univ., Los Angeles.

PROCEEDINGS OF THE UCLA INTERNATIONAL CONFERENCE ON RADIATION AND REMOTE PROBING OF THE ATMOSPHERE

Jacob G. Kuriyan, ed. 1974 422 p refs Conf. held at Los Angeles, 28-30 Aug. 1973 Sponsored by DOT, NASA, TRW Systems, and Army

(NASA-CR-140816) Avail: NTIS HC \$10.50 CSCL 04A

Various articles are presented on multiple scattering problems and radiative transfer in the atmosphere. Particle size distribution and molecular absorption are also discussed.

THE FINAL SKYLAB MISSION: MAN AT HOME AND AT WORK IN SPACE

15 Feb. 1974 8 p

(NASA-TM-X-70394; MR-15) Avail: NTIS HC \$3.25 CSCL 22C

The accomplishments of the Skylab 4 mission are discussed. The medical experiments and dietary aspects of the mission are reported. The observation of the Comet Kohoutek is described. The remote sensing of earth resources is examined to show the areas of coverage. The repair of the space station and the accomplishment of unscheduled requirements are discussed. Statistical data of all the Skylab missions are tabulated.

N75-12390*# Purdue Univ., Lafayette, Ind. Lab. for Applications of Remote Sensing.

AN INTERDISCIPLINARY ANALYSIS OF MULTISPECTRAL SATELLITE DATA FOR SELECTED COVER TYPES IN THE COLORADO MOUNTAINS, USING AUTOMATIC DATA PROCESSING TECHNIQUES Monthly Progress Report, Oct. 1974

Roger M. Hoffer, Principal Investigator Oct. 1974 7 p EREP (Contract NAS9-13380)

(E75-10002; NASA-CR-140688) Avail: NTIS HC \$3.25 CSCL

The author has identified the following significant results. Documentation is presented of the capability of the middle infrared portion of the electromagnetic spectrum to spectrally differentiate clouds from snow. Other portions of the spectrum cannot provide this capability.

N75-12416# Committee on Science and Astronautics (U. S. House).

EARTH RESOURCES SURVEY SYSTEM

Washington GPO 1974 286 p refs Hearings on H.R. 14978 and H.R. 15781 before Subcomm. on Space Sci. and Appl. of Comm. on Sci. and Astronaut., 93d Congr., 2d Sess., No. 46, 3, 4, and 9 Oct. 1974

(GPO-41-665) Avail: Subcomm. on Space Sci. and Appl.

The hearings to explore the desirability of creating an operational satellite system for surveying the earth's resources are reported. The ERTS program is reviewed, and the technology for data acquisition, dissemination, and utilization are assessed along with remote sensing techniques.

F.O.S.

N75-12424*# ECON, Inc., Princeton, N.J.

IMPROVED (ERTS) INFORMATION AND ITS IMPACT ON U.S. MARKETS FOR AGRICULTURAL COMMODITIES: A QUANTITIATIVE ECONOMIC INVESTIGATION OF PRODUCTION, DISTRIBUTION AND NET EXPORT EFFECTS 31 Oct. 1974 154 p refs

(Contract NASw-2558)

(NASA-CR-141044; Rept-74-2001-6) Avail: NTIS HC \$6.25 CSCL 05B

An econometric investigation into the markets for agricultural commodities is summarized. An overview of the effort including the objectives, scope, and architecture of the analysis and the estimation strategy employed is presented. The major empirical results and policy conclusions are set forth. These results and conclusions focus on the economic importance of improved crop forecasts, U.S. exports, and government policy operations. A number of promising avenues of further investigation are suggested.

N75-13010# Committee on Aeronautical and Space Sciences (U. S. Senate).

EARTH RESOURCES SATELLITES

Washington GPO 1974 341 p refs Hearings on S. 2350 and S. 3484 before Comm. on Aeron. and Space Sci., 93d Congr., 2d Sess., 6, 8, and 9 Aug. and 18 Sep. 1974 (GPO-40-560) Avail: SOD HC \$3.00

The arguments for and against the expansion of earth resources satellite capabilities and preparation for an operational satellite system were presented in a hearing before Congress whose purpose was to gather information on two impending

bills which would increase the federal government's activities in remote sensing programs. Significant accomplishments of ERTS 1 are reviewed, along with some potential capabilities which are as yet unexplored. The ways in which ERTS data is being used are also described, with emphasis on the role of the EROS Data Center in Sioux Falls, S. Dak. in disseminating information of use to farmers, environmental experts, regional planners, and researchers. Plans for an operational earth resources satellite system, which is expected to become a more pressing issue with the launch of ERTS-B, are discussed in terms of public vs. private control and the possibility that such a system will present difficulties in foreign relations. A.A.D.

N75-13370*# National Aeronautics and Space Administration, Washington, D.C.

ERTS-B (EARTH RESOURCES TECHNOLOGY SATELLITE) 14 Jan. 1975 74 p

(NASA-News-Release-74-329) Avail: NASA Scientific and Technical Information Facility, P. O. Box 33, College Park, Md. 20740 CSCL 05B

Mission plans and objectives of the ERTS 2 Satellite are presented. ERTS 2 follow-on investigations in various scientific disciplines including agriculture, meteorology, land-use, geology, water resources, oceanography, and environment are discussed. Spacecraft design and its sensors are described along with the Delta launch vehicle and launch operations. Applications identified J.M.S. from ERTS 1 investigations are summarized.

N75-13373# Earth Satellite Corp., Washington, D.C. THE ERTS COST-BENEFIT STUDY Quarterly Report, Feb. Apr. 1974

May 1974 44 p

(Contract DI-14-08-0001-13519)

(PB-234065/1; USGS-DO-74-010; QR-5) NTIS HC \$3.75 CSCL 08F

The report makes an assessment of potential benefits in marine resources management and analytical work for the case studies in land use, forestry and environmental monitoring. Interim case study reports were drafted, mission profiles for case studies to be used in developing alternative system costs and analyzing cloud cover were developed. GRA

N75-14203 *# ECON, Inc., Princeton, N.J.

THE ECONOMIC VALUE OF REMOTE SENSING OF EARTH RESOURCES FROM SPACE: AN ERTS OVERVIEW AND THE VALUE OF CONTINUITY OF SERVICE. VOLUME 1: SUMMARY

George A. Hazelrigg, Jr. and Klaus P. Heiss 20 Dec. 1974 82 p

(Contract NASw-2580)

(NASA-CR-141268; Rept-74-2002-10-1-Vol-1) Avail: NT/S HC \$4.75 CSCL 05C

An overview of the ERTS program is given to determine the magnitude of the benefits that can be reasonably expected to flow from an Earth Resources Survey (ERS) Program, and to assess the benefits foregone in the event of a one or two-year gap in ERS services. An independent evaluation of the benefits attributable to ERS-derived information in key application areas is presented. These include two case studies in agriculturedistribution, production and import /export, and one study in water management. The cost-effectiveness of satellites in an ERS system is studied by means of a land cover case study. The annual benefits achieveable from an ERS system are measured by the in-depth case studies to be in the range of \$430 to \$746 million. Benefits foregone in the event of a one-year gap in ERS service are estimated to be \$147 to \$220 million and \$274 to \$420 million for a two-year gap in ERS service. Author

N75-14204 # ECON, Inc., Princeton, N.J. THE ECONOMIC VALUE OF REMOTE SENSING BY SATELLITE: AN ERTS OVERVIEW AND THE VALUE OF

CONTINUITY OF SERVICE. VOLUME 2: SOURCE DOCUMENT

John Andrews, Alan Donziger, Jr. George A. Hazelrigg , Klaus Heiss, Francis Sand, and Peter Stevenson 20 Dec. 1974

(Contract NASw-2580)

(NASA-CR-141193; Rept-74-2002-10-2-Vol-2) Avail: NTIS HC \$7.00 CSCL 05C

The economic value of an ERS system with a technical capability similar to ERTS, allowing for increased coverage obtained through the use of multiple active satellites in orbit is presented. A detailed breakdown of the benefits achievable from an ERS system is given and a methodology for their estimation is established. The ECON case studies in agriculture, water use, and land cover are described along with the current ERTS system. The cost for a projected ERS system is given.

N75-14205 *# ECON, Inc., Princeton, N.J.

THE ECONOMIC VALUE OF REMOTE SENSING OF EARTH RESOURCES FROM SPACE: AN ERTS OVERVIEW AND THE VALUE OF CONTINUITY OF SERVICE. VOLUME 3: INTENSIVE USE OF LIVING RESOURCES: AGRICULTURE. PART 1: OVERVIEW

Alain L. Kornhauser and Lawrence B. Wilson 20 Dec. 1974 148 p

(Contract NASw-2580)

(NASA-CR-141265; Rept-74-2002-10-Vol-3-Pt-1) Avail: NTIS HC \$5.75 CSCL 05C

Potential economic benefits obtainable from a state-of-the-art ERS system in the resource area of intensive use of living resources, agriculture, are studied. A spectrum of equal capability (cost saving), increased capability, and new capability benefits are quantified. These benefits are estimated via ECON developed models of the agricultural marketplace and include benefits of improved production and distribution of agricultural crops. It is shown that increased capability benefits and new capability benefits result from a reduction of losses due to disease and insect infestation given ERS's capability to distinguish crop vigor and from the improvement in world trade negotiations given ERS's worldwide surveying capability.

N75-14206 # ECON, Inc., Princeton, N.J.

THE ECONOMIC VALUE OF REMOTE SENSING OF EARTH RESOURCES FROM SPACE: AN ERTS OVERVIEW AND THE VALUE OF CONTINUITY OF SERVICE. VOLUME 3: INTENSIVE USE OF LIVING RESOURCES, AGRICULTURE. PART 2: DISTRIBUTION EFFECTS

David F. Bradford and Harry H. Kelejian 31 Oct. 1974 115 p

(Contract NASw-2558)

(NASA-CR-141194; Rept-74-2002-10-Vol-3-Pt-2) Avail: NTIS HC \$5.25 CSCL 05C

The results of an investigation of the value of improving information for forecasting future crop harvests are described. A theoretical model is developed to calculate the value of increased speed of availability of that information. The analysis of U.S. domestic wheat consumption was implemented. New estimates of a demand function for wheat and of a cost of storage function were involved, along with a Monte Carlo simulation for the wheat spot and future markets and a model of market determinations of wheat inventories. Results are shown to depend critically on the accuracy of current and proposed measurement techniques.

Author

N75-14207 *# ECON, Inc., Princeton, N.J.

THE ECONOMIC VALUE OF REMOTE SENSING OF EARTH RESOURCES FROM SPACE: AN ERTS OVERVIEW AND THE VALUE OF CONTINUITY OF SERVICE. VOLUME 3: INTENSIVE USE OF LIVING RESOURCES, AGRICULTURE. PART 3: THE INTEGRATED IMPACT OF IMPROVED (ERS) INFORMATION ON US AGRICULTURAL COMMODITIES
Andrew D. Seidel 20 Dec. 1974 159 p refs

(Contract NASw-2580)

(NASA-CR-141262; Rept-74-2002-10-Vol-3-Pt-3) Avail: NTIS HC \$6.25 CSCL 05C

The economic value of information produced by an assumed operational version of an earth resources survey satellite of the ERTS class is assessed. The theoretical capability of an ERTS system to provide improved agricultural forecasts is analyzed and this analysis is used as a reasonable input to the econometric methods derived by ECON. An econometric investigation into the markets for agricultural commodities is summarized. An overview of the effort including the objectives, scopes, and architecture of the analysis, and the estimation strategy employed is presented. The results and conclusions focus on the economic importance of improved crop forecasts, U.S. exports, and government policy operations. Several promising avenues of further investigation are suggested.

N75-14208*# ECON, Inc., Princeton, N.J.
THE ECONOMIC VALUE OF REMOTE SENSING OF EARTH
RESOURCES FROM SPACE: AN ERTS OVERVIEW AND

RESOURCES FROM SPACE: AN ERTS OVERVIEW AND THE VALUE OF CONTINUITY OF SERVICE. VOLUME 4: FORESTRY, WILDLIFE AND RANGELAND

John Andrews and Peter Stevenson 31 Oct. 1974 136 p

refs (Contract NASw-2580)

(NASA-CR-141263; Rept-74-2002-10-Vol-4) Avail: NTIS HC \$5.75 CSCL 05C

The economic value of ERS information in the resource management area of extensive use of living resources, forestry, wildlife, and rangeland, is determined. Timber and forage resources are quantitatively evaluated. It is shown that these resources have economic value in the tens of billions of dollars, but the economic benefits of improved management of the forests and rangelands are not limited to efficiency in the production of these commercial resources. Multiple-use values including watershed, wildlife, and recreation are also involved.

N75-14209*# ECON, Inc., Princeton, N.J.
THE ECONOMIC VALUE OF REMOTE SENSING OF EARTH
RESOURCES FROM SPACE: AN ERTS OVERVIEW AND
THE VALUE OF CONTINUITY OF SERVICE. VOLUME 5:
INLAND WATER RESOURCES

Elliot Wetzler, Wes Peterson, and Martin Putnam 31 Oct. 1974 413 p

(Contract NASw-2580)

(NASA-CR-141269; Rept-74-2002-10-Vol-5) Avail: NTIS HC \$10.50 CSCL 08H

The economic value of an ERTS system in the area of inland water resources management is investigated. Benefits are attributed to new capabilities for managing inland water resources in the field of power generation, agriculture, and urban water supply. These benefits are obtained in the area of equal capability (cost savings) and increased capability (equal budget), and are estimated by applying conservative assumptions to Federal budgeting information, Congressional appropriation hearings, and ERTS technical capabilities.

N75-14210*# ECON, Inc., Princeton, N.J.

THE ECONOMIC VALUE OF REMOTE SENSING OF EARTH RESOURCES FROM SPACE: AN ERTS OVERVIEW AND THE VALUE OF CONTINUITY OF SERVICE. VOLUME 6: LAND USE. PART 1: INTRODUCTION AND OVERVIEW Keith R. Lietzke and Peter A. Stevenson 20 Dec. 1974 56 p (Contract NASW-2580)

(NASA-CR-141266) Avail: NTIS HC \$4.25 CSCL 05C

The utility of an ERS system as an effective tool in land use management is analyzed. The potential new capabilities of a space based ERS system are qualitatively examined. A variety of resource management functions are postulated within which ERTS activities might occur and the present ERS investigations in these areas are outlined.

N75-14211 # ECON, Inc., Princeton, N.J.
THE ECONOMIC VALUE OF REMOTE SENSING OF EARTH
RESOURCES FROM SPACE: AN ERTS OVERVIEW AND
THE VALUE OF CONTINUITY OF SERVICE. VOLUME 6:
LAND USE. PART 2: THE ROLE OF ERTS IN THE
ESTABLISHMENT AND UPDATING OF A NATIONWIDE
LAND COVER INFORMATION SYSTEM
Robert J. Christie 31 Oct. 1974 302 p

(Contract NASw-2558)

(NASA-CR-141267) Avail: NTIS HC \$9.25 CSCL 05C

The utility of an ERS system as an effective tool in land use management is analyzed. The cost effectiveness of satellites as a component of an ERS system is presented based on various projected levels of demand. It is indicated that a cost savings potential of \$7.9 to \$37.1 million annually is attributable to the inclusion of ERS-like satellites in the ERTS system.

N75-14212*# ECON, Inc., Princeton, N.J.

THE ECONOMIC VALUE OF REMOTE SENSING OF EARTH RESOURCES FROM SPACE: AN ERTS OVERVIEW AND THE VALUE OF CONTINUITY OF SERVICE. VOLUME 7: NONREPLENISHABLE NATURAL RESOURCES: MINERALS, FOSSIL FUELS AND GEOTHERMAL ENERGY SOURCES Keith R. Lietzke 31 Oct. 1974 85 p refs

(Contract NASw-2580)

(NASA-CR-141264; Rept-74-2002-10-Vol-7) Avail: NTIS HC \$4.75 CSCL 05C

The application of remotely-sensed information to the mineral, fossil fuel, and geothermal energy extraction industry is investigated. Public and private cost savings are documented in geologic mapping activities. Benefits and capabilities accruing to the ERS system are assessed. It is shown that remote sensing aids in resource extraction, as well as the monitoring of several dynamic phenomena, including disturbed lands, reclamation, erosion, glaciation, and volcanic and seismic activity.

N75-14213*# ECON, Inc., Princeton, N.J.
THE ECONOMIC VALUE OF REMOTE SENSING OF EARTH
RESOURCES FROM SPACE: AN ERTS OVERVIEW AND
THE VALUE OF CONTINUITY OF SERVICE. VOLUME 8:
ATMOSPHERE

Richard Miles and Gregg Fawkes 31 Oct. 1974 96 p

(Contract NASw-2580)

(NASA-CR-141191; Rept-74-2002-10-Vol-8) Avail: NTIS HC \$4.75 CSCL 05C

The economic value of an ERS system in the resource area of atmosphere is determined. Benefits which arise from air pollution and cloud observations correlated to ground stations are discussed along with cost savings associated with air pollution monitoring by satellite. Social benefits due to more precise knowledge of the effects of pollution are presented.

Author

N76-14214*# ECON, Inc., Princeton, N.J.

THE ECONOMIC VALUE OF REMOTE SENSING OF EARTH RESOURCES FROM SPACE: AN ERTS OVERVIEW AND THE VALUE OF CONTINUITY OF SERVICE. VOLUME 9: OCEANS

Keith R. Lietzke 20 Dec. 1974 134 p

(Contract NASw-2580)

(NASA-CR-141270; Rept-74-2002-10-Vol-9) Avail: NTIS HC \$5.75 CSCL 05C

The impact of remote sensing upon marine activities and oceanography is presented. The present capabilities of the current Earth Resources Technology Satellite (ERTS-1), as demonstrated by the principal investigators are discussed. Cost savings benefits are quantified in the area of nautical and hygrographic mapping and charting. Benefits are found in aiding coastal zone management and in the fields of weather (marine) prediction, fishery harvesting and management, and potential uses for ocean vegetation. Difficulties in quantification are explained, the primary factor being that remotely sensed information will be of greater benefit as input to forecasting models which have not yet been constructed.

N75-14215*# ECON, Inc., Princeton, N.J.

THE ECONOMIC VALUE OF REMOTE SENSING OF EARTH RESOURCES FROM SPACE: AN ERTS OVERVIEW AND THE VALUE OF CONTINUITY OF SERVICE. VOLUME 10: INDUSTRY

Keith R. Lietzke 31 Oct. 1974 36 p refs (Contract NASw-2580)

(NASA-CR-141192; Rept-74-2002-10-Vol-10) Avail: NTIS HC \$3.75 CSCL 05C

The economic benefits of an ERS system in the area of industrial resources are discussed. Contributions of ERTS imagery to the improvement of shipping routes, detection of previously unknown and potentially active faults in construction areas, and monitoring industrial pollution are described. Due to lack of economic research concerning the subject of ERS applications in this resource area the benefit estimations reported are regarded as tentative and preliminary.

N75-14216# European Space Research Organization, Paris (France).

EUROPEAN EARTH-RESOURCES SATELLITE EXPERI-MENTS

Bruce T. Battrick, ed. and Nguyen T. Duc, ed. May 1974 469 p refs Partly in ENGLISH and partly in FRENCH Proc. of a Symp. held at Frascati, Italy, 28 Jan. - 1 Feb. 1974 Original contains color illustrations

(ESRO-SP-100) Avail: NTIS HC \$11.50

Proceedings of the symposium are presented. The main topic was the use of imagery data from ERTS-1. Data processing methods were discussed for ERTS, EREP, and Skylab imagery. Applications of these data were considered in the fields of oceanography, hydrology, and glaciology; geology and geomorphology; and agriculture, forestry, and vegetation.

N75-14224 East Anglia Univ., Norwich (England), GENERALISATION OF ERTS DATA FOR GLOBAL-SCALE INVESTIGATIONS

I. E. Hill, A. C. Armstrong, and K. M. Clayton In ESRO European Earth-Resources Satellite Expts. May 1974 p 75-80 T

A computer-based system is described for analysis of the imagery from the ERTS project. Information contained in ERTS imagery is of use on many spatial scales. To achieve a global scale, generalization to units of the order of 50 km by 50 km is necessary. This involves problems of both data retrieval and efficient and meaningful generalization. Some results of using a computer system based on analysis of the grey-scale histograms are presented and possibilities for future developments are discussed.

Author (ESRO)

N75-14246 Centre National de la Recherche Scientifique, Toulouse (France). Service de la Carte de la Vegetation. ARNICA/ERTS-1 EXPERIMENT. RESULTS AND PERSPECTIVES [EXPERIENCE ARNICA/ERTS-1. RESULTATS ET PERSPECTIVES]

P. A. Rey In ESRO European Earth Resources Satellite Expts. May 1974 p 325-347 In FRENCH; ENGLISH summary Original contains color illustrations

The ARNICA program, concerned with processing of ERTS imagery covering southwest France and the Pyrenees, is discussed. Three fundamental fields have been treated: analog exploitation of the images, digital exploitation combined with automatic cartography, and ground truth experiments together with a study of atmospheric image transfer.

N75-14259# Army Foreign Science and Technology Center, Charlottesville, Va.

SCIENTIFIC RESEARCH AND EXPERIMENTAL DESIGN PLANNING AND COORDINATION FOR 1973

A. E. Sitalov 15 Apr. 1974 7 p Transl. into ENGLISH from Geodez. i Kartograf. (Moscow), no. 2, Feb. 1973 p 72-74 (AD-785998; FSTC-HT-23-0158-74) Avail: NTIS CSCL 08B

The report gives the results of a meeting discussing coordination of plans and projects in geodesy, aerial photography and cartography. New techniques, instruments and cooperation are introduced for 1973 to increase the efficiency of work done by the geodetic service. The topographical, geodetic and cartographic services are now part of the Scientific and Technical Mining Society.

N75-14358*# National Aeronautics and Space Administration. Lyndon B. Johnson Space Center, Houston, Tex.

THE PROCEEDINGS OF THE SKYLAB LIFE SCIENCES
SYMPOSIUM, VOLUME 1
Richard S. Johnston and Lawrence E. Diottoin New 1974

Richard S. Johnston and Lawrence F. Dietlein Nov. 1974 430 p refs Presented at Houston, Tex., 27-29 Aug. 1974 2 Vol.


(NASA-TM-X-58154-Vol-1; JSC-09275-Vol-1) Avail: NTIS HC \$11.25 CSCL 06B

The three manned Skylab missions resulted in biomedical experiment data in the areas of neurophysiology, musculoskeletal physiology, biochemistry, hematology, cytology, cardiovascular and respiratory metabolic functions: as well as detailed test objectives involving crew health and environment procedures. Major emphasis was placed on results from the last mission, Skylab 4, which covered 84 days of in-flight data collection. Many new norms were defined for normal man living and operating in a unique environment. While man is quite adaptable to this unique environment, many of the changes observed in Skylab require additional research for future flights lasting very long periods of time such as a Mars mission requiring 18 months.

Earth Resources/A Continuing Bibliography (Issue 5)

OCTOBER 1975

Typical Subject Index Listing



The subject heading is a key to the subject content of the document. The title is used to provide a description of the subject matter. When the title is insufficiently descriptive of the document content, the title extension is added, separated. from the title by three hyphens. The (NASA or AIAA) accession number and the page number are included in each entry to assist the user in locating the abstract in the abstract section (of this supplement). If applicable, a report number is also included as an aid in identifying the document. Under any one subject heading, the accession numbers are arranged in sequence with the AIAA accession numbers appearing first.

A

ABSORPTION SPECTRA

Scattering and absorption from poly-dispersed aerosols p0019 N75-11553

ABSORPTION SPECTROSCOPY

A microwave correlation spectr eter for air pollution p0063 A75-11732 monitorina Remote atmospheric sensing with an airborne laser absorption spectrometer p0066 A75-16485

ACQUISTO-OPTICS

Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation

p0063 A75-11443

AERIAL PHOTOGRAPHY A satellite-aircraft thermal study of the upwelled waters off Spanish Sahara p0041 A75-10645

Agricultural inventory techniques with orbital and p0001 A75-12872 altitude imagery

Automated remote measurement system in agricultural meteorology --- for crop survey by serial pho p0064 A75-12947

eric particles from A camera for photography of atmospl aircraft p0053 A75-13526

Terrain measurement from holographic stereograms p0001 A75-16682

Cloud screening from aerial photography applying coherent optical pattern recognition techniques p0054 A75-16690

International Symposium on Remote Sensing or Environment, 9th, University of Michigan, Ann Arbor, Mich., April 15-19, 1974, Proceedings. Volumes 1, 2 & 3 p0068 A75-17101-Some features of the urban environment of Tokyo by

sensing --- infrared surface temperature p0012 A75-17102 A method of specifying remotely

p0002 A75-17106 sample points data from multiplexed p0054 A75-17117 Extraction of urban land cover synthetic aperture radar imagery

Investigation of radar discrimination of sea ice p0042 A75-17121.

Areal extent of snow in forested regions - A practical estimation technique using ERTS-1 data

p0002 A75-17122 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado p0013 A75-17137

Flood inundation in the southeastern United States fro p0047 A75-17138 aircraft and satellite imagery The use of remote sensing and natural indicators to

use of remote sensing una state floodplains - Preliminary findings p0067 A75-17142

Evaluation of ERTS-1 and aircraft data for assessing internal drainage in irrigated agriculture

p0002 A75-17143 rmal infrared remote p0002 A75-17144 ng to study soil moisture

Modifications in a computer-implem sted method for the detection and extraction of objects in serial photographs

angle on automatic p0014 A75-17157 classification of ERTS-1 data Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites

p0048 A75-17160 Scanning thermal plumes ... from power plant p0014 A75-17162

The use of remote sensing in lim utrophication of lakes 50048 A75-17163 Preliminary results of fisheries inv estigation associated ith Skylab-3 p0003 A75-17167 Boundaries of ERTS and aircraft data within which useful

water quality information can be obtained p0014 A75-17182

Texture analysis with Fourier series --- for crop identification from color serial photographs

p0004 A75-17187 Land use classification and around truth correlations from simultaneously acquired aircraft and ERTS-1 MSS date

p0014 A75-17188 Photographs from balloons - Their use in agrono anagement of environment p0055 A75-17191
A motion-compensated spatial scanner --- MSS design management of environment

p0068 A75-17213 Automatic contouring on the Gestalt Photomapper 2-1 p0028 A75-18055

An intensive natural resources in tory aids land use planning and resources management in Puerto Rico

p0005 A75-18056 The use of high altitude photography and ERTS-1 imagery for wildlife habitat inventory in Kansas

p0005 A75-18066 Early results from the experimental Automated Image

Data Extraction System --- for serial terrain sensing p0069 A75-18072 The PUCE programme for terrain evaluation for igineering purposes. 2: Procedures for terrain

engineering purposes. classificati [PB-233847/3] p0029 N75-10625

47/3] a panoramic serial camera in serosurveying p0071 N75-11450 [AD-784162] Correlation of missions 191, 51M and helicopter photography --- aerial photography and mapping of three eas in Test Site 175 in Texas

[NASA-CR-134289] Image selection system' --- computerized data storage

[NASA-CR-2475] p0058 N75-12422 From aerospace photographs to forecasting and calculating flows

[NASA-TT-F-16006] p0059 N75-13387

A comparison of ERTS imagery with conventional aerial photography for land-resource surveys in less developed countries. Examples from the Rift Valley Lakes Basin. p0061 N75-14249

Investigation of accuracy of determination of plane coordinates by means of the Central Scientific Research Institute of Geodesy, aerial photography and cartography rangefinder

[AD-785886] p0061 N75-14255 Morphographic and morphometric analysis of vegetative cover photographs in serial photography for the mapping tatio

[AD-785932] n0009 N75-14258 rch and experimental design planning and coordination for 1973

[AD-785998] The availability of local serial photography in southern California --- for solution of urban planning problems
[NASA-CR-141314] p0025 N75-1

n0025 N75-15126 Use of forest vegetation as an indicator in the interpretation and large scale mapping of soils in a forest

[AD-785967] p0010 N75-15146

AERIAL RECONNAISSANCE

Special color analysis techniques p0064 A75-12874 sensor imagery of high-power pulses of current

soidal shape for an serial variant of the method transient processes --- for serial prosp p0053 A75-13310

Pattern recognition relevant to reconnaissance and remote sensing [PHL-1974-23] ··· noting classification methods p0070 N75-10573

panoramic aerial camera p0071 N75-11450 [AD-784162]

Correlation of missions 191, 51M and helicopter photography --- aerial photography and mapping of three areas in Test Site 175 in Texas [NASA-CR-134289] p0058 N75-12279

Contribution of space platforms to a ground and airborne remote-sensing programme over active Italian volcano p0073 N75-14236

Ground-truth measurements in relation to aircreft and satellite studies of agricultural land use and land classification in Britain --- ERTS-1 and serial remote sensing p0009 N75-14251

Agricultural resources investigations in northern Italy, southern France and Madagascar --- ERTS-B and aerial rice growing and forestry surveys p0009 N75-14252

Determination of the serosol content in the atmospher from ERTS-1 data p0013 A75-17130

The point visibility meter - A forward scatter instrument for the measurement of serosol extinction coef

p0070 A75-19266 Quantitative determination of stratospheric aerosol

p0016 N75-10535 Scattering and absorption from poly-

p0019 N75-11553

Multiple scattering in cloud layers; some results --emphasizing aerosol parameters on global basis p0019 N75-11554

Lidar observations of atmospheric particulate content

p0019 N75-11559 Quantitative determination of stratospheric aerosol

[E75-10010] p0020 N75-12393 Quantitative determination of stratospheric aerosol

[E75-10079] p0023 N75-14198 Determination of the earth's aerosol albedo using Skylab

data --- Lake Michigan [E75-10096] p0025 N75-15114

AEROSPACE MEDICINE

The Proceedings of the Skylab Life Sciences Symposium, [NASA-TM-X-58154-VOL-1] p0081 N75-14358

AEROSPACE SYSTEMS

The potential benefits for development derived from space applications
[IAF PAPER 74-147] p0075 A75-13761

AFRICA
The detection of dust storms over land and water with satellite visible and infrared measurement: o0012 A75-16675

An examination of the extent of fire in the grassland and savanna of Africa along the southern side of the share p0002 A75-17104
Comparison of the Southern and Northern Hemisphere

general circulation characteristics as determined by satellite

[UCRL-75556] p0017 N75-10638 Cruise data report. CHAIN 115 leg 2, Southlant expedition --- to survey geophysical parameters

[AD-784421] n0029 N75-10641 Mapping of the major structures of the African rift system Somalia, and Yemen Ethioo

p0030 N75-13340 [E75-10024] Utilization of ERTS-1 imagery in cultivation and settlement site identification and carrying capacity estimates in Upper Volta and Niger

[E75-10029] p0008 N75-13345 Preliminary evaluation of ERTS-1 imagery of Central p0006 N75-14249

A comparison of ERTS imagery with conventional serial photography for land-resource surveys in less developed countries. Examples from the Rift Valley Lakes Basin, Ethiopia p0061 N75-14249

AFRICAN RIFT SYSTEM

ERTS-1 data on Afar tectonics p0031 N75-14239 A comparison of ERTS imagery with conventional serial photography for land-resource si rveys in less developed lift Valley Lakes 8 asin, p0061 N75-14249 Examples from the Rift

Ethiopia

AGRICULTURE

GRICULTURE
Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872
Automated remote measurement system in agricultural meteorology --- for crop survey by serial photo-

p0064 A75-12947 Agricultural resources investigations in Southern France d Northern Italy - p0004 A75-17173 and Northern Italy Statistical separability of agricultural cover types in

subsets of one to twelve spectral channels p0005 A75-17216

Geographic applications of ERTS-1 imagery to landscape change --- Mississippi River and Great Smoky Mountains of Tennessee and North Carolina

[E74-10810] p0034 N75-10537 Agriculture, forestry, range resources

n0006 N75-10558 Feasibility study ASCS remote sensing/compliance

determination system [NASA-CR-134288] p0007 N75-12419 The value of improved (ERS) information based on

domestic distribution effects of U.S. agriculture crops [NASA-CR-141046] p0007 N75-12423 remote measuring Automated

ing system in p0007 N75-12518 prometeorology p0007 N75-12518
Utilization of ERTS-1 imagery in cultivation and settlement site identification and carrying capacity estimates in Upper Volta and Niger [E75-10029] p0008 N75-13345

The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources: Agriculture. Part 1: Overview [NASA-CR-141265] p0079 N75-14205

The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources, agriculture. Part 2: Distribution effects

[NASA-CR-141194] p0079 N75-14206

[NASA-LH-141194] D0079 N75-14206 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources, agriculture. Part 3: The integrated impact of improved (ERS) information on US agricultural commodities.

[NASA-CR-141262] p0079 N75-14207 Agricultural resources investigations in northern Italy, southern France and Madagascar --- ERTS-B and aerial rice growing and forestry surveys p0009 N75-14252
Future US activities based on experience gained from
ERTS-1 and EREP --- land use, forestry, agriculture, and p0009 N75-14253
Irrigation scheduling, freeze warning and soil salinity detecting - Cameron and Starr Counties, Texas
[E75-10109]

AIR POLLUTION

The 1973 program of measurement of the minor constituents of the stratosphere using the Concorde 001

p0063 A75-11720
A microwave correlation spectrometer for air pollution onitoring eter for air pollution p0063 A75-11732 monitoring On evaluating compliance with air pollution levels 'not to be exceeded more than once a year

p0011 A75-12803

Trace gas analysis of power plant plumes via aircraft measurement - 03, NOx, and SO2 chemistry p0064 A75-13148

Airborne survey of major air basins in California p0011 A75-14360

On the detectability of atmospheric carbon monoxide by microwave remote sensing p0013 A75-17129 Determination of the aerosol content in the atmosph from ERTS-1 data p0013 A75-17130

from power plant p0014 A75-17162 Scanning thermal plumes --condensers Conference on Environmental Sensors and Applications.

London, England. November 18, 19, 1974, Proceedings p0070 A75-19265 Feasibility study of a Gunn-diode microwave cavity spectrometer as a formaldehyde monitor

p0071 N75-11311 [UCID-16488] Observation and measurement of atmospheric pollution

p0018 N75-11476 [WMO-368] A technique for measurement of sulphur dioxide and sulphate concentrations from an aircraft

p0018 N75-11498 Aircraft measurements of the concentration and size distribution of atmospheric sulfate particles

n0057 N75-11504 Correlation spectroscopy techniques for the regional and global measurements of air pollution p0018 N75-11505 Microwave rotational spectroscopy: A physical technique specific pollutant monitoring p0018 N75-11529

for specific pollutant monitoring Measurements of meteorological parameters and air pollution spread over Stockholm, Sweden, by means of remote sensing and airborne instrumentation

p0018 N75-11531

Airborne vapor surveillance system p0021 N75-12459 [AD-785585]

AIR SAMPLING

Observation and measurement of atmospheric pollution [WMQ-368] p0018 N75-11476

Methods for the automatic recording of atmospheric dust concentrations and some of the results obtained p0057 N75-11502

AIR WATER INTERACTIONS

Monitoring and predicting El Nino invasions --- equatorial Pacific Ocean-atmosphere interactions

n0041 A75-12719 The wind vector in a layer over water p0041 A75-15179

AIRBORNE EQUIPMENT

Performance, comparison of technic p0063 A75-10075 stereo radar images

The 1973 program of measurement of the minor constituents of the stratosphere using the Concorde 001 p0063 A75-11720

Mathematical modeling for the detection of fish by an aborne laser p0065 A75-15953 Remote atmospheric sensing with an airborne lase

p0066 A75-16485 absorption spectrometer Passive microwave sensing of moist soils

p0066 A75-17124

Rapid stock assessment of pilchard populations by craft-borne remote sensors p0003 A75-17168 aircraft-borne remote sensors A technique for measurement of sulphur dioxide and sulphate concentrations from an aircraft

p0018 N75-11498 Aircraft measurements of the concentration and size

distribution of atmospheric sulfate particles p0057 N75-11504 Airborne vapor surveillance system

[AD-785585] p0021 N75-12459

ALABAMA

The remote-sensing program of the Geological p0069 A75-17230 of Alabama ERTS-1 imagery interpretation techniques in the

Tennessee Valley --- land use and soil mapp [E74-10811] p0016 N75-10538

A review of initial investigations to utilize ERTS-1 data in determining the availability and distribution of living marine resources --- harvest and management of fisheries resources in Mississippi Sound and Gulf weter p0043 N75-10555

MSFC integrated experiments preliminary report --- for the Skylab program [NASA-TM-X-64881]

p0078 N75-10967 Determination of points of entry for potential contaminants into limestone aguifers using thermal infrared

imagery [NASA-CR-120540] p0021 N75-12546 Computer and photogra of central north Alabama [NASA-TR-R-431] otogrammetric general land use study

p0022 N75-13372

A summary of ERTS data applications in Alaska p0077 A75-17236

The application of ERTS-1 data to the Corps of Engineers Civil Works Mission --- for ecological monitoring p0015 A75-18064

Evaluate ERTS imagery for mapping and detection of changes of snowcover on land and on glaciers --- Alaska and Washington p0049 N75-10539 [E74-10812]

Environmental surveys in Alaska based upon ERTS sta p0017 N75-10551 Mineral resources, geological structures, and landform p0035 N75-10561

ALGORITHMS

clustering algorithm for classification p0002 A75-17113 Remote sensing geophysics from Skylab --- rationing-type

algorithms to enhance multispectral data p0030 N75-13352

ALPS MOUNTAINS (EUROPE)

Methodological aspects and regional examples of snow-cover mapping from ERTS-1 and EREP imagery of the Swiss Alps p0051 N75-14232 Geological features on ERTS-1 images of

nOO37 N75-14237 the west-central Alps --- Italy ALTIMETERS

Terrain properties and topography from Skylab altimetry

[£75-10059] p0030 N75-13359 Calibration and evaluation of Skylab altimetry for geodetic determination of the geoid p0030 N75-13364

AMPHIBIOUS VEHICLES

The use of color infrared imagery for the study of ma buggy tracks p0061 N75-15127 ANDESITE

Remote sensing geophysics from Skylab reflectance anomalies of andesites and basalts in Southern

California and Nevada p0036 N75-13351 [E75-10051]

ANOMALOUS TEMPERATURE ZONES Geologic interpretation of infrared thermal images in East Oatrani area, Western Desert, Egypt ,p0033 A75-17215

ANTARCTIC REGIONS Microbial abundance and thermol antarctic dry valley soils

p0005 A75-17973 Overall evaluation of ERTS-1 imagery for cartographic application --- Australia, Antarctica, U.S., and U.S.S.R. [F74-10800] p0029 N75-10528

Earth Resources Technology Satellite-1 (ERTS-1) data and anthropology: Use of these data in carrying capacity estimates for sites in Upper Volta and Niger DOO78 N75-11406 [E75-10001]

APOLLO TELESCOPE MOUNT

Apollo Telescope Mount experiments technology p0064 A75-13589

AOUIFFRS

Determination of points of entry for potential contaminants into limestone aquifers using thermal infrared [NASA-CR-120540]

ARCTIC OCEAN

A study of microwave emission properties of sea ice -AIDJEX 1972 p0041 A75-17120 Investigation of radar discrimination of sea ice p0042 A75-17121

ARCTIC REGIONS

A satellite view of weather systems over the North
American Arctic p0063 A75-10790
Use of ERTS-1 in sea-ice studies p0045 N75-14234 AIDJEX bulletin number 25 --- study of pack ice, sea ice, and Beaufort Sea ice

[PB-235298/7] n0045 N75-14262

ARGENTINA

The interpretation and use of false-colour infra-red and true colour photography of part of Argentina obtained by Skylark earth resources rockets p0067 A75-17192 A comparison of the capabilities of ERTS-1 and the

A comparison of the capeumuses
Skylark earth-resources rocket for resources surveying in p0023 N75-14250 ARID LANDS Operational use of satellite and high altitude remote

sensing for the generation of input data for water demand p0067 A75-17148 A study to explore the use of orbital remote sensing to determine native and plant distribution --- Arizona Re Ecological Test Site [E74-10818] p0006 N75-

p0006 N75-10544 Dust storms according to data of space research spaceborne photography of and zones [NASA-TT-F-18021] p0021 N75-1244

p0021 N75-12463

Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p0013 A75-17136

Remote detection of soil surface moisture p0003 A75-17146
A study to explore the use of orbital remote sensing to

determine native ario Ecological Test Site e arid plant distribution --- Arizona Regional p0006 N75-10544 [E74-10818]

Land use and mapping p0017 N75-10560 In situ spectroradiometric calibration of EREP imagery and oceanography of Block Island Sound ··· Arizona

geology [E75-10042] p0072 N75-12411 Application of Skylab EREP photographs to study of the modern episode of accelerated erosion in southern

Arizona [E75-10054] p0036 N75-13354 Preliminary results in the comparison of Skylab, ERTS

and RB-57 images for the detection of lineaments and fractures in Precambrian, Paleozoic, and late Tertiary rocks on and near the Colorado Plateau, North-Central Arizona 75-10060] p0037 N75-13360 Cartographic evaluation of Skylab-A S-192 scanner

images --- Chesapeake Bay, Phoenix, and San Francisco

(E75-10078) p0031 N75-14197 Urban and regional land use analysis: CARETS and census cities experiment package --- New Haven, Connecticut; Pontiac, Michigan; Phoenix, Arizona; District of Columbia

[E75-10091] p0024 N75-15109 ARIZONA REGIONAL ECOLOGICAL TEST SITE

A study to explore the use of orbital remote sensing to determine native arid plant distribution --- Arizona Regional Ecological Test Site [E74-10818] p0006 N75-10544

ARKANSAS

Detection of moisture and moisture related phenomena from Skylab --- Kansas, Oklahoma, and Arkan [E75-10035] p0007 N75-12405

ASTRONAUT PERFORMANCE

The Proceedings of the Skylab Life Sciences Symposium,

[NASA-TM-X-58154-VOL-1] p0081 N75-14358 ATCHAFALAYA RIVER BASIN (LA)

Evaluation of satellite remote sensing and automatic data techniques for characterization of wetlands and coastal marshlands --- detection and measurement of accretion in estuary system of Atchafalaya River in southern

[E75-10025] p0044 N75-13341 Evaluation of satellite remote sensing and automatic data

techniques for characterization of wetlands and coastal marshlands --- Atchafalaya River Basin, Louisiana p0050 N75-13344 [E75-10028]

ATLANTIC OCEAN

The detection by ERTS-1 of wind-induced ocean surface features in the lee of the Antilles Islands

p0041 A75-10644

A satellite-aircraft thermal study of the upwelled waters f Spanish Sahara p0041 A75-10645 off Spanish Sahara Rapid stock assessment of pilchard populations by

hard populations by p0003 A75-17168 aircraft-borne remote sensors Ocean internal waves off the North American and African p0042 A75-17183 coasts from ERTS-1

SUBJECT INDEX **CALIFORNIA** Comparison of the Southern and Northern Hemisphere Atmospheric release advisory capability: Research and BLACK HILLS (SD-WY) Experiment to evaluate feasibility of utilizing Skylab-EREP general circulation characteristics as determined by satellite progress p0025 N75-15202 remote sensing data of tectonic analysis of the Bighorn Mountains region, Wyoming-Montana --- Black Hills, South JUCRL-75644-REV-21 [UCRL-75556] p0017 N75-10638 ATMOSPHERIC SCATTERING Dakota/Wyoming Cruise data report. CHAIN 115 leg 2, Southlant Ground-based determination of atmospheric radiance for correction of ERTS-1 data
ATMOSPHERIC TEMPERATURE p0054 A75-16476 --- to survey geophysical parameters p0029 N75-10641 Inventory of forest and rangeland resources, including forest stress --- Colorado, Georgia, and Black Hills, South [AD-784421] Nimbus 4 observations of cha An assessment of the potential contributions to oceanography from Skylab visual observations and hand-held photography --- Pacific Ocean, Atlantic Ocean es in total ozone and stratospheric temperatures during a sud Dakota dden warming p0012 A75-14841 [E75-10032] p0007 N75-12402 Theoretical investigations on the determination of the ulf of Mexico BLIGHT vertical temperature distribution in the atmosphere Spectral reflectance studies on mineral deficiency in co [E75-10017] assuming multispectral thermal radiation measurements Interpretation of ERTS-1 data for investigating the French plants n0003 A75-17172 60 GHz [BMFT-FB-W-74-04] Thermal behaviour of some rice fields affected by a western coastal zones (Fralit program) p0024 N75-14331 p0051 N75-14230 vellows-type disease p0004 A75-17174 MOSPHERIC WINDOWS

Estimation of sea surface temperature from remote ATMOSPHERIC ATTENUATION BLOCK ISLAND SOUND (RI) Deteriorating effects on 3 mm wave pa e passive imagery p0055 A75-17198 In situ spectroradiometric calibration of EREP imagery sensing in the 11to 13-micron window re p0065 A75-14803 and oceanography of Block Island Sound --- Arizona Measurements of the extraterrestrial solar radiant flux from 2981 to 4000 A and its transmission through the geology [E75-10042] ATOMIC SPECTRA p0072 N75-12411 Life cycle cost study of Army Spectrometric Oil Analysis Program (ASOAP) from 2981 to 4000 A and its definition through the earth's atmosphere as it is affected by dust and ozone p0070 A75-19547 BONNEVILLE SALT FLATS (UT) [AD-786501] p0073 N75-14870 Experimental evaluation of atmospheric effects on Study of atmospheric effects in Skylab data 75-10006l p0020 N75-12391 radiometric measurements using the EREP of Skylab Salt Flats, Utah [F75-10006] p0053 A75-11822 The community satellite, III AUSTRALIA [E75-10036] p0020 N75-12406 ATMOSPHERIC CHEMISTRY Overall evaluation of ERTS-1 imagery for cartographic BOUNDARY LAYERS Trace gas analysis of power plant plumes via aircraft measurement - 03, NOx, and SO2 chemistry p0064 A75-13148 annlication --- Australia, Antarctica, U.S. Remote sensing of ocean current boundary layer ---[E74-10800] p0029 N75-10528 The PUCE programme for terrain evaluation for ngineering purposes. 2: Procedures for terrain [E74-10801] n0042 N75-10529 Quality control in global and regional air chemistry engineering purposes. An assessment of the potential contributions to oceanography from Skylab visual observations and hand-held photography --- Pacific Ocean, Atlantic Ocean monitoring networks: Lessons of the European Air Chemistry Network p0018 N75-11540 p0018 N75-11540 [PB-233847/3] ATMOSPHERIC CIRCULATION o0029 N75-10625 A study of the usefulness of Skylab EREP data for earth Monitoring and predicting El Nino invasions --- equatorial Pacific Ocean-atmosphere interactions and Gulf of Mexico cources studies in Australia --- Australia geology 75-10047] p0036 N75-12415 [E75-10017] p0044 N75-12396 [E75-10047] n0041 A75-12719 Remote sensing of ocean current boundary layer --- Gulf Australian and Soviet gravity surveys along the Australian Comparison of the Southern and Northern Hemi general circulation characteristics as determined by satellite [E75-10031] p0045 N75-14185 Remote sensing of ocean current boundary layer --- Gulf of Mexico Recognition and interpretation of spectral signatures of egetation from aircraft and satellite imagery in Western tueensland, Australia p0008 N75-14241 [UCRL-75556] DO017 N75-10638 ATMOSPHERIC COMPOSITION [E75-10067] p0045 N75-14186 Queensland, Australia The 1973 program of measurement of the minor Remote sensing of ocean current boundary layer --- Gulf constituents of the stratosphere using the Concorde 001 of Mexico p0063 A75-11720 В [E75-10098] p0046 N75-15116 A camera for photography of atmospheric particles fro BRIGHTNESS TEMPERATURE p0053 A75-13526 A radio picture of the earth --- microwave remote Spectrophotometric observations BACKSCATTERING earth from satellites p0066 A75-16191 Sovuz-13 Active microwave measurement of soil water content [IAF PAPER 74-156] Signatures of various earth surfaces measured by the p0065 A75-13765 p0001 A75-16242 [NASA-CR-141685] Nimbus-5 microwave spectrometer p0066 A75-17119 Airborne survey of major air basins in Californi Simultaneous active and passive microwave response A study of microwave emission properties of sea ice -AIDJEX 1972 p0041 A75-17120 n0011 A75-14360 of the earth - The Skylab radscat experiment Remote atmospheric sensing with an airborne p0012 A75-17114 Design data collection with Skylab/EREP microwave instrument S-193 --- Texas, Utah, Illinois, and Minnesota [E75-10080] p0072 N75-14199 p0066 A75-16485 absorption spectrometer Radar response to vegetation --- soil moisture mapping On the detectability of atmospheric carbon m via microwave backscattering p0005 A75-18556
A study of a dual polarization laser backscatter system p0013 A75-17129 [E75-10080] microwave remote sensing BUOYS The potentialities for determining content of minor gas for remote identification and measurement of water Report of the meeting on drifting buoys for the first GARP constituents in the atmosphere by satellites p0015 A75-17932 global experiment [GARP-SPEC-REPT-13] [NASA-CR-140665] p0017 N75-10563 p0043 N75-10674 Methods for the automatic recording of atmospheric dust BALLOON-BORNE INSTRUMENTS
Detection of synchronous changes in fluxes of energetic electrons and gamma-quanta during a magnetic concentrations and some of the results obtained p0057 N75-11502 during a magnetic p0064 A75-12317 C Aircraft measurements of the concentration and size BASALT distribution of atmospheric sulfate particles p0057 N75-11504
Geophysical monitoring for climatic change. The NOAA
rogram p0018 N75-11510 Remote sensing geophysics from Skylab --- near infrared reflectance anomalies of andesites and basalts in Southern CALIBRATING Transference of ERTS-1 spectral signatures in time and program p0018 N75-11510
Measurement of trace stratospheric gases by inverting California and Nevada n0066 A75-17111 Boundaries of ERTS and aircraft data within which useful p0036 N75-13351 BEAUFORT SEA (NORTH AMERICA) thermal infrared limb radiance profiles water quality information can be obtained p0014 A75-17182 AIDJEX bulletin number 25 --- study of pack ice, seace, and Beaufort Sea ice p0018 N75-11523 ATMOSPHERIC CONDUCTIVITY Australian and Soviet gravity surveys along the Australian Ice particulates in the n
ATMOSPHERIC EFFECTS mesosphere p0012 A75-16752 [PB-235298/7] p0045 N75-14262 p0031 N75-13454 Design data collection with Skylab/EREP microwave instrument S-193 --- Utah and Texas Rad/Scat BESMEX: Bering Sea marine mammal experiment ---CALIFORNIA vith the primary target species being the walrus and A simulation of the San Andreas fault experiment p0033 A75-11068 measurements bowhead whale A comparison of Skylab and ERTS data for agricultural [E75-10070] n0059 N75-14189 [NASA-TM-X-62399] p0008 N75-13503 Design data collection with Skylab/EREP microwave BIGHORN MOUNTAINS (MT-WY) crop and natural vegetation interpretation AIAA PAPER 74-1219] p0001 A75-12033 Airborne survey of major air basins in California Experiment to evaluate feasibility of utilizing Skylab-EREP emote sensing data of tectonic analysis of the Bighorn p0072 N75-14199 [E75-10080] p0011 A75-14360 ATMOSPHERIC HEATING Mountains region, Wyoming-Montana --- Black Hills, South Nimbus 4 observations of changes in total ozone and Dakota/Wyoming Remote detection of soil surface moisture p0003 A75-17146 stratospheric temperatures during a sudo p0030 N75-12395 DO012 A75-14841 Operational use of satellite and high altitude remote BIOGEOCHEMISTRY ATMOSPHERIC MOISTURE Remote sensing to detect the toxic effects of metals on vegetation for mineral exploration p0034 A75-17226 sensing for the generation of input data for water demand models p0067 A75-17148 The effect of atmospheric water vapor on automatic classification of ERTS data p0013 A75-17131 p0013 A75-17131 Determining range condition and forage production potential in California from ERTS-1 imagery BIOMEDICAL DATA ATMOSPHERIC OPTICS The Proceedings of the Skylab Life Sciences Symposium. agery p0003 A75-17169 On the natural limitations of target differentiation by means of spectral discrimination techniques --- ERTS remote sensors p0054 A75-17132 ERTS 1 --- coastal p0048 A75-17179 [NASA-TM-X-58154-VOL-1] California nearshore processes p0081 N75-14358 BIRDS currents and sediments The point visibility meter - A forward scatter instru Surface currents along the California Utilization of Skylab (EREP) system for appraising for the measurement of aerosol extinction coef ERTS imagery p0048 A75-1718U
The usefulness of imaging passive microwave for rural changes in continental migratory bird habitat [E74-10817] p0006 o0070 A75-19266 p0006 N75-10543 ATMOSPHERIC PRESSURE Utility of ERTS for monitoring the breeding habit of migratory waterfowl --- United States and Canada and urban terrain analysis p0055 A75-17199 Report of the meeting on drifting buoys for the first GARP Oil slick detection by X-band synthetic aperture radar global experiment [GARP-SPEC-REPT-13] DO068 A75-17209 o0006 N75-10557 p0043 N75-10674 Unsupervised mapping of geologic features and soils in alifornia p0034 A75-17225 ATMOSPHERIC RADIATION
Proceedings of the UCLA International Conference on
Radiation and Remote Probing of the Atmosphere ... Utilization of Skylab (EREP) system for appraising changes in continental migratory bird habitat Evaluate ERTS imagery for mapping and detection of p0008 N75-13361 changes of snowcover on land and on glaciers --- Alaska Utilization of Skylab (EREP) system for appraising emphasizing radiative transfer [NASA-CR-140816] changes in continental migratory bird habitat [E75-10087] p0009 p0078 N75-11549 [E74-10812]

n0017 N75-10560

p0035 N75-11407

Land use and mapping

Southern California

[E75-10003]

Lineaments in basement terrane of the Peninsular Ranges,

p0009 N75-15105

p0043 N75-10673

BISTATIC REFLECTIVITY

[NASA-CR-137469]

Bistatic radar sea state monitoring

Influence of the atmosphere on spectral radiance and contrasts of natural formations measured from space ---

p0019 N75-11557

using data from Soyuz and Salvut spacecraft

Fault tectonics and earthquake hazards in the Peninsular Applications of satellite geodesy in the exploration of CLINICAL MEDICINE Southern California the internal structure of the earth p0029 A75-18123 ental and crew microbiology studies p0024 N75-14366 Ranges, South [£75-10005] Skylab enviro The gravity field and the definition of stationary sea Remote sensing geophysics from Skylab --- Southern p0042 A75-18124 CLOUD COVER surface topography California and Nevada minerals and rocks Possibilities of calculating the earth's local gravitational ald from satellite observations p0029 A75-18524 p0030 N75-12413 satellite visible and infrared measurements [E75-10045] field from satellite observations Fault tectonics and earthquake hazards in the Peninsular DO012 A75-16675 Cloud screening from aerial photography applying Ranges, Southern California Urban and regional land use analysis: CARETS and [E75-10022] p0030 N75-13338 coherent optical pattern recognition technical census cities experiment package [E75-10056] Analysis of pseudocolor transformations of ERTS-1 images of Southern California area --- geological faults and p0054 A75-16690 p0022 N75-13356 The calculation of cloud shadows in modelling of the arth's surface survey from space p0014 A75-17204 Urban and regional land use analysis: CARETS and earth's surface survey from space p0014 A75-17204
Observing and forecasting local effects from satellite data census cities experiment package p0036 N75-13346 [E75-10100] n0025 N75-15118 p0016 A75-18153 Remote sensing geophysics from Skylab --- near infrared --- mesoscale cloud patterns reflectance anomalies of andesites and basalts in Southern CENTRAL AFRICAN REPUBLIC An examination of the extent of fire in the grassland and savanna of Africa along the southern side of the INASA-TT-F-7981 California and Nevada p0057 N75-11419 p0036 N75-13351 Geographical correlation of TV and IR images obtained [E75-10051] Sahara n0002 A75-17104 Fault tectonics a nd earthquake hazards in the Peninsular from weather satellites --- cloud cover photography CENTRAL ATLANTIC REGIONAL ECOL TEST SITE Ranges, Southern California --- Gulf of California p0071 N75-11433 Urban and regional land use analysis: System for electronic transformation and geographic -100741 p0031 N75-14193 CARETS and Cartographic evaluation of Skylab-A S-192 scanner census cities experiment package correlation of satellite television information --- cloud cover p0071 N75-11434 --- Chesapeake Bay, Phoenix, and San Francisco [E75-10019] n0020 N75-12398 Geographical correlation of earth cloud cover pictures test sites Urban and regional land use analysis: CARETS and DO071 N75-11435 [E75-10078] n0031 N75-14197 obtained from unoriented satellites nsus cities experiment package [E75-10056] p0022 N75-13356 Multiple scattering in cloud layers; some results ---Study to develop improve d spacecraft snow survey Urban and regional land use analysis: CARETS and ensus cities experiment package --- New Haven, methods using Skylab/EREP data --- Sierra Nevada,
Cascades, and Great Plains
[E75-10081] p0073 N75-14200 emphasizing aerosol parameters on global bas p0019 N75-11554 census cities experiment package --- New Haven. Connecticut; Pontiac, Michigan; Phoenix, Arizona; District Environmental satellite imagery, September 1974 Generalisation of ERTS data p0058 N75-11571 for global-scale of Columbia investigations --- Central Valley, Californi CLOUD HEIGHT INDICATORS p0024 N75-15109 [E75-10091] p0081 N75-14224 The role of lidar in environmental sensing --- laser Urban and regional land use analysis: CARETS and The remote identification of terrain features and materials ceilometer and visual range measurement appli census cities experiment package [E75-10100] at a California test site. An investigative study of p0070 A75-19274 p0025 N75-15118 CLOUD PHOTOGRAPHS CENTRAL EUROPE [PB-235991/7] p0031 N75-14261 Cloud screening from aerial photography applying Use of terrestrial gravity data in combination with satellite results for determining geoid heights in Central Europe Fault tectonics and earthquake hazards in the Peninsular coherent optical pattern recognition technique Ranges, Southern California [E75-10108] p0054 A75-16690 p0029 A75-18128 n0039 N75-15124 CLOUD PHOTOGRAPHY CHEMICAL PROPERTIES The availability of local aerial photo graphy in southern A camera for photography of atmospheric particles from Monitoring physical and chemical parameters of Delaware Bay waters with an ERTS-1 data collection platform [E75-10090] p0051 N75-15108 California ---, for solution of urban planning prob [NASA-CR-141314] p0025 N p0053 A75-13526 DO025 N75-15126 System for electronic transformation and geographic Geologic structure in California: [E75-10090] correlation of satellite television information ERTS-1 imagery CHESAPEAKE BAY (US) p0071 N75-11434 photography The application of ERTS-1 data to the Corps of Engineers [NASA-TM-X-70799] n0039 N75-15132 graphical correlation of earth cloud cover pictures Civil Works Mission --- for ecological mon p0071 N75-11435 Usefulness of remote sensing techniques for the environmental monitoring of the Sacramento-San Joaquin obtained from unoriented satellites monitoring p0015 A75-18064 A cloud physics investigation utilizing Skylab data p0025 N75-15135 [E75-10021] p0020 N75-12400 Skylab-EREP investigations of wetlands ecology --- New Meteorological data. Volume 139, no. 4: The European climatology, 1973. Meteorological satellite-data from the Berlin zone of reception, part 4, fourth quarter CAMERAS Jersey and Chesapeake Bay A camera for photography of atmospheric particles from [E74-10816] p0049 N75-10542 p0053 A75-13526 A survey for the use of remote sensing in the Chesapeake p0023 N75-13483 The Skylab S-190B Earth Terrain Can Bay regin p0069 A75-18071 CLOUD PHYSICS [NASA-CR-137476] p0043 N75-10571 A cloud physics investigation utilizing Skylab data [E75-10021] n0020 N75-1 CANADA Cartographic evaluation of Skylab-A S-192 scar n0077 N75-10550 The Canadian ERTS program p0020 N75-12400 images --- Chesapeake Bay, Phoenix, and San Francisco A cloud physics investigation utilizing Skylab data [E75-10099] p0025 N75-1 Utility of ERTS for monitoring the breeding habit of p0025 N75-15117 migratory waterfowl --- United States and Canada [E75-10078] p0031 N75-14197 p0006 N75-10557 COACHELLA VALLEY (CA) CHRONOLOGY Fault tectonics and earthquake hazards in the Peninsular Ranges, Southern California Water survey of Canada: Application for use of ERTS-A Remote sensing geophysics from Skylab --- stratigraphic analysis of weathered rocks for retransmission of water resources data p0049 N75-11409 [E74-10805] [E75-10022] p0030 N75-13338 p0037 N75-13358 [E75-10058] Water survey of Canada: Application for use of ERTS-A COAL CITIES retransmission of water resources data Classification and mapping of coal refuse, vegetative Urban and regional land use analysis: CARETS and [E75-10026] p0050 N75-13342 cover types, and forest types by digital processing ERTS-1 ties experiment package Water survey of Canada: Application for use of ERTS-A p0020 N75-12398 p0054 A75-17110 (E75-10019) nsmission of water resources data Urban and regional land use analysis: CARETS and Automated strip mine and reclamation mapping from [E75-10053] p0050 N75-13353 ERTS --- Ohio coal p0035 N75-10556 ensus cities experiment package CAPE KENNEDY LAUNCH COMPLEX [E75-10056] p0022 N75-13356 A demonstration of ERTS-1 analog and digital techniques NASA-KSC's earth resources benefits from space Urban and regional land use analysis: CARETS and applied to strip mining in Maryland and West Virgin exploration p0075 A75-13592 p0039 N75-15133 census cities experiment package [NASA-TM-X-70798] CARBON DIOXIDE [E75.10100] n0025 N75-15118 COASTAL CURRENTS Optical-acoustic detection of low concentrations of CLASSIFICATIONS California nearshore processes -currents and sediments FRTS 1 drogen fluoride, nitrous oxide, and carbon dioxide in gases clustering algorithm for p0048 A75-17179 using pulsed hydrogen-fluoride laser radia p0002 A75-17113 Surface currents along the California coast observed on RTS imagery p0048 A75-17180 p0063 A75-11443 The PUCE programme for terrain evaluation fo ERTS imagery CARBON DIOXIDE LASERS ngineering purposes. 2: Procedures for terrain Application of a single laser Doppler system to the easurement of atmospheric winds COASTAL ECOLOGY classificati Coastal wetlands - Prospects for satellite inventor [PB-233847/3] n0029 N75-10625 [NASA-TM-X-64891] p0047 A75-13580 p0021 N75-12462 Digital land-use classification using an ERTS-1 i p0023 N75-14226 CARBON MONOXIDE Inventories of Delaware's coastal vegetation and land-use On the detectability of atmospheric carbon monoxide by parison of various numerical techniques for ication of multispectral data. Application to geology microwave remote sensing p0013 A75-17129 Application of ERTS-1 data to the protection and CARIBBEAN SEA --- ERTS-1 imagery of Rhone Valley (France) p0038 N75-14244 detection by ERTS-1 of wind-induced ocean surface nagement of New Jersey's coastal en p0042 A75-17233 Ground-truth measurements in relation to aircraft and features in the lee of the Antilles Islands p0041 A75-10644 satellite studies of agricultural land use and land classification in Britain --- ERTS-1 and aerial remote sensing The application of ERTS-1 data to the Corps of Engineers CASCADE RANGE (CA-OR-WA) Civil Works Mission --- for ecological monitoring p0015 A75-18064 Evaluate ERTS imagery for mapping and detection of changes of snowcover on land and on glaciers --- Alaska in Great Britain p0009 N75-14251 The ERTS-1 investigation (ER-600). Volume 2: ERTS-1 Ground-based measurements of solar radiation by nd Washin coastal/estuarine analysis --- Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 [E74-10812] geophysical monitoring for climatic change (USA) --- using pyroheliometers, pyranometers, and radiometers p0049 N75-10539 Study to develop improved spacecraft snow sur-Feasibility of remote sensing for detecting thermal p0018 N75-11527 methods using Skylab/EREP data --- Sierra Nevada. Cascades, and Great Plains pollution. Part 1: Feasibility study. Part 2: Implementation Urban and regional land use analysis: CARETS and plan --- coastal ecology [NASA-CR-134453] [E75-10081] p0073 N75-14200 census cities experiment package n0025 N75-15199 [E75-10056] p0022 N75-13356 CAVITY RESONATORS COASTAL RANGES (CA) Meteorological data. Volume 145 no. 1: The European climatology 1974 meteorological satellite data from the Feasibility study of a Gunn-diode microwave cavity pectrometer as a formaldehyde monitor Fault tectonics and earthquake hazards in the Peninsular Ranges, Southern California [E75-10022] Berlin zone of reception, part 1, 1st quarter [UCID-16488] p0071 N75-11311 p0030 N75-13338 p0023 N75-13479 CELESTIAL GEODESY Meteorological data. Volume 139, no. 4: The European climatology, 1973. Meteorological satellite-data from the COASTAL WATER Earth parameters from global satellite triangulation and Skylab and FRTS-1 investigations of coastal land-use p0064 A75-12985 and water properties in Delaware Bay [AIAA PAPER 74-1220] trilateration Bertin zone of reception, part 4, fourth quarter p0041 A75-12034 Transformation of terrestrial survey data to Doppler n0023 N75-13483 Monitoring and predicting El Nino invasions --- equatorial satellite datum p0054 A75-16569 ERTS image contribution to geological knowledge

covering different types of climates (France, Togo, Dahome

Madagascar)

p0038 N75-14240

Pacific Ocean-atmosphere interactions

p0041 A75-12719

New methods for studying the dynamics of the entire p0016 A75-18108

SUBJECT INDEX Coastal and estuarine studies with ERTS-1 and Skylab p0047 A75-16240 Computer derived coastal water classifications via p0042 A75-17177 spectral signatures Correlation of coastal water turbidity and current circulation with ERTS-1 and Skylab imagery p0048 A75-17181 Ocean internal waves off the North American and African p0042 A75-17183 A review of initial investigations to utilize ERTS-1 data in determining the availability and distribution of living marine resources --- harvest and management of fisheries resources in Mississippi Sound and Gulf p0043 N75-10555 Evaluation of satellite remote sen ing and automatic data techniques for characterization of wetlands and coastal marshlands --- detection and measurement of accretion in estuary system of Atchafalaya River in southern [E75-10025] p0044 N75-13341 Evaluation of satellite remote sensing and automatic data techniques for characterization of wetlands and coastal marshlands --- Atchafalaya River Basin, Louisiana [E75-10028] The ERTS-1 investigation (ER-600). Volume 2: ERTS-1 coastal/estuarine analysis --- Galveston Bay, Texas p0051 N75-13371 [NASA-TM-X-58118] Interpretation of ERTS-1 data for investigating the French western coastal zones (Fralit program) p0051 N75-14230 COHERENT LIGHT Coherent optics in mapping: Proceedings of the Tutorial Seminar and Technology Utilization Pro N.Y., March 27-29, 1974 Program, Rochester, p0027 A75-16676 COLOR PHOTOGRAPHY Special color analysis techniques --- for reconnaissance p0064 A75-12874 Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135 Flood inundation in the southeastern United States from aircraft and satellite imagery p0047 A75-17138 Texture analysis with Fourier series --- for crop identification from color aerial photographs p0004 A75-17187

The interpretation and use of false-colour infra-red and true colour photography of part of Argentina obtained by Skylark earth resources rockets p0067 A75-17192 Optical film density values from color IR photography wetland soils mapping p0005 A75-18073 Evaluation of Skylab imagery as an information service for investigating land use and natural resources --- Long Island, New York

[E74-10730] p0056 N75-10526 Analysis of pseudocolor transformations of ERTSimages of Southern California area --- geological faults and

p0036 N75-13346 Colour composites of ERTS multispectral imagery for applications in hydrology and oceanography --- Italian Mediterranean and North Sea areas p0045 N75-14231 COLORADO

A comparison of Skylab and ERTS data for agricultural crop and natural vegetation interpretation
[AIAA PAPER 74-1219] p0001 A75-12033
Land type analysis for regional land use planning from
photomorphic mapping - An example for Boulder County,

p0013 A75-17137 Geologic and mineral and water resources investigations

Western Colorado, using Skylab EREP p0034 N75-10530 [E74-10802] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques

p0057 N75-10531 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques --- hydrology. ecology, geology, vegetation, and mineral deposits

[275-10002] p0078 N75-12390 Inventory of forest and rangeland resources, including forest stress --- Colorado, Georgia, and Black Hills, South Dakota

[E75-10032] p0007 N75-12402 Index of Skylab data available at Colorado School of

-- multispectral photography of Colorado p0072 N75-12410
Geologic and mineral and water resources investigations
Western Colorado. using Skylab EREP data --- Utah
75-100431 [E75-10041]

[E75-10043] p0058 N75-12412

Geologic and mineral and water resources investigations m Colorado, using Skylab EREP data p0037 N75-14190 [E75-10071]

An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques — geology, hydrology, vegetation, and snow cover [E75-10072] p0037 N75-14191

An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using utomatic data processing techniques

p0073 N75-15122 Geologic and mineral and water resources investigations western Colorado, using Skylab EREP data

p0025 N75-15123

COLORADO PLATEAU (US)

Preliminary results in the comparison of Skylab, ERTS and RB-57 images for the detection of lineaments and fractures in Precambrian, Paleozoic, and late Tertiary rocks near the Colorado Plateau, North-Central [E75-10060] p0037 N75-13360 COMMUNITIES

Earth Resources Technology Satellite-1 (ERTS-1) data and anthropology: Use of these data in carrying capacity estimates for sites in Upper Volta and Niger [E75-10001] p0078 N75-11406

Utilization of ERTS-1 imagery in cultivation and settlement site identification and carrying capacity estimates in Upper Volta and Niger [E75-10029]

p0008 N75-13345 COMPARATORS

The ITC 8-image comparator and its possible use in ERTS p0060 N75-14222 COMPUTER GRAPHICS

Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing. natural resources, social and economic data

Coastal wetlands analysis from ERTS MSS digital data didied spectral measurements p0048 A75-17178 Color terrain man of Vallanda (1997) and field spectral measurements pOO48 A75 17178
Color terrain map of Yellowstone National Park, computer-derived from ERTS MSS data

p0004 A75-17185 Image selection system --- computerized data storage

and retrieval system [NASA-CR-2475] p0058 N75-12422

COMPUTER PROGRAMMING

Use of feature extraction techniques for the texture and context information in ERTS imagery: Spectral and textural processing of ERTS imagery --- classification of Kansas n0059 N75.13335

[E75-10011] Planning applications in east central Florida --- maximum likelihood classification programming [E75-10015] p0021 N75 13337

Reformatting Skylab and MSS data over northern

[E75-10016] p0061 N75-15101 Developing processing techniques for Skylab data [E75-10082] p0061 N75-1 p0061 N75-15102

COMPUTER PROGRAMS

SICLOPS - A system of computer programs for rectified mapping of airborne scanner imagery p0055 A75-17156

The GLOBAL computer program and GLOBEMAP: A special purpose three-dimensional plotter subroutine [AD-785869] p0032 N75-15141

COMPUTER SYSTEMS DESIGN

An analysis of applications development systems for remotely sensed, multispectral data for the Earth Observations Division of the NASA Lyndon B. Johnson

Space Center [NASA-CR-140345] D0071 N75-11447

COMPUTER TECHNIQUES

Satellites and computer in the service of meteorology: WWW - The global weather observation system p0053 A75-11274

Electronics in meteorological instrumentation p0065 A75-14726

Mapping soil features from multispectral scanner data p0027 A75-15435

Design concepts for land use and natural resource inventories and information systems p0013 A75-17133 Interactive computer processing for land use planning p0054 A75-17134

A hydrogeomorphic approach to evaluating flood potential in central Texas from orbital and suborbital remote sensing p0047 A75-17140

of MSS data Adaptive processing of decision-directed Kalman filter data using a p0055 A75-17151 Implementation of an advanced table look-up classifier for large area land-use classification p0014 A75-17153

Modifications in a computer-implemented method for the detection and extraction of objects in aerial photographs DO055 A75-17154

Automatic classification of eutrophication of inland lakes p0048 A75-17164

Computer derived coastal water classifications via p0042 A75-17177 spectral signatures

Geometronics-earth resource measurements and displays the Forest Service p0005 A75-18063 for the Forest Service Automatic mapping of strip mine operations from

spacecraft data p0034 A75-18065 Early results from the experimental Automated Image Data Extraction System --- for aerial terrain sensing

p0069 A75-18072 Automatic pattern recognition and mapping utilizing ermal infrared imagery p0070 A75-18074

thermal infrared imagery Evaluation of Skylab imagery as an information service for investigating land use and natural resources --- Long York

[E74-10730] p0056 N75-10526 Remote sensing of ocean current boundary layer ---Florida

[E74-10801] p0042 N75-10529 An interdisciplinary analysis of multispectral satellite data

for selected cover types in the Colorado Mountains, using utomatic data processing techniques [F74-10803] p0057 N75-10531 Use of a digital computer to construct isoline maps based

nodel of terrain p0057 N75-11453 [AD-784170] Computer and photogrammetric general land use study of central north Alabama

[NASA-TR-R-431] n0022 N75-13372 Fully automated compilation of gravity and geophysical

maps [AD-785838] o0032 N75-15147

COMPUTERIZED SIMULATION A simulation of the San Andreas fau

p0033 A75-11068 On evaluating compliance with air pollution levels 'not to be exceeded more than once a year

p0011 A75-12803

COMSAT PROGRAM

The community satellite. III p0053 A75-11822

CONCORDE AIRCRAFT

The 1973 program of measurement of the minor constituents of the stratosphere using the Concorde 001 p0063 A75-11720

CONFERENCES

Energy development: Proceedings of the Energy Sources Conference, Anaheim, Calif., July 14-19, 1974 p0075 A75-12986

Technology today for tomorrow; Proceedings of the Eleventh Space Congress, Cocoa Beach, Fla., April 17-19, 1974. Volumes 1 & 2 p0011 A75-13576

Coherent optics in mapping: Proceedings of the Tutorial eminar and Technology Utilization Program, Rochester, Y., March 27-29, 1974 p0027 A75-16676

International Symposium on Remote Sensing of Environment, 9th, University of Michigan, Ann Arbor, Mich., April 15-19, 1974, Proceedings, Volumes 1, 2 & 3

p0066 A75-17101 American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings

p0056 A75-18052 Geodesy and physics of the earth; Proceedings of the Second International Symposium, Potsdam, East German May 7-11, 1973. Parts 1 & 2 p0028 A75-1810 p0028 A75-18106

Conference on Weather Forecasting and Analysis, 5th, St. Louis, Mo., March 4-7, 1974, Preprints p0016 A75-18151

Conference on Environmental Sensors and Applications, London, England, November 18, 19, 1974, Proceedings p0070 A75-19265

Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results

[NASA-SP-356] p0077 N75-10549 Observation and measurement of atmospheric pollution

p0018 N75-11476 Proceedings of the UCLA International Conference on Radiation and Remote Probing of the Atmosphere ---

emphasizing radiative transfer [NASA-CR-140816] p0078 N75-11549

European earth-resources satellite experiments
[ESRO-SP-100] p0081 N75-14216

The Proceedings of the Skylab Life Sciences Symposium,

[NASA-TM-X-58154-VOL-1] n0081 N75-14358

CONGRESSIONAL REPORTS

National Environmental Policy Act of 1969. Environmental indices, status of development pursuant to Sections 102(2)(B) and 204 of the act [GPO-25-314] D0077 N75-10617

Earth resources survey system [GPO-41-665] D0078 N75-12416

Earth resources satellites [GPO-40-560] p0078 N75-13010

CONNECTICUT

Investigation of land use of northern megalopolis using ERTS-1 imagery INASA-CR-24591 n0017 N75-10545

Urban and regional land use analysis: CARETS and ensus cities experiment package ... New Haven, census cities experiment package --- New Haven, Connecticut; Pontiac, Michigan; Phoenix, Arizona; District of Columbia

[E75-10091] p0024 N75-15109

CONSERVATION

Land use mapping using ERTS-1 imagery for The Four Rivers Resource Conservation and Development District p0016 A75-18070

CONTAMINANTS

Skylab induced environment --- spacecraft-originated contaminant effects

AIAA PAPER 74-1225) p0011 A75-12037 CONTAMINATION

Determination of points of entry for potential contaminants into limestone aquifers using thermal infrared imagery

-CR-120540] pCO21 N75-12546 CONTINENTAL SHELVES

Offshore investigation: Producible shut-in leases as of January 1974, phase 2 --- an estimation of natural gas

reserves in offshare wells p0036 N75-11458 Use of a digital computer to construct isoline maps based

on a stereomodel of terrain [AD-784170] p0057 N75-11453

CONTROL MOMENT GYROSCOPES Skylab. II - The three month vigil p0065 A75-13963 SUBJECT INDEX

COOK INLET (AK) COOK INLET (AK) DATA ACQUISITION DATA SYSTEMS The application of ERTS-1 data to the Corps of Engineers Orientation of methods of information processing for the Automated remote measurement system in agricultural Civil Works Mission --- for ecological monitoring meteorology --- for crop survey by aerial photometry control of terrestrial resources p0015 A75-18064 p0064 A75-12947 [IAF PAPER 74-140] p0053 A75-13758 COORDINATES Comparison of existing earth resources data processing ERTS-1 and data for national land use planning poo75 A75-13581 Geographic correlation of television pictures obtained from weather satellites p0071 N75-11432 facilitie [IAF PAPER 74-144] p0053 A75-13760 Extraction of urban land cover data from multiplexed rothetic aperture radar imagery p0054 A75-17117 Geographical correlation of TV and IR images obtained Electronics in meteorological instrumentation synthetic aperture radar imagery p0065 A75-14726 from weather satellites --- cloud cover photograph Multi-aspect techniques in remote sensing --- for MSS pools A75-17217 p0071 N75-11433 Interactive machine assessment of critical land resources System for electronic transformation and geographic using ERTS-1 data p0013 A75-17149 Remote sensing and land use inventory and mapping correlation of satellite television information Earth resources sensor data handling system: NASA p0015 A75-18057 p0071 N75-11434 photography JSC version Geographical correlation of earth cloud cover pictures obtained from unoriented satellites p0071 N75-11435 DATA COLLECTION PLATFORMS [NASA-CR-140334] p0072 N75-12418 Surface water modeling Everglades Water Basin, DATA TRANSMISSION Water survey of Canada: Application for use of ERTS-A r retransmission of water resources data Florida [E74-10814] p0049 N75-10540 Spectral reflectance studies on mineral deficiency in co p0003 A75-17172 niants ERTS program of the US Army Corps of Engineers --water resources p0042 N75-10554 75-10026] p0050 N75-13342 Water survey of Canada: Application for use of ERTS-A [E75-10026] COSMIC RAYS Investigations of diffuse cosmic gamma radiation in the Water survey of Canada: Application for use of ERTS-A for retransmission of water resources data [F75-10053] range 28 keV-4.1 MeV p0015 A75-17667 p0050 N75-13353 retransmission of water resources data COST ANALYSIS p0049 N75-11409 DÉFOLIATION [E74-10805] Forest defoliation assessment with satellite imagi The ERTS cost-benefit study Water survey of Canada: Application for use of ERTS-A [PB-234065/1] p0003 A75-17171 p0079 N75-13373 retransmission of water resources data Life cycle cost study of Army Spectrometric Oil Analysis p0050 N75-13342 DELAWARE [E75-10026] Program (ASOAP) Inventories of Delaware's coastal vegetation and land-use Water survey of Canada: Application for use of ERTS-A utilizing digital processing of ERTS-1 imagery p0004 A75-17188 p0073 N75-14870 or retransmission of water resources data CROP GROWTH p0050 N75-13353 [E75-10053] Determining range condition and forage production potential in California from ERTS-1 imagery Urban and regional land use analysis: CARETS and Monitoring physical and chemical parameters of Delaware census cities experiment package Bay waters with an ERTS-1 data collection platform [E75-10090] p0051 N75-1 p0003 A75-17169 [E75-10019] p0020 N75-12398 DO051 N75-15108 Investigation into the spectral signature of agricultural Remote sensing geophysics from Skylab --- rationing-type DATA PROCESSING p0004 A75-17189 crops during their state of growth algorithms to enhance multispectral data Automated image processing in forestry 75-10052] Radar cross sections of vegetation canopies determined p0053 A75-10435 p0030 N75-13352 Urban and regional land use analysis: CARETS and ensus cities experiment package --- New Haven, by monostatic and bistatic scatterometry Satellites and computer in the service of meteorology: census cities experiment package --- New Haven, Connecticut: Pontiac, Michigan: Phoenix, Arizona; District p0005 A75-17190 WWW - The global weather observation system p0053 A75-11274 Applications of remote sensing by the state of ebraska p0069 A75-17232 Comparison of existing earth resources data processing of Columbia Nebraska [E75-10091] p0024 N75-15109 Detection of moisture and moisture related phenomena Urban and regional land use analysis: CARETS and [IAF PAPER 74-144] p0053 A75-13760 from Skylab --- Texas and Kansas census cities experiment package [E75-10100] [E75-10073] Early results from the experimental Automated Image Data Extraction System --- for aerial terrain sensing p0069 A75-18072 p0008 N75-14192 p0025 N75-15118 CROP IDENTIFICATION DELAWARE RIVER BASIN (US) A comparison of Skylab and ERTS data for agricultural Skylab and ERTS-1 investigations nd water properties in Delaware Bay ns of coastal land-use JSC earth resources data analysis capabilities available crop and natural vegetation interpretation [AIAA PAPER 74-1219] DOOD1 A75-12033 [AIAA PAPER 74-1220] IAA PAPER 74-1220] p0041 A75-12034 Coastal and estuarine studies with ERTS-1 and Skylab [NASA-TM-X-72029] n0057 N75-10570 Agricultural inventory techniques p0001 A75-12872 Correlation and registration of ERTS multispectral high-altitude imagery p0047 A75-16240 Correlation of coastal water turbidity and current magery --- by a digital processing technique Automated remote measurement system in agricultural p0058 N75-12420 [NASA-CR-134294] circulation with ERTS-1 and Skylab imagen meteorology --- for crop survey by aerial photometry Spectroradiometric and multispectral scanner over Lake p0064 A75-12947 p0048 A75-17181 Monroe Monitoring physical and chemical parameters of Delaware clustering algorithm for unsupervised [E75-10066] p0072 N75-13366 Bay waters with an ERTS-1 data collection platform [E75-10090] p0002 A75-17113 classification Evaluation of ERTS-1 data by analogue and digital rechargus --- Germany and Italy Processing of ERTS CCT imagery of Sweden. A pilot study Generalisation of ERTS data for global-scale Agricultural resources investigations s in Southern France p0004 A75-17173 and Northern Italy DELTAS Usefulness of remote sensing techniques for the environmental monitoring of the Sacramento-San Joaquin First results from the crop identification technology Generalisation of ERTS data for investigations --- Central Valley, California assessment for remote sensing /CITARS/ p0004 A75-17175 DEPTH MEASUREMENT n0081 N75-14224 Texture analysis with Fourier series --- for crop Reformatting Skylab and MSS data over northern identification from color aerial photographs Michigan [E75-10048] p0004 A75-17187 p0061 N75-15101 [E75-10016] Developing processing techniques for Skylab data [E75-10082] Land use classification and ground truth correlations from DESERTS simultaneously acquired aircraft and ERTS-1 MSS data p0061 N75-15102 p0014 A75-17188 satellite visible and infrared measurements Analysis of Skylab MSS data, and calibration of thermal The interpretation and use of false-colour infra-red and data over Lake Monroe, Indiana true colour photography of part of Argentina obtained p0073 N75-15113 [E75-10095] Skylark earth resources rockers p0067 A75-17192 Earth resources data analysis pro [NASA-CR-141477] DATA PROCESSING EQUIPMENT rces data analysis program, phase 2 Statistical separability of agricultural cover types in p0074 N75-15333 [E74-10818] subsets of one to twelve spectral channels p0005 A75-17216 Orientation of methods of information processing for the control of terrestrial resources [IAF PAPER 74-140] The application of pattern classification techniques to Skylark earth resources rocket imagery p0053 A75-13758 DESIGN ANALYSIS p0056 A75-19268 DATA RECORDING Methods for the automatic recording of atmospheric dust A study of the early detection of insect infestations and density/distribution of host plants [E75-10062] concentrations and some of the results obtained p0057 N75-11502 p0008 N75-13362 DATA REDUCTION CROP VIGOR at 1.43 GHz More information and less data from space --- data Spectral reflectance studies on mineral deficiency in corn [NASA-CR-2458] ction in remote sensing systems p0063 A75-11730 niants p0003 A75-17172 DIFFUSE RADIATION Remote detection of soil surface moisture Thermal behaviour of some rice fields affected by a Investigations of diffuse cosmic gamma radiation range 28 keV-4.1 MeV p0015 A75p0003 A75-17146 yellows-type disease p0004 A75-17174 Reduction of information redundancy in ERTS-1 and CROPS DIGITAL DATA EREP data. A proposal for an experimental study The value of improved (ERS) information based on p0060 N75-14225 domestic distribution effects of U.S. agriculture crops
[NASA-CR-141046] p0007 N75-12423 using ERTS-1 data Developing processing techniques for Skylab data p0061 N75-15102 [E75-10082] CRUDE OIL Earth resources data analysis program, phase 2 and field spectral measurements Crude and refined petroleum oil structured luminescence [NASA-CR-141477] p0074 N75-15333 signatures induced by UV laser or lamp and their remote DATA RETRIEVAL Michigan test site sensing applications p0033 A75-17207 Image selection system --- computerized data storage Prediction of the Fraunhofer line detectivity of luminescent materials --- crude and refined oils, effluents Image services of property of the property of DIGITAL SIMULATION p0068 A75-17221 and vegetation Generalisation of ERTS data for investigations --- Central Valley, California Life cycle cost study of Army Spectrometric Oil Analysis [AD-784170] o0081 N75-14224 n (ASOAP) Program (ASO [AD-786501] p0073 N75-14870 DATA SAMPLING

DAHOMEY ERTS image contribution to geological knowledge overing different types of climates (France, Togo, Dahomey, fadagascar) p0038 N75-14240 Madagascar)

D

More information and less data from space reduction in remote sensing systems p0063 A75-11730
On evaluating compliance with air pollution levels 'not to be exceeded more than once a year p0011 A75-12803

DATA STORAGE

Image selection system --- computerized data storage n0058 N75-12422 [NASA-CR-2475]

p0025 N75-15135 Skylab water depth determination --- Lake Erie and Lake p0050 N75-13348 The detection of dust storms over land and water with p0012 A75-16675 A study to explore the use of orbital remote sensing to determine native arid plant distribution --- Arizona Regional Ecological Test Site p0006 N75-10544 ERTS image contribution to geological knowledge covering different types of climates (France, Togo, Dahomey, Madagascar) p0038 N75-14240 Design concepts for land use and natural resource inventories and information systems . p0013 A75-17133 DIELECTRIC PROPERTIES Measurements of the dielectric properties of sea water p0045 N75-13496 p0015 A75-17667 Interactive machine assessment of critical land resource p0013 A75-17149 Coastal wetlands analysis from ERTS MSS digital data of field spectral measurements p0048 A75-17178 Developing processing techniques for Skylab data --p0058 N75-12409 Use of a digital computer to construct isoline maps based on a stereomodel of terrain p0057 N75-11453 DIGITAL TECHNIQUES Oil exploration needs for digital processing of imagery p0033 A75-10437 clustering algorithm for unsupervised unsupervised crop p0002 A75-17113 classification Correlation and registration of ERTS multispectral imagery --- by a digital process* technique p0058 N75-12420 Digital land-use classifica n ERTS-1 image √75-14226

A demonstration of ERTS-1 analog and digital techniques applied to strip mining in Maryland and West Virginia **EARTH MOVEMENTS** The economic value of remote sensing of earth resources Geodesy and physics of the earth; Proceedings of the from space: An ERTS overview and the value of continuity [NASA-TM-X-70798] p0039 N75-15133 Volume 3: Intensive use of living resources Part 1: Overview Second International Symposium, Potsdam, East German -11, 1973. Parts 1 & 2 p0028 A75-18106 DISPLAY DEVICES New methods for studying the dynamics of the entire p0016 A75-18108 [NASA-CR-141265] p0079 N75-14205 p0005 A75-18063 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources, for the Forest Service DISTORTION The effect of the secular change of the terrestrial gravity Geometric distortions of opticommercial panoramic field on elevation level differences p0028 A75-18122 agriculture. Part 2: Distribution effects p0057 N75-11425 ision systems On the determination of secular variation in the earth's p0079 N75-14208 [NASA-CR-141194] DISTRICT OF COLUMBIA gravity field and of secular polar motion The economic value of remote sensing of earth resources p0017 N75-10560 o0029 A75-18125 Land use and mapping from space: An ERTS overview and the value of continuity Urban and regional land use analysis: CARETS and **EARTH PLANETARY STRUCTURE** of service. Volume 3: Intensive use of living resources, agriculture. Part 3: The integrated impact of improved (ERS) information on US agricultural commodities Plutonic structure of the earth's crust on space imag ensus cities experiment package p0020 N75-12398 DO027 A75-17125 [E75-10019] Remote sensing geophysics from Skylab --- rationing-type Elements of the deep structure of the earth's crust on p0079 N75-14207 [NASA-CR-141262] algorithms to enhance multispectral data [E75-10052] the space images of the East Caucasus The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rengeland p0080 N75-14208 p0030 N75-13352 p0028 A75-17228 Urban and regional land use analysis: CARETS and ansus cities experiment package --- New Haven, Applications of satellite geodesy in the exploration of a internal structure of the earth p0029 A75-18123 census cities experiment package --- New Haven, Connecticut; Pontiac, Michigan; Phoenix, Arizona; District the internal structure of the earth EARTH RESOURCES The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 5: Inland water resources of Columbia Economic effectiveness of using the space systems of [E75-10091] p0024 N75-15109 distant indication of natural resources
[IAF PAPER 74-165] Urban and regional land use analysis: CARETS and p0075 A75-13770 p0080 N75-14209 [NASA-CR-141269] census cities experiment package Investigation of transient earth resources phenomena: The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction [F75-10100] p0025 N75-15118 Continuation study DIURNAL VARIATIONS [NASA-CR-139163] p0073 N75-15131 Morning-evening asymmetry of the thermosphere during geomagnetic disturbance p0011 A75-11103 EARTH RESOURCES PROGRAM ARTH RESOUNCES PROGRAM
Summary of flight performance of
Resources Experiment Package /EREP/ p0011 A75-11103 of the Skylab Earth [NASA-CR-141266] p0080 N75-14210 DOPPLER EFFECT The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a nationwide Transformation of terrestrial survey data to Doppler stellite datum p0054 A75-16569 p0076 A75-17211 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources, agriculture. Parl 2: Distribution effects DOPPLER RADAR The wind vector in a layer over water p0041 A75-15179
Measurement of sea surface currents using airborne
Doppler radar and inertial navigation systems land cover information system p0080 N75-14211 [NASA-CR-141267] p0079 N75-14206 [NASA-CR-141194] The economic value of remote sensing of earth resources EARTH ROTATION p0042 A75-17203 from space: An ERTS overview and the value of continuity Problems and consequences or modern the global tectonics for geodesy and gravimetry p0028 A75-18109 DRAINAGE PATTERNS of service. Volume 7: Nonreplenishable natural resources:
Minerals, fossil fuels and geothermal energy sources
[NASA-CR-141264] p0080 N75-14212 Geologic interpretation of ERTS-1 satellite images for West Aswan area, Egypt p0033 A75-17109 Evaluation of ERTS-1 and aircraft data for assessing The economic value of remote sensing of earth resources internal drainage in irrigated agriculture Spectrophotometric observations of earth from optical data processing analysis of stream patterns achibited on ERTS-1 imagery po056 A75-17219
A Skylab program for the International Hydrological from space: An ERTS overview and the value of continuity of service. Volume 8: Atmosphere [NASA-CR-141191] [IAF PAPER 74-156] p0065 A75-13765 Signatures of various earth surfaces measured by the imbus-5 microwave spectrometer p0086 A75-17119 n0080 N75-14213 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity Nimbus-5 microwave spectrometer Decade (IHD) --- Lake Ontario Basin The calculation of cloud shadows in modelling of the of service. Volume 9: Oceans [E75-10039] p0050 N75-12408 p0014 A75-17204 p0080 N75-14214 earth's surface survey from space [NASA-CR-141270] DUST The economic value of remote sensing of earth resources Orbital geology --- an aid to mineral exploration and Measurements of the extraterrestrial solar radiant flux from 2981 to 4000 A and its transmission through the mapping the earth's surface [NASA-TT-F-16000] from space: An ERTS overview and the value of continuity of service. Volume 10: Industry [NASA-CR-141192] n0035 N75-10887 earth's atmosphere as it is affected by dust and ozone n0080 N75-14215 **EARTHQUAKES ECONOMIC DEVELOPMENT** Satellite techniques for the study of secular variations Methods for the automatic recording of atmospheric dust Evaluation of ERTS data utilization in deve p0064 A75-12980 Dust storms according to data of space research --spaceborne photography of arid zones
[NASA-TT-F-16021]
BY STORMS concentrations and some of the results obtained Nuntries
Remote sensing for regional resource analysis
p0015 A75-18061 countries Fault tectonics and earthquake hazards in the Peninsular Ranges, Southern California p0035 N75-11408 [F75-10005] **ECONOMIC FACTORS** Fault tectonics and earthquake hazards in the Peninsular Economic effectiveness of using the space systems of Ranges, Southern California [E75-10022] DUST STORMS distant indication of natural resources p0030 N75-13338 The detection of dust storms over land and water with AF PAPER 74-165] p0075 A75-13770 satellite visible and infrared measurements --- isoseismal maps p0072 N75-13455 Catalog of earthquake photographs **ECOSYSTEMS** p0012 A75-16675 BESMEX: Bering Sea marine mammal experiment --with the primary target species being the walrus and
bowhead whale Fault tectonics and earthquake hazards in the Peninsular Ranges, Southern California --- Gulf of California Ε [E75-10074] p0031 N75-14193 NASA-TM-X-62399) p0008 N75-13503 Fault tectonics and earthquake hazards in the Peninsular Ranges, Southern California
[E75-10108] p0039 N75-15124 EGYPT EARTH (PLANET) Geologic interpretation of ERTS-1 satellite ima p0033 A75-17108 The earth as a radio source - Terrestrial kilor East Aswan area, Egypt p0011 A75-11211 Geologic interpretation of ERTS-1 satellite images to **ECOLOGY** The radio emission of the earth as a pla West Aswan area, Egypt g0033 A75-17109 Water-management models in Florida from ERTS-1 ata p0047 A75-13579 Geologic interpretation of infrared thermal images in East p0076 A75-15319 data EARTH ALBEDO Qatrani area, Western Desert, Egypt p0033 A75-17215
ELECTRIC POWER PLANTS Microbial abundance and thermoluminescence Determination of the earth's aerosol albedo using Skylab antarctic dry valley soils DO005 A75-17973 data --- Lake Michigan Trace gas analysis of power plant plumes via aircraft measurement - 03, NOx, and SO2 chemistry Skylab-EREP investigations of wetlands ecology --- New p0064 A75-13148 p0025 N75-15114 EARTH ATMOSPHERE y and Chesapeake Bay [E74-10816] Spectrophotometric observations of earth from Soyuz-13' p0049 N75-10542 ELECTROMAGNETIC MEASUREMENT
A Shuttle/SpaceLab RF environment su An interdisciplinary analysis of multispectral satellite data [IAF PAPER 74-156] for selected cover types in the Colorado Mountains, using p0065 A75-13765 p0016 A75-19457 automatic data processing techniques --- hydrology. atmospheric effects in Skylab data ELECTROMAGNETIC WAVE TRANSMISSION ecology, geology, vegetation, and mineral deposits [E75-10002] p0020 N75-12391 An examination of whiteouts and transmission through ice fog in the visible and near infrared [DREO-TN-74-18] p0021 N75-12487 [E75-10006] p0078 N75-12390 Experimental evaluation of atmospheric effects on radiometric measurements using the EREP of Skytab ---ERTS-1 contributions to the ecogenetical knowledge of Bonneville Salt Flats, Utah the Llanos of the Orinoco (Colombia and Venezuela) ELECTRON DENSITY (CONCENTRATION) p0023 N75-14245 lce particulates in the mesosphere p0012 A75-16752 ELECTRON FLUX DENSITY [E75-10036] p0020 N75-12406 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 8: Atmosphere [NASA-CR-141191] p0080 N75-14213 Qualitative study of some ERTS-1 images of the Golion Detection of synchronous changes in fluxes of energetic project. Ecological aspects of terrestrial environments p0023 N75-14248 electrons and gamma-quanta during a DO080 N75-14213 disturbance p0064 A75-12317 **ECONOMIC ANALYSIS** ELECTRONIC EQUIPMENT **EARTH CORE** The economic impact of remote sensing data as the source of nonpoint pollution monitoring and control Applications of satellite geodesy in the exploration of An electronic drawing aid for map compiling --- from stereo-orthophotos p0027 A75-10076 mal structure of the earth p0029 A75-18123 p0076 A75-17166 EARTH CRUST Electronics in meteorological instrume ntation The value of improved (ERS) information based on Plutonic structure of the earth's crust on space im DO065 A75-14726 tion effects of U.S. agriculture crops Electrically scanning microwave radiometers -[NASA-CR-141046] p0007 N75-12423 Elements of the deep structure of the earth's crust on le space images of the East Caucasus satellite-borne remote sensing p0058 A75-17202 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity p0028 A75-17228 Emissivities and forward scattering of natural and

(NASA-CR-141268)

[NASA-CR-141193]

Volume 2: Source document

The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service.

p0079 N75-14203

p0079 N75-14204

À.

The effect of the secular change of the terri

TH LIMB phis inc.

Measurement of trace stratospheric gases by inverting

p0028 A75-18122

p0018 N75-11523

d on elevation level differences

thermal infrared limb regionce profiles

p0023 1

EARTH LIMB

p0087 A75-17197

p0075 A75-12986

n-made material at three millimeter v

Energy development; Proceedings of the Energy Sources Conference, Anaheim, Calif., July 14-19, 1974

ENERGY SOURCES

ENERGY TECHNOLOGY Methodological aspects and regional examples of U.S. energy resources - Outlook for the future A study of Minnesota forests and lakes using data from p0075 A75-12999 Earth Resources Technology Satellites snow-cover mapping from ERTS-1 and EREP nd EREP imagery of p0051 N75-14232 **ENERGY TECHNOLOGY** [NASA-CR-140865] p0007 N75-12421 the Swiss Alps Geological features on ERTS-1 images ges of a test area in p0037 N75-14237 ergy development; Proceedings of the Energy Sources Earth resources satellites the west-central Alps --- Italy p0037 N75-14237
ARNICA/ERTS-1 experiment. Results and perspectives
--- image processing for southwest France and Pyrenees Conference, Anaheim, Calif., July 14-19, 1974 [GPO-40-560] p0078 N75-13010 p0075 A75-12986 Investigation of the utility of kinetically controlled and U.S. energy resources - Outlook for the future catalytic reactions to perform environmental determinations p0075 A75-12999 p0081 N75-14246 by parallel spectrophotometric analysis **EUROPEAN SPACE PROGRAMS ENVIRONMENT EFFECTS** p0021 N75-13063 [TID-26594] Geophysical monitoring for climatic change. The NOAA Geoscientific research with rockets and satellites
[IAF PAPER 74-116] p0053 A75-Environmental definition program -- satellite monitoring p0053 A75-13741 p0018 N75-11510 the stratosphere and troposphere Applicability of Skylab remote sensing for detection and monitoring of surface mining activities — strip mining and reclamation in Kentucky and Tennessee EUTROPHICATION [UCID-16565] DO022 N75-13458 The use of remote sensing in limnological studies Environmental satellites and ERTS-1 l imagery p0059 N75-14217 eutrophication of lakes p0048 A75-17163 [E75-10038] p0044 N75-12407 Automatic classification of eutrophication of inland lakes DO048 A75-17164 Usefulness of remote sensing techniques for the Applicability of Skylab remote sensing for detection ar from snacecraft data environmental monitoring of the Sac Monitoring trophic status of inland monitoring of surface mining activities --- evironmental p0025 N75-15135 p0049 A75-18067 effects and reclamation in Tennessee and Kentucky application of ERTS imagery EVERGLADES (FL) **ENVIRONMENTAL QUALITY** Investigation of the environmental change pattern of California nearshore processes currents and sediments FRTS 1 -Surface water modeling Everglades Water Basin. p0048 A75-17179 Florida Japan [E75-10086] [E74-10814] n0024 N75-15104 p0049 N75-10540 Investigation of the environmental change pattern of **ENVIRONMENT MANAGEMENT EXHAUST GASES** Trace gas analysis of power plant plume measurement - 03, NOx, and SO2 chemistry Photographs from balloons - Their use in agronomy and [E75-10086] p0024 N75-15104 plant plumes via aircraft management of environment o0055 A75-17191 ENVIRONMENTAL SURVEYS Trends in the use of remote ser DO064 A75-13148 Airborne survey of major air basins in California EXPERIMENTAL DESIGN DO005 A75-18068 and environmental management p0011 A75-14360 **ENVIRONMENT MONITORING** Planning and management of science programs on Application of ERTS data to enviro ntal invento DO011 A75-13577 Skylab earth resources pro s program p0016 A75-18069 assessment [AIAA PAPER 74-1215] o0075 A75-12030 **ENVIRONMENTAL TESTS** EXTRAVEHICULAR ACTIVITY
Skylab. II - The three month vigil Skylab induced environment -- spacecraft-originated The 1973 program of measurement of the minor constituents of the stratosphere using the Concorde 001 p0063 A75-11720 contaminant effects p0065 A75-13963 [AIAA PAPER 74-1225] p0011 A75-12037 **ENVIRONMENT PROTECTION** F National Environmental Policy Act of 1969. Environmental indices, status of development pursuant to Sections 102(2)(B) and 204 of the act **EPHEMERIDES** Calibration and evaluation of Skylab altimetry for geodetic etermination of the geoid FARM CROPS n0030 N75-13364 [GPO-25-314] [E75-10064] p0077 N75-10617 Improved (ERTS) information and its impact on U.S. ENVIRONMENT SIMULATION EREP markets for agricultural commodities: Summary of flight performance of the Skylab Earth Resources Experiment Package /EREP/ Water-management models in Florida from ERTS-1 economic investigation of production, distribution and net p0047 A75-13579 export effects p0076 A75-17211 ENVIRONMENTAL CONTROL [NASA-CR-141044] p0078 N75-12424 Trends in the use of remote sensing for state resource and environmental management p0005 A75-180 Methodological aspects and regional examples of Utilization of ERTS-1 imagery in cultivation and p0005 A75-18068 snow-cover mapping from ERTS-1 and EREP imagery settlement site identification and carrying capacity estimates p0051 N75-14232 Modern trends in environmental testi in Upper Volta and Niger EROSION [F75-10029] p0008 N75-13345 ENVIRONMENTAL MONITORING FEASIBILITY ANALYSIS · Investigation of the environmental change pattern of Very high resolution radiometer p0064 A75-12109
Monitoring and predicting El Nino invasions --- equatorial Estimation of population density in Tokyo districts from p0076 A75-17103 [E75-10086] D0024 N75-15104 Pacific Ocean-atmosphere interactions On the feasibility of remote monitoring of soil moist FRROR ANALYSIS p0041 A75-12719 A simulation of the San Andreas fault with microwave sensors n0067 A75-17147 Thermal emission of agglomerations as recorded on p0033 A75-11068 inary feasibility study on a synthetic aperture environmental satellite imagery --- of European urban radar /SARSAT/ for earth resources surveys Analytical estimates of precision in determining p0067 A75-17194 parameters of motion of artificial earth satellite whe [IAF PAPER 74-139] n0011 A75-13757 analyzing imprecise measurements with correlated erro Investigation of microwave hologram techniques for The potential benefits for development derived from space Single flight stereo radar capabilities --- for terrain p0028 A75-17201 nolication to earth resources n0067 A75-17195 annlications FILTER WHEEL INFRARED SPECTROMETERS [IAF PAPER 74-147] p0075 A75-13761 Quantitative determination of stratospheric aerosol On the feasibility of remote monitoring of soil moisture th microwave sensors p0067 A75-17147 ERROR CORRECTING DEVICES characteristics [E75-10079] with microwave sensors p0067 A75-17147
Radar cross sections of vegetation canopies determined p0023 N75-14198 SICLOPS - A system of computer programs for re mapping of airborne scanner imagery p0055 A75-17156 Feasibility of using S-191 infrared spectra for geological studies from space --- Nevada by monostatic and bistatic scatterometry **ESTUARIES** p0005 A75-17190 Coastal and estuarine studies with ERTS-1 and Skylab [E75-10089] p0038 N75-15107 Electrically scanning microwave radiometers p0047 A75-16240 FINLAND p0056 A75-17202 satellite-borne remote sensing Evaluation of satellite remote sensing and automatic data at Sodankyla during a Geophysical variations geomagnetic storm, 17-18 December 197*
[REPT-9] The remote Raman spectrometer is a viable instrument techniques for characterization of wetlands and coastal note sensing of the environment marshlands --- detection and measurement of accretion p0030 N75-13432 DO068 A75-17214 in estuary system of Atchafalaya River in southern FIRE DAMAGE Conference on Environmental S ors and Applications, An examination of the extent of fire in the grassland An examination of the southern side of the and savanna of Africa along the southern side of the p0002 A75-17104 London, England, November 18, 19, 1974, Proceedings p0070 A75-19265 The role of lidar in environmental sensing --- laser ceilometer and visual range measurement application FIRE PREVENTION Systems study of an automated fire weather data p0070 A75-19274 An examination of the extent of fire in the grassland A Shuttle/SpaceLab RF environment survey facility [NASA-TM-X-62402] p0006 N75-12277 d savanna of Africa along the southern side of the p0016 A75-19457 FISHES p0002 A75-17104 Modern trends in environmental testing Mathematical modeling for the detection of fish by an Mapping of the major structures of the African rift system p0016 N75-10469 [SLA-73-5931] airborne laser p0065 A75-15953 Ethiopia, Somalia, and Yemen Environmental surveys in Alaska based upon ERTS ata p0017 N75-10551 Preliminary results of fisheries investigation associated ith Skylab-3 p0003 A75-17167 [E75-10024] D0030 N75-13340 Natural-resources mapping of Ethiopia from ERTS-1 nagery p0031 N75-14235 with Skylab-3 ERTS program of the US Army Corps of Engineers --ater resources p0042 N75-10554 Rapid stock assessment of pilchard populations b p0003 A75-17168 water resources FRTS-1 data on Afar tectonics p0031 N75-14239 aircraft-borne remote sensors A comparison of ERTS imagery with conventional aerial Remote sensing applications in marine science programs A review of initial investigations to utilize ERTS-1 data in determining the availability and distribution of living marine resources --- harvest and management of fisheries at VIMS photography for land-resource surveys in less developed [NASA-CR-137475] p0043 N75-10566 Examples from the Rift Valley Lakes Basin, p0061 N75-14249 A survey for the use of remote sensing in the Chesapeake resources in Mississippi Sound and Gulf waters p0043 N75-10555 EUROPE Bay region [NASA-CR-137476] p0043 N75-10571 Use of terrestrial gravity data in combination with satellite The Coho Project: Living resources prediction feasibility study, volume 1 --- meteorological forecasting of fish results for determining geoid heights in Central Europe National Environmental Policy Act of 1969 p0029 A75-18128 Environmental indices, status of development pursuant to concentrations Comparison of the Southern and Northern Hernispl DO044 N75-10681 Sections 102(2)(B) and 204 of the act [PB-234057/8] eneral circulation characteristics as determined by satellite The Coho Project: Living resources prediction feasibility udy. Volume 2: Environmental report [GPO-25-314] p0077 N75-10617 ozone data [UCRL-75556] Application of remote sensing to state and regional p0017 N75-10638 study. Volume 2 [PB-234058/6] p0044 N75-10682 problems --- for Mississippi [NASA-CR-140805] Quality control in global and regional air chemistr monitoring networks: Lessons of the European A The Coho Project: Living resources prediction feasibility Geophysical monitoring for climatic change. The NOAA ogram p0018 N75-11510 Volume 3: System evaluation report Chemistry Network p0018 N75-11540 p0044 N75-10683 Meteorological data. Volume 145 no. 1: The European climatology 1974 meteorological satellite data from the [PB-234059/4]

ne of reception, part 1, 1st quarter

Berlin zone of reception, part 4, fourth quarter

Meteorological data. Volume 139, no. 4: The European

climatology, 1973. Meteorological satellite-data from the

p0023 N75-13479

p0023 N75-13483

FLIGHT CHARACTERISTICS
ERTS 1 flight evaluation report, 23 October 1973 - 23
January 1974 --- orbits 6371 to 7651

ERTS 1 flight evaluation report, 23 January - 23 April 1974 --- orbits 7652 to 8907 [NASA-CR-140810] p0077 N75-10547

[NASA-CR-140712]

p0077 N75-10548

program

Ground-based measurements of solar radiation by geophysical monitoring for climatic change (USA) --- using pyroheliometers, pyranometers, and radiometers

Quality control in global and regional air chemistry monitoring networks: Lessons of the European Ai Chemistry Network p0018 N75-11540

p0018 N75-11527

p0018 N75-11540

GEOLOGICAL FAULTS SURJECT INDEX ERTS 1 flight evaluation report, 23 April 1974-23 July Use of forest vegetation as an indicator in the interpretation and large scale mapping of soils in a forest Problems and consequences of modern hypotheses on 1974 --- orbits 8908 to 10182 the global tectonics for geodesy and gravimetry p0028 A75-18109 o0077 N75-10568 [AD-785967] p0010 N75-15146 Soviet satellite geodesy: Basis and status of Soviet laser FLOOD DAMAGE geodesy and long-base line interferometry [AD-781138/3] pOX FORMALDEHYDE lood inundation in the southeastern United States from Feasibility study of a Gunn-diode microwave cavity pectrometer as a formaldehyde monitor p0047 A75-17138 p0029 N75-10524 aircraft and satellite imagery spectrometer a [UCID-16488] Mapping of the 1973 Mississippi River floods by the OAA-2 satellite p0028 A75-17139 Interaction of marine geodesy, satellite technology and p0071 N75-11311 NOAA-2 satellite cean physics FORWARD SCATTERING p0043 N75-10671 Water resource and hazard planning report for the Clark [NASA-CR-137470] Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 Fork River Valley above Missoula, Missoula County. Status and prospects of development of space geodesy [JPRS-63231] p0029 N75-10869 [PB-236041/0] Calibration and evaluation of Skylab altimetry for geodetic The point visibility meter - A forward scatter instrument FLOOD PLAINS ermination of the geold for the measurement of aerosol extinction coefficient p0070 A75-19266 The use of remote sensing and natural indicators to p0030 N75-13364 [E75-10064] delineate floodplains - Preliminary findings Scientific research and experimental design planning and FOSSIL FUELS p0067 A75-17142 coordination for 1973 The economic value of remote sensing of earth resour-FLOOD PREDICTIONS p0081 N75-14259 from space: An ERTS overview and the value of continuity of service. Volume 7: Nonreplenishable natural resources: Minerals, fossil fuels and geothermal energy sources [NASA-CR-141264] p0080 N75-14212 A hydrogeomorphic approach to evaluating flood potential The GLOBAL computer program and GLOBEMAP: A in central Texas from orbital and suborbital remote sensing imagery p0047 A75-17140 special purpose three-dimensional plotter subroutine [AD-785869] p0032 N75-15141 Water resource and hazard planning report for the Clark Fork River Valley above Missoula, Missoula County, FOURIER SERIES The next problems in the area of the implementation of Texture analysis with Fourier series --- for crop planning for cartographical-geodetic work
[AD-785882] p0 Montana [PB-236041/0] identification from color aerial photographs p0032 N75-15143 p0051 N75-14260 p0004 A75-17187 Optical quantum generators in geodetic measurer [AD-787281] p0032 N75-1 **FOURIER TRANSFORMATION** FLOODS p0032 N75-15237 Application and sensitivity investigation of Fourier transforms for microwave radiometric inversions Geographic applications of ERTS-1 imagery to landscape GEODETIC SATELLITES change --- Mississippi River and Great Smoky Mountains of Tennessee and North Carolina Interaction of marine geodesy, satellite technology and p0070 N75-11310 [NASA-CR-140754] ocean physics [NASA-CR-137470] n0034 N75-10537 [E74-10810] p0043 N75-10671 FLORIDA Agricultural resources investigations in Southern France **GEODETIC SURVEYS** and Northern Italy p0004 A75-17173
Optical processing of ERTS-1 (MSS) images: Examples Water-management models in Florida from ERTS-1 Earth parameters from global satellite triangulation and p0047 A75-13579 p0064 A75-12985 for the Arcachon basin and Rhone delta -Rhone Delta Remote sensing of ocean current boundary layer --p0060 N75-14223 Geodetic aspects of laser distance measurements to the and Arcachon Basin, France moon and radio-interference measurements to quasars Recent and present sedimentation complexity of the [E74-10801] n0042 N75-10529 Rhone delta and the French mediterraneal project) --- ERTS-1 imagery p00 p0027 A75-14646 Surface water modeling Everglades Water Basin, p0045 N75-14228 The gravity field and the definition of stationary sea Interpretation of ERTS-1 data for investigating the French surface topography p0042 A75-18124 [E74-10814] p0049 N75-10540 western coastal zones (Fralit program) On the determination of secular variation in the earth's Planning applications in east central Florida --- maximum likelihood classification programming p0051 N75-14230 gravity field and of secular polar motion ERTS image contribution to geological knowledge covering different types of climates (France, Togo, Dahomey p0029 A75-18125 [F75-10015] n0021 N75-13337 Use of terrestrial gravity data in combination with satellite FLOW MEASUREMENT Madagascar) p0038 N75-14240 Comparison of various numerical techniques for classification of multispectral data. Application to geology results for determining geoid heights in Central Europe From aerospace photographs to forecasting and p0029 A75-18128 Cruise data report. CHAIN 115 leg 2, Southlant calculating flows [NASA-TT-F-16006] nO059 N75-13367 --- ERTS-1 imagery of Rhone Valley (France) -- to survey geophysical parameters p0038 N75-14244 p0029 N75-10641 FLUORESCENCE [AD-784421] Development of an experimental airborne laser remote sensing system for the detection and classification of oil ARNICA/ERTS-1 experiment. Results and perspectives Australian and Soviet gravity surveys along the Australian ··· image processing for southwest France and Py p0068 A75-17206 p0081 N75-14246 p0031 N75-13454 Qualitative study of some ERTS-1 images of the Golion FOAMS Some first accuracy estimates for applications of aerial project. Ecological aspects of terrestrial environments Multi-frequency radiometric measurements of foam and gradiometry [AD-786418] p0023 N75-14248 p0032 N75-14335 a mono-molecular slick --- microwave sensing of ocean surface p0068 A75-17208 Agricultural resources investigations in northern Italy, southern France and Madagascar --- ERTS-B and aerial rice growing and forestor surveys 90009 N75-14252 surface GEODIMETERS Analysis of methods for the geodetic support of complex FOG rice growing and forestry surveys p0009 N75-14252 An examination of whiteouts and transmission through ice fog in the visible and near infrared [DREO-TN-74-18] p0021 N75-12467 sea geophysical surveys HOFER LINE DISCRIMINATORS p0031 N75-13498 Prediction of the Fraunholer line detectivity of luminescent materials --- crude and refined oils, effluents and vegetation p0068 A75-17221 GEOELECTRICITY Report of the geomagnetic and geoelectric observations, FOREST MANAGEMENT 1972 (rapid variations) Automated image processing in forestry FREEZING p0053 A75-10435 [REPT-13] p0032 N75-15228 Irrigation scheduling, freeze warning and soil salinity detecting --- Cameron County, Texas GEOGRAPHY Forest defoliation assessment with satellite imagery p0003 A75-17171 Experience in integrated geological and geographical [E75-10046] p0007 N75-12414 Irrigation scheduling, freeze warning and soil salinity interpretation of space photographs obtained Agricultural resources investigations in Southern France p0036 N75-11437 and Northern Italy p0004 A75-17173 detecting · · · Cameron County, Texas **GEOIDS** p0050 N75-13350
Irrigation scheduling, freeze warning and soil salinity detecting --- Cameron and Starr County, Texas
[E75-10084] Geometronics-earth resource measurements and displays Use of terrestrial gravity data in combination with satellite for the Forest Service p0005 A75-18063 results for determining geoid heights in Central Europe Agriculture, forestry, range resources o0029 A75-18128 p0006 N75-10558 Calibration and evaluation of Skylab altimetry for geodetic Irrigation scheduling, freeze warning and soil salinity detecting --- Cameron and Starr Counties, Texas [E75-10109] p0010 N75-15125 Utilization of EREP data in geological evaluation regional determination of the geoid planning, forest management, and water management in North Carolina [E75-10064] n0030 N75-13364 GEOLOGICAL FAULTS [E75-10063] p0008 N75-13363 A simulation of the San Andreas fault experiment p0033 A75-11068 G Classification and mapping of coal refuse, vegetative Orbital geology --- an aid to mineral exploration and cover types, and forest types by digital processing ERTS-1 data p0054 A75-17110 mapping the earth's surface [NASA-TT-F-16000] GAMMA RAYS p0035 N75-10887 Areal extent of snow in forested regions - A practical Lineaments in basement terrane of the Peninsular Ranges. Detection of synchronous changes in fluxes of energetic and gamma-quanta during a magnetic ce p0064 A75-12317 Southern California [E75-10003] estimation technique using ERTS-1 data p0002 A75-17122 p0035 N75-11407 disturbance Fault sectonics and earthquake hazards in the Peninsular Ranges, Southern California
[E75-10005] p0035 N75-11408 Geographic applications of ERTS-1 imagery to landscape change --- Mississippi River and Great Smoky Mountains Investigations of diffuse cosmic gamma range 28 keV-4.1 MeV p0015 A75-17667 of Tennessee and North Carolina ANALYSIS p0034 N75-10537 Hydrologic significance of lineaments in central Optical-acoustic detection of low concentrations of Systems study of an automated fire weather data hydrogen fluoride, nitrous oxide, and carbon dioxide in gases Tennesse [E75-10030] p0050 N75-12401 using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 [NASA-TM-X-62402] n0006 N75-12277 Fault tectonics and earthquake hazards in the Peninsular Inventory of forest and rangeland resources, including forest stress --- Colorado, Georgia, and Black Hills, South Ranges, Southern California Trace gas analysis of power plant plumes via aircraft measurement - 03. NOx, and SO2 chemistry 5-10022] p0030 N75-13338 p0064 A75-13148 Analysis of pseudocolor transformations of ERTS-1 Dakota [E75-10032] n0007 N75-12402 images of Southern California area --- geological faults and **GAS DETECTORS** A study of Minnesota forests and lakes using data from lineaments Microwave rotational spectroscopy: A physical technique [E75-10037] Earth Resources Technology Satellites p0036 N75-13346 for specific pollutant monitoring p0018 N75-11529 Preliminary results in the comparison of Skylab, ERTS and RB-57 images for the detection of linearments and fractures in Precambrian, Paleozoic, and late Tertiary rocks on and near the Colorado Plateau, North-Central Arizona p0007 N75-12421 GAS SPECTROSCOPY The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] p0080 N75-14208 The potentialities for determining content of minor gas constituents in the atmosphere by satellites p0015 A75-17932 [E75-10060] 60037 N75-13360 Fault tectonics and earthquake hazards in the Peninsular Agricultural resources investigations in northern Italy, southern France and Madagascar --- ERTS-B and aerial GEODESY Skylab altimeter applica [AIAA PAPER 74-1221] dications and scientific results Ranges, Southern California --- Gulf of California Geodesy and physics of the earth; Proceedings of the rice growing and forestry surveys p0009 N75-14252
Future US activities based on experience gained from
ERTS-1 and EREP --- land use, forestry, agriculture, and [E75-10074] p0031 N75-14193 Investigation of the environmental change pattern of

Second International Symposium, Potsdam, East Germany

p0028 A75-18106

[E75-10086]

May 7-11, 1973. Parts 1 & 2

p0024 N75-15104

GEOLOGICAL SURVEYS SUBJECT INDEX

Fault tectonics and earthquake hazards in the Peninsular GEOMORPHOLOGY GRADIENTS Ranges, Southern California [E75-10108] A technology to renovate the search for new m Some first accuracy estimates for applications of aerial p0039 N75-15124 Three studies with p0063 A75-10434 deposits Geologic structure in California: [AD-786418] A hydrogeomorphic approach to evaluating flood potential ERTS-1 imagery in central Texas from orbital and suborbital remote GRASSLANDS [NASA-TM-X-70799] An examination of the extent of fire in the grassland and savanna of Africa along the southern side of the p0039 N75-15132 p0047 A75-17140 imagery GEOLOGICAL SURVEYS Elements of the deep structure of the earth's crust on Geologic interpretation of ERTS-1 satellite images for n0002 A75-17104 the space images of the East Caucasus p0033 A75-17108 East Aswan area, Egypt Grass canopy bidirectional spectral reflectance DO028 A75-17228 Geologic interpretation of ERTS-1 satellite images for p0003 A75-17170 Applications of Skylab EREP photographs to mapping West Aswan area, Egypt p0033 A75-17109

Development of research in the USSR on the use of radar images for geological purposes p0033 A75-17126

Skylab S-192 ratio codes of soil, mineral, and rock spectra GRAVIMETRY of landforms and environmental geology in the Great Plains and Midwest --- Illinois, Iowa, Kansas, Missouri, Nebraska, Problems and consequences of modern hypotheses on the global tectonics for geodesy and gravimetry and South Dakota p0028 A75-18109 p0037 N75-13355 for ratio image selection and interpretation Possibilities of calculating the earth's local gravitational GEOPHYSICAL OBSERVATORIES p0029 A75-18524 p0003 A75-17158 field from satellite observations Geologic interpretation of infrared thermal images in East Qatrani area, Western Desert, Egypt p0033 A75-17215 Remote sensing of rock type in the visible and Report of the geomagnetic and geoelectric observations, 1972 (rapid variations) Australian and Soviet gravity surveys along the Australian calibration line p0032 N75-15228 in the visible and p0033 A75-17220 [BULL-161] GRAVITY ANOMALIES p0031 N75-13454 GEOPHYSICS Geodesy and physics of the earth; Proceedings of the Second International Symposium, Potsdam, East Germany, Application of radar imagery to environmental geologic The gravity field and the definition of stationar The gravity new and the definition of stationary sea surface topography
Use of terrestrial gravity data in combination with satellite results for determining geold heights in Central Europe p0034 A75-17222 May 7-11, 1973. Parts 1 & 2 p0028 A75-18106 Surface compositional mapping in the Wind River Range and Basin, Wyoming by multispectral techniques applied to ERTS-1 data p0028 A75-17224 New methods for studying the dynamics of the entire p0016 A75-18108 p0029 A75-18128 Unsupervised mapping of geologic features and soils in Possibilities of calculating the earth's local gravitational field from satellite observations p0029 A75-18524 Cruise data report. CHAIN 115 leg 2, Southlant p0034 A75-17225 The remote-sensing program of the Geological Survey Alahama p0069 A75-17230 to survey geophysical parameters Fully automated compilation of gravity and geophysical AD-7844211 p0029 N75-10641 of Alahama Geological and geophysical studies from space and Geologic and mineral and water re urces investigations prospects for their development D0032 N75-15147 p0035 N75-11436 in Western Colorado, using Skylab EREP data GRAVITY GRADIOMETERS Remote sensing geophysics from Skylab --- Southern alifornia and Nevada minerals and rocks [E74-10802] p0034 N75-10530
Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State --- Lake Ontario [E74-10802] Some first accuracy estimates for applications of aerial p0030 N75-12413 [E75-10045] [AD-786418] p0032 N75-14335 Remote sensing geophysics from Skylab --- rationing-type p0034 N75-10536 GRAVITY WAVES Environmental surveys in Alaska based upon ERTS p0017 N75-10551 orithms to enhance multispectral data Ocean internal waves off the North American and African p0030 N75-13352 p0042 A75-17183 coasts from ERTS-1 Geologic evaluation and applications of ERTS-1 imagery Remote sensing geophysics from Skylab --- solar heating GREAT BASIN (US) over Georgia p0034 N75-10552
An evaluation of the suitability of ERTS data for the of ground surface and infrared reflectance of rock e Great Basin investigation [E75-10076] p0037 N75-14195 [E75-10057] p0037 N75-13357 purposes of petroleum exploration --- Anadarko Basin in The Great Basin investigation --- photomapping lithology Remote sensing geophysics from Skylab --- stratigraphic analysis of weathered rocks
[E75-10058] p0037 N75-13358 Oklahoma and Texas klahoma and Texas p0035 N75-10553 A study of the usefulness of Skylab EREP data for earth nd geological structures p0038 N75-15120 [E75-10102] resources studies in Australia --- Australia geology 75-10047] p0036 N75-12415 Utilization of EREP data in geological evaluation regional Analysis of methods for the geodetic support of complex GREAT BRITAIN Ground-truth measurements in relation to aircraft and satellite studies of agricultural land use and land classification in Britain --- ERTS-1 and aerial remote sensing sea geophysical surveys planning, forest management, and water managem North Carolina [JPRS-63621] p0031 N75-13498 Fully automated compilation of gravity and geophysical p0009 N75-14251 [E75-10063] p0008 N75-13363 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using GREAT LAKES (NORTH AMERICA) [40-785838] n0032 N75-15147 Observing and forecasting local effects from satellite data
-- mesoscale cloud patterns p0016 A75-18153 GEOPOTENTIAL automatic data processing techniques hydrology, vegetation, and snow cover [E75-10072] p003 p0016 A75-18153 The effect of the secular change of the terrestrial gravity field on elevation level differences p0028 A75-18122 GREAT PLAINS CORRIDOR (NORTH AMERICA) p0037 N75-14191 Applications of Skylab EREP photographs to mapping of landforms and environmental geology in the Great Plains The gravity field and the definition of stationary sea Geologic evaluation of central Italy from ERTS-1 and kylab data p0038 N75-14238 p0042 A75-18124 surface topography and Midwest --- Illinois, Iowa, Kansas, Missouri, Nebraska, On the determination of secular variation in the earth's Feasibility of using S-191 infrared spectra for geological studies from space --- Nevada nd South Dakota gravity field and of secular polar motion [E75-10055] n0037 N75-13355 p0029 A75-18125 Study to develop improved spacecraft snow survey methods using Skylab/EREP data --- Sierra Nevada. Cascades, and Great Plains [E75-10088] p0038 N75-15106 Feasibility of using S-191 infrared spectra for geological studies from space --- Nevada Possibilities of calculating the earth's local gravitational field from satellite observations p0029 A75-18524 [E75-10089] p0038 N75-15107 **GEORGIA** [E75-10081] p0073 N75-14200 Geologic evaluation and applications of ERTS-1 imagery An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using GREAT SMOKY MOUNTAINS (NC-TN) over Georgia p0034 N75-10552 Geographic applications of ERTS-1 imagery to landscape change --- Mississippi River and Great Smoky Mountains of Tennessee and North Carolina Inventory of forest and rangeland resources, including utomatic data processing techniques forest stress --- Colorado, Georgia, and Black Hills, South [E75-10104] p0073 N75-15122 [E74-10810] p0034 N75-10537 GEOLOGY [E75-10032] p0007 N75-12402 An interdisciplinary analysis of multispectral satellite data GROUND STATIONS GEOTHERMAL RESOURCES for selected cover types in the Colorado Mountains, using automatic data processing techniques --- hydrology, Comparison of existing earth resources data processing Geothermal reconnaissance from quantitative analysis of thermal infrared images n0056 A75-17218 [IAF PAPER 74-144] ecology, geology, vegetation, and mineral deposits p0053 A75-13760 e of remote sensing of earth resources Transformation of terrestrial survey data to Doppler tellite datum p0054 A75-16569 p0078 N75-12390 [E75-10002] from space: An ERTS overview and the value of continuity of service. Volume 7: Nonreplenishable natural resources: Minerals, fossil fuels and geothermal energy sources New theoretical models and ratio imaging techniques satellite datum associated with the NASA earth resources spectral Ground-based measurements of solar radiation by geophysical monitoring for climatic change (USA) --- using pyroheliometers, pyranometers, and radiometers information system p0080 N75-14212 [NASA-CR-141264] p0058 N75-12417 [NASA-CR-140357] Applications of Skylab EREP photographs to mapping of landforms and environmental geology in the Great Plains and Midwest --- Illinois, Iowa, Kansas. Missouri, Nebraska, Meteorological data. Volume 145 no. 1: The European climatology 1974 meteorological satellite data from the Berlin zone of reception, part 1, 1st quarter GROUND TRUTH Remote detection of soil surface moistu and South Dakota p0003 A75-17146 p0023 N75-13479 [E75-10055] Meteorological data. Volume 139, no. 4: The European climatology, 1973. Meteorological satellite-data from the First results from the crop identification technology ERTS image contribution to geological knowledge assessment for remote sensing /CITARS/ covering different types of climates (France, Togo, Dahomey, Madagascar) p0038 N75-14240 DÓ004 A75-17175 Berlin zone of reception, part 4, fourth quarter Madagascar) p0023 N75-13483 Land use classification and ground truth correlations from Evaluation of ERTS-1 data by analogue and digital chniques --- Germany and Italy p0059 N75-14218 Comparison of various numerical techniques for simultaneously acquired aircraft and ERTS-1 MSS data lassification of multispectral data. Application to geology -- ERTS-1 imagery of Rhone Valley (France) p0014 A75-17188 Mapping of water bodies in northern Germany from ERTS p0051 N75-14227 The usefulness of imaging passive microwave for rural p0038 N75-14244 and urban terrain analysis p0055 A75-17199 GEOMAGNETISM Ground-truth measurements in relation to aircraft and Morning-evening asymmetry of the thermosphere during a geomagnetic disturbance p0011 A75-11103 Evaluate ERTS imagery for mapping and detection of changes of snowcover on land and on glaciers --- Alaska and Washington satellite studies of agricultural land use and land classification in Britain --- ERTS-1 and aerial remote sensing Magnetopause rotational forms p0011 A75-11221 p0009 N75-14251 in Great Britain [F74-10812] n0049 N75-10539 Solar wind and geomagnetic activity GULF OF CALIFORNIA (MEXICO) GLACIOLOGY p0027 A75-11975 Fault tectonics and earthquake hazards in the Peninsular Ranges, Southern California --- Gulf of California [E75-10074] p0031 N75-14193 ydrology: Norwegian p0051 N75-14233 ERTS-1 images in glaciology and hydro A generator of high-power current pulses of semisinusoidal shape for an aerial variant of the method GLOBAL AIR POLLUTION of transient processes --- for aerial prospecting
p0053 A75-13310 Correlation spectroscopy techniques for the regional and global measurements of air pollution p0018 N75-11505 Global measurements of air pollution from satellites — employing radiometer techniques p0018 N75-11521 Satellite photography of eddies in the gulf loop current Geophysical variations at Sodankyla comagnetic storm, 17-18 December 1971 p0041 A75-11255 Sodankyla during a A review of initial investigations to utilize ERTS-1 date p0030 N75-13432 GLOBAL ATMOSPHERIC RESEARCH PROGRAM in determining the availability and distribution of living Report of the meeting on drifting buoys for the first GARP Report of the geomagnetic and geoelectric observations. marine resources --- harvest and management of fisheries resources in Mississippi Sound and Gulf waters

global experiment [GARP-SPEC-REPT-13]

p0043 N75-10674

p0043 N75-10555

p0032 N75-15228

1972 (rapid variations) [REPT-13]

SUBJECT INDEX INFORMATION SYSTEMS

Offshore investigation: Producible shut-in leases as of Operational use of satellite and high altitude remote IMAGERY January 1974, phase 2 --- an estimation of natural reserves in offshore wells p0036 N75-11 European earth-resources satellite experiments sensing for the generation of input data for water de p0067 A75-17148 p0036 N75-11458 [ESRO-SP-100] p0081 N75-14216 An assessment of the potential contributions to oceanography from Skylab visual observations and hand-held photography --- Pacific Ocean, Atlantic Ocean Water resources --- monitoring and management from Evaluation of ERTS-1 data by analogue and digital p0059 N75-14218 p0043 N75-10559 techniques --- Germany and Italy. An interdisciplinary analysis of multispectral satellite data Processing of ERTS CCT imagery of Sweden. A pilot for selected cover types in the Colorado Mountains, using p0060 N75-14219 and Gulf of Mexico study automatic data processing techniques --- hydrology, [E75-10017] p0044 N75-12396 The potential of small-scale mapping from ERTS-1 ecology, geology, vegetation, and mineral deposits p0060 N75-14221 Oil pollution detection, monitoring and law enforcement [E75-10002] p0078 N75-12390 isiana Gulf Coast waters Generalisation of ERTS data for global-scale Hydrologic significance of lineaments in central p0022 N75-13343 [E75-10027] investigations --- Central Valley, California Remote sensing of ocean current boundary layer --- Gulf p0081 N75-14224 [E75-10030] p0050 N75-12401 of Mexico Meteorological interpretation of high resolution images A Skylab program for the International Hydrological Decade (IHD) --- Lake Ontario Basin [E75-10031] n0045 N75-14185 for remote sensing of earth resources, nO060 N75-14229 Remote sensing of ocean current boundary layer --- Gulf p0050 N75-12408 ERTS-1 images in glaciology and hydrology: Norwegian [E75-10039] p0051 N75-14233 An interdisciplinary analysis of multispectral satellite data experience p0045 N75-14186 Arrimerusciplinary analysis or multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques — geology, hydrology, vegetation, and snow cover [E75-10072] p0037 N75-14191 [E75-10067] Use of ERTS-1 in sea-ice studies p0045 N75-14234 Remote sensing of ocean current boundary layer --- Gulf Natural-resources mapping of Ethiopia from ERTS-1 p0031 N75-14235 n0046 N75-15116 [E75-10098] Geological features on ERTS-1 images of a test area in the west-central Alps --- Italy p0037 N75-14237 Colour composites of ERTS multispectral imagery for GULF STREAM applications in hydrology and oceanography Mediterranean and North Sea areas p0045 f Satellite photography of eddies in the gulf loop of ERTS-1 data on Afar tectonics p0031 N75-14239 p0045 N75-14231 p0041 A75-11255 ERTS-1 images in glaciology and hydrology: Norwegian xperience p0051 N75-14233 ERTS image contribution to geological knowledge **GUNN DIODES** overing different types of climates (France Feasibility study of a Gunn-diode microwave cavity nce, Togo, Dahomey, p0038 N75-14240 Madagascar) spectrometer as a formaldehyde monitor Study of recreational land and onen space using Skylah agery --- snow and ice hydrology of southeast p0071 N75-11311 Recognition and interpretation of spectral signatures of [UCID-16488] Michigan vegetation from aircraft and satellite imagery in Western [E75-10092] p0024 N75-15110 Queensland, Australia p0008 N75-14241 н A Skylab program for the International Hydrological Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 Decade (IHD) --- Lake Ontario Basin [E75-10093] p0052 N75-15111 Preliminary results of the interpretation of ERTS-1 imagery for soil surveys in the Merida region. Spain p0009 N75-14247 HABITATS A Skylab program for the International Hydrological The use of high altitude photography and ERTS-1 imagery for wildlife habitat inventory in Kansas Decade (IHD) --- Lake Ontario Basin p0052 N75-15112 [E75-10094] Qualitative study of some ERTS-1 images of the Golion p0005 A75-18066 An interdisciplinary analysis of multispectral satellite data project. Ecological aspects of terrestrial environments Utilization of Skylab (EREP) system for appraising r selected cover types in the Colorado mountains, using p0023 N75-14248 changes in continental migratory bird habitat [E74-10817] automatic data processing techniques IMAGING TECHNIQUES (F75, 10104) p0073 N75-15122 p0006 N75-10543 Performance comparison of techniques for obtaining stereo radar images p0063 A75-1007 p0063 A75-10075 HYDROMETEOROLOGY Earth Resources Technology Satellite-1 (ERTS-1) data system Automated remote and anthropology: Use of these data in carrying capacity estimates for sites in Upper Volta and Niger A technology to renovate the search for new mineral p0007 N75-12518 p0063 A75-10434 p0078 N75-11406 HYDRONIUM IONS Automated image processing in forestry [E75-10001] Ice particulates in the mesosphere p0012 A75-16752 Utilization of Skylab (EREP) system for appraising p0053 A75-10435 changes in continental migratory bird habitat [E75-10061] p0008 N75-13361 Oil exploration needs for digital processing of imagery p0033 A75-10437 [E75-10061] 1 Utilization of Skylab (EREP) system for appraising ns to mapping p0027 A75-14683 Coherent optics potential application changes in continental migratory bird habitat Deteriorating effects on 3 mm wave passive imagery p0055 A75-17198 n0009 N75-15105 [E75-10087] n examination of whiteouts and transmission through HAZE Multi-aspect techniques in remote sensing --- for MSS nagery p0068 A75-17217 Effect of atmospheric haze and sun angle on automatic ice fog in the visible and near infrared tion of ERTS-1 data p0014 A75-17157 IDREO.TN-74-18] p0021 N75-12467 ICE ENVIRONMENTS The application of remote sensing techniques: Technical HELICOPTERS Development of an experimental airborne laser remote and methodological issues ice particulates in the mesosphere p0012 A75-16752 sensing system for the detection and classification of oil snills p0068 A75-17206 [NASA-CR-140674] New theoretical models and ratio imaging techniques Performance of a hologram matrix radar p0065 A75-14792 Correlation of missions 191, 51M and helicopte associated with the NASA earth resources spectral photography -- aerial photography and mapping of three areas in Test Site 175 in Texas [NASA-CR-134289] p0058 N75-12279 A study of microwave emission properties of sea ice -DJEX 1972 p0041 A75-17120 information system [NASA-CR-140357] AID IFX 1972 nO058 N75-12417 The ITC 8-image comparator and its possible use in ERTS pools in ERTS pools po Study of recreational land and open space using Skylab HEURISTIC METHODS --- snow and ice hydrology of southeast and EREP investigations A clustering classification INDEXES (DOCUMENTATION) algorithm for p0002 A75-17113 Index of Skylab data available at Colorado School of [E75-10092] n0024 N75-15110 HE LAREDS Mines ... multispectral photography of Colorado **ICELAND** Optical-acoustic detection of low concentrations of [E75-10041] p0072 N75-12410 Environmental studies of Iceland with ERTS-1 image hydrogen fluoride, nitrous oxide, and carbon dioxide in gases INDIANA n0012 A75-17105 using pulsed hydrogen-fluoride laser radiation First results from the crop identification technology IDAHO p0063 A75-11443 assessment for remote sensing / CITARS/ Color terrain map of Yellowstone National Park 00004 A75-17175 HIGH ALTITUDE BALLOONS computer-derived from ERTS MSS data Photographs from balloons - Their us Mineral resources, geological structures, and landform in agro p0004 A75-17185 management of environment p0055 A75-17191 p0035 N75-10561 SURVEYS Geothermal reconnaissance from quantitative analysis of ENERGY ELECTRONS Spectroradiometric and multispectral scanner over Lake thermal infrared images p0056 A75-17218 Detection of synchronous changes in fluxes of energetic Monroe ILLINOIS [E75-10066] during a magnetic p0064 A75-12317 p0072 N75-13366 gamma-quanta First results from the crop identification technology Reformatting Skylab and MSS data over northern assessment for remote sensing /CITARS/ HIGH RESOLUTION Indiana p**ÓO**O4 A75-17175 Very high resolution radiometer [E75-10016] p0064 A75-12109 p0061 N75-15101 Applications of Skylab EREP photographs to mapping HIGHWAYS Analysis of Skylab MSS data, and calibration of thermal Multidisciplinary analysis for highway engineering of landforms and environmental geology in the Great Plains and Midwest --- Illinois, Iowa, Kansas, Missouri, Nebraska. data over Lake Monroe, Indiana [E75-10095] p0073 N75-15113 purposes --- Maine nd South Dakota n0025 N75-15121 [E75-10103] INDUSTRIES HOLOGRAMMETRY [£75-10055] p0037 N75-13355 The economic e of remote sensing of earth resources Hologrammetry - Concepts and applications from space: An ERTS overview and the value of continuity Design data collection with Skylab/EREP microwave p0065 A75-15434 of service. Volume 10: Industry [NASA-CR-141192] nstrument S-193 --- Texas, Utah, Illinois, and Minne p0072 N75-14199 DOORO N75-14215 HOLOGRAPHY [E75-10080] Performance of a hologram matrix rada INERTIAL NAVIGATION IMAGE CORRELATORS p0065 A75-14792 Measurement of sea surface currents using airborne Correlation and registration of ERTS multispectral Donnler radar and inertial navigation systematical Terrain measurement from holographic stereogram imagery --- by a digital processing technique p0001 A75-16682 p0042 A75-17203 p0058 N75-12420 [NASA-CR-134294] INFESTATION HURRICANES IMAGE ENHANCEMENT A remote sensing study of Pacific hurricane Ava A study of the early detection of insect infestations and density/distribution of host plants Evaluation of Skylab imagery as an information service p0012 A75-17112 for investigating land use and natural resources --- Hudson p0008 N75-13362 HYDROCARBON FUELS [E75-10062] y and Long Island, New York Exploration for fossil and nuclear fuels from orbital ORMATION [E75-10013] p0059 N75-13336 The value of improved (ERS) information based on altitudes --- results of ERTS program for oil exploration Analysis of pseudocolor transformations of ERTS-1 images of Southern California area --- geological faults and domestic distribution effects of U.S. agriculture crops
[NASA-CR-141046] p0007 N75-12423 [NASA-TM-X-70781] p0017 N75-11413 HYDROGEOLOGY INFORMATION SYSTEMS Optical data processing analysis of stream patterns [E75-10037] p0036 N75-13346 More information and less data from space exhibited on ERTS-1 imagery reduction in remote sensing systems p0063 A75-11730 Remote sensing geophysics from Skylab --- rationing-type HYDROLOGY Design concepts for land use and natural resource A hydrogeomorphic approach to evaluating flood potential algorithms to enhance multispectral data in central Texas from orbital and suborbital remote sensing [E75-10052] p0030 N75-13352 inventories and information systems p0013 A75-17133

IMAGE MOTION COMPENSATION
A motion-compensated spatial scanner -

p0068 A75-17213

imagery p0047 A75-17140
Application of remote sensing to the location of hydrologically active /source/ areas p0066 A75-17141

Evaluation of Skylab imagery as an information service for investigating land use and natural resources [E75-10018] p0020 N75-12397

J

The role of ERTS in the establishment and of a nationwide	Remote sounding of stratospheric temperatures using	J
land cover information system 1	high resolution radiance measurements from the IRIS-D	
[NASA-CR-141045] p0020 N75-12425 INFRARED DETECTORS	[NASA-CR-141130] p0022 N75-13452	JAPAN
Oil-on-Water sensor for detecting oil by emitting	Feasibility of using S-191 infrared spectra for geological	Some features of the urban environment of Tokyo by
wideband IR pulse	studies from space Nevada [E75-10088] p0038 N75-15106	remote sensing infrared surface temperature measurements p0012 A75-17102
[AD-784104] p0072 N75-11454	INLAND WATERS	Estimation of population density in Tokyo districts from
INFRARED IMAGERY	The economic value of remote sensing of earth resources	ERTS-1 data p0076 A75-17103
A satellite view of weather systems over the North American Arctic p0063 A75-10790	from space: An ERTS overview and the value of continuity	Investigation of the environmental change pattern of
Thermal emission of agglomerations as recorded on	of service. Volume 5: Inland water resources	Japan (575 10096) -0034 NZ5 15104
environmental satellite imagery of European urban	[NASA-CR-141269] p0080 N75-14209	[E75-10086] p0024 N75-15104
centers	INSECTS	1,0
[IAF PAPER 74-139] p0011 A75-13757	Forest defoliation assessment with satellite imagery p0003 A75-17171	K
The detection of dust storms over land and water with satellite visible and infrared measurements	A study of the early detection of insect infestations and	
p0012 A75-16675	density/distribution of host plants	KALMAN FILTERS
Use of visible, near-infrared, and thermal infrared remote	[E75-10062] p0008 N75-13362	Adaptive processing of MSS data using a
sensing to study soil moisture p0002 A75-17144	INTERFEROMETRY	decision-directed Kalman filter p0055 A75-17151 KANSAS
Thermal behaviour of some rice fields affected by a	Soviet satellite geodesy: Basis and status of Soviet laser	Moisture detection from Skylab soil studies for Texas
yellows-type disease p0004 A75-17174 Geologic interpretation of infrared thermal images in East	geodesy and long-base line interferometry [AD-781138/3] p0029 N75-10524	and Kansas sites p0002 A75-17145
Qatrani area, Western Desert, Egypt p0033 A75-17215	INTERNATIONAL COOPERATION	Optical data processing analysis of stream patterns
Geothermal reconnaissance from quantitative analysis of	The Canadian ERTS program p0077 N75-10550	exhibited on ERTS-1 imagery p0056 A75-17219
thermal infrared images p0056 A75-17218	A Skylab program for the International Hydrological	The application of ERTS-derived information to the regional land use planning process p0015 A75-18058
Automatic pattern recognition and mapping utilizing	Decade (IHD) Lake Ontario Basin	regional land use planning process p0015 A75-18058 The use of high altitude photography and ERTS-1 imagery
thermal infrared imagery p0070 A75-18074 The application of pattern classification techniques to	[E75-10093] p0052 N75-15111	for wildlife habitat inventory in Kansas
Skylark earth resources rocket imagery	A Skylab program for the International Hydrological	p0005 A75-18066
p0056 A75-19268	Decade (IHD) Lake Ontario Basin [E75-10094] p0052 N75-15112	Land use mapping using ERTS-1 imagery for The Four
Determination of points of entry for potential	INTERPLANETARY MAGNETIC FIELDS	Rivers Resource Conservation and Development District in Kansas p0016 A75-18070
contaminants into limestone aquifers using thermal infrared	Solar wind and geomagnetic activity	Detection of moisture and moisture related phenomena
imagery [NASA-ČR-120540] p0021 N75-12546	p0027 A75-11975	from Skylab Kansas, Oklahoma, and Arkansas
An application of infrared remote sensing techniques to	INVERSIONS	[E75-10035] p0007 N75-12405
ecological problems	Application and sensitivity investigation of Fourier	Use of feature extraction techniques for the texture and
[AD-786028] p0022 N75-13419	transforms for microwave radiometric inversions	context information in ERTS imagery: Spectral and textural
Environmental satellites and ERTS-1 imagery p0059 N75-14217	[NASA-CR-140754] p0070 N75-11310 ION DENSITY (CONCENTRATION)	processing of ERTS imagery classification of Kansas land use
Feasibility of using S-191 infrared spectra for geological	Ice particulates in the mesosphere p0012 A75-16752	[E75-10011] p0059 N75-13335
studies from space Nevada	IONOGRAMS	Applications of Skylab EREP photographs to mapping
[E75-10088] p0038 N75-15106	lonospheric meeting on improved ionograms and	of landforms and environmental geology in the Great Plains
Feasibility of using \$-191 infrared spectra for geological	ionspheric sounding	and Midwest Illinois, Iowa, Kansas, Missouri, Nebraska,
studies from space Nevada [E75-10089] p0038 N75-15107	[REPT-3] p0022 N75-13437	and South Dakota [E75-10055] p0037 N75-13355
The use of color infrared imagery for the study of marsh	IONOSPHERIC SOUNDING	Detection of moisture and moisture related phenomena
buggy tracks p0061 N75-15127	Investigations of the ionosphere by space techniques p0065 A75-13868	from Skylab Texas and Kansas
INFRARED LASERS	Ionospheric meeting on improved ionograms and	[E75-10073] p0008 N75-14192
Remote atmospheric sensing with an airborne laser	ionspheric sounding	Skylab study of water quality Kansas reservoirs and
absorption spectrometer p0066 A75-16485 INFRARED PHOTOGRAPHY	[REPT-3] p0022 N75-13437	lakes
Land type analysis for regional land use planning from	IOWA	[E75-10075] p0051 N75-14194
photomorphic mapping - An example for Boulder County,	Applications of Skylab EREP photographs to mapping	KENTUCKY ERTS-1 imagery interpretation techniques in the
Colorado p0013 A75-17137	of landforms and environmental geology in the Great Plains and Midwest Illinois, Iowa, Kansas, Missouri, Nebraska.	Tennessee Valley land use and soil mapping
Application of remote sensing to the location of hydrologically active /source/ areas p0068 A75-17141	and South Dakota	[E74-10811] p0016 N75-10538
Spectral reflectance studies on mineral deficiency in corn	[E75-10055] p0037 N75-13355	Applicability of Skylab remote sensing for detection and
plants p0003 A75-17172	IRON COMPOUNDS	monitoring of surface mining activities strip mining and
Optical film density values from color IR photography	Mapping exposed silicate rock types and exposed ferric	reclamation in Kentucky and Tennessee [E75-10038] p0044 N75-12407
for wetland soils mapping p0005 A75-18073	and ferrous compounds from a space platform [E75-10008] p0036 N75-12392	Applicability of Skylab remote sensing for detection and
Geographical correlation of TV and IR images obtained	IRRIGATION	monitoring of surface mining activities evironmental
from weather satellites cloud cover photography p0071 N75-11433	Evaluation of ERTS-1 and aircraft data for assessing	effects and reclamation in Tennessee and Kentucky
An interdisciplinary analysis of multispectral satellite data	internal drainage in irrigated agriculture	[E75-10085] p0038 N75-15103
for selected cover types in the Colorado Mountains, using	p0002 A75-17143	KOLMOGOROFF-SMIRNOFF TEST
automatic data processing techniques geology,	Irrigation scheduling, freeze warning and soil salinity	Modifications in a computer-implemented method for the detection and extraction of objects in aerial photographs
hydrology, vegetation, and snow cover	detecting Cameron County, Texas [E75-10046] p0007 N75-12414	p0055 A75-17154
·[E75-10072] p0037 N75-14191	Satellite pictures can help guide Sandhills irrigation	·
INFRARED RADIATION Measurement of trace stratospheric gases by inverting	center-pivot irrigation in Sand Hills Region, Nebraska	•
thermal infrared limb radiance profiles	[E75-10023] p0021 N75-13339	L
p0018 N75-11523	Irrigation scheduling, freeze warning and soil salinity	
Methods of calculating infrared transfer: A review	detecting Cameron County, Texas [E75-10050] p0050 N75-13350	LABORATORY EQUIPMENT
in planetary atmospheres p0019 N75-11560	•	JSC earth resources data analysis capabilities available
Infrared remote sounding p0072 N75-11562	Irrigation scheduling, freeze warning and soil salinity detecting Cameron and Starr County, Texas	to EOD revision B [NASA-TM-X-72029] p0057 N75-10570
INFRARED RADIOMETERS	[E75-10084] p0008 N75-14202	LAKE ERIE
Images from the NOAA-3 very high resolution radiometer over the North Sea and adjoining countries	Irrigation scheduling, freeze warning and soil salinity	Remote sensing for regional resource analysis
p0066 A75-16188		0015 475 40051
	detecting Cameron and Starr Counties, Texas	p0015 A75-18061
Design data collection with Skylab/EREP microwave	detecting Cameron and Starr Counties, Texas [E75-10109] p0010 N75-15125	Skylab water depth determination Lake Erie and Lake
Design data collection with Skylab/EREP microwave instrument S-193 Utah and Texas Rad/Scat	detecting Cameron and Starr Counties, Texas [E75-10109] p0010 N75-15125 ITALY	Skylab water depth determination Lake Erie and Lake Michigan
instrument S-193 Utah and Texas Rad/Scat measurements	detecting Cameron and Starr Counties, Texas [F75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348
instrument S-193 Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189	detecting Cameron and Starr Counties, Texas [F75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN
instrument S-193 ··· Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological	detecting Cameron and Starr Counties, Texas [F75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France	Skylab water depth determination · · · Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during
instrument S-193 Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189	detecting Cameron and Starr Counties, Texas [E75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN
instrument S-193 ··· Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological studies from space ··· Nevada	detecting Cameron and Starr Counties, Texas [E75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0498 A75-17160 Skylab water depth determination Lake Erie and Lake
instrument S-193 Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological studies from space Nevada [E75-10088] p0038 N75-15106 INFRARED SCANNERS Some features of the urban environment of Tokyo by	detecting — Cameron and Starr Counties, Texas [E75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223 Contribution of space platforms to a ground and airborne	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160 Skylab water depth determination Lake Erie and Lake Michigan
instrument S-193 ··· Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological studies from space ··· Nevada [E75-10088] p0038 N75-15106 INFRARED SCANNERS Some features of the urban environment of Tokyo by remote sensing ··· infrared surface temperature	detecting Cameron and Starr Counties. Texas [E75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-1723 Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcances	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160 Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348
instrument S-193 ··· Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological studies from space ··· Nevada [E75-10088] p0038 N75-15106 INFRARED SCANNERS Some features of the urban environment of Tokyo by remote sensing ··· infrared surface temperature measurements p0012 A75-17102	detecting Cameron and Starr Counties. Texas [E75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223 Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcances	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160 Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 Determination of the earth's aerosol albedo using Skylab
instrument S-193 ··· Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological studies from space ··· Nevada [E75-10088] p0038 N75-15106 INFRARED SCANNERS Some features of the urban environment of Tokyo by remote sensing ··· infrared surface temperature measurements Remote sensing techniques applied to the study of Italian	detecting Cameron and Starr Counties. Texas [E75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0068 A75-17223 Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcanoes [E75-10069] Evaluation of ERTS-1 data by analogue and digital techniques Germany and Italy p0059 N75-14218	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160 Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348
instrument S-193 ··· Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological studies from space ··· Nevada [E75-10088] p0038 N75-15106 INFRARED SCANNERS Some features of the urban environment of Tokyo by remote sensing ··· infrared surface temperature measurements p0012 A75-17102	detecting Cameron and Starr Counties, Texas [E75-10109] pO010 N75-15125 ITALY Agricultural resources investigations in Southern France and Northern Italy Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223 Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcanoes [E75-10069] p0037 N75-14188 Evaluation of ERTS-1 data by analogue and digital techniques Germany and Italy p0059 N75-14218 Digital land-use classification using an ERTS-1 image	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160 Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 Determination of the earth's aerosol albedo using Skylab data Lake Michigan [E75-10096] p0025 N75-15114 LAKE ONTARIO
instrument S-193 ··· Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological studies from space ··· Nevada [E75-10088] p0038 N75-15106 INFRARED SCANNERS Some features of the urban environment of Tokyo by remote sensing ··· infrared surface temperature measurements p0012 A75-17102 Remote sensing techniques applied to the study of Italian volcanic areas · The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223	detecting Cameron and Starr Counties. Texas [E75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223 Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcanoes [E75-10069] p0037 N75-14188 Evaluation of ERTS-1 data by analogue and digital techniques Germany and Italy Digital land-use classification using an ERTS-1 image roccession of the control of the	Skylab water depth determination ··· Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160 Skylab water depth determination ··· Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 Determination of the earth's aerosol albedo using Skylab data ··· Lake Michigan [E75-10096] p0025 N75-15114 LAKE ONTARIO Lake Ontario water mass delineation from ERTS-1
instrument S-193 Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological studies from space Nevada [E75-10088] p0038 N75-15106 INFRARED SCANNERS Some features of the urban environment of Tokyo by remote sensing infrared surface temperature measurements p0012 A75-17102 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p00069 A75-17223 INFRARED SPECTRA	detecting Cameron and Starr Counties, Texas [E75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223 Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcanoes [E75-10069] Evaluation of ERTS-1 data by analogue and digital techniques Germany and Italy p0059 N75-14218 Digital land-use classification using an ERTS-1 image p0023 N75-14226 Colour composites of ERTS multispectral imagery for	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160 Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 Determination of the earth's aerosol albedo using Skylab data Lake Michigan [E75-1096] p0025 N75-15114 LAKE ONTARIO Lake Ontario water mass delineation from ERTS-1 p0042 A75-17184
instrument S-193 Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological studies from space Nevada [E75-10088] p0038 N75-15106 INFRARED SCANNERS Some features of the urban environment of Tokyo by remote sensing infrared surface temperature measurements p0012 A75-17102 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223 INFRARED SPECTRA Remote sensing of rock type in the visible and	detecting — Cameron and Starr Counties, Texas [E75-10109] ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-1723 Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcanoes [E75-10069] Evaluation of ERTS-1 data by analogue and digital techniques — Germany and Italy Digital land-use classification using an ERTS-1 image Italy Colour composites of ERTS multispectral imagery for applications in hydrology and oceanography — Italian	Skylab water depth determination ··· Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160 Skylab water depth determination ··· Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 Determination of the earth's aerosol albedo using Skylab data ··· Lake Michigan [E75-10096] p0025 N75-15114 LAKE ONTARIO Lake Ontario water mass delineation from ERTS-1 p042 A75-17184 Assessment of ERTS-1 imagery as a tool for regional
instrument S-193 ··· Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological studies from space ··· Nevada [E75-10088] p0038 N75-15106 INFRARED SCANNERS Some features of the urban environment of Tokyo by remote sensing ··· infrared surface temperature measurements p0012 A75-17102 Remote sensing techniques applied to the study of Italian volcanic areas · The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223 INFRARED SPECTRA Remote sensing of rock type in the visible and near-infrared p0033 A75-17220	detecting Cameron and Starr Counties. Texas [E75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-1723 Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcanoes [E75-10069] p0037 N75-14188 Evaluation of ERTS-1 data by analogue and digital techniques Germany and Italy p0059 N75-14218 Digital land-use classification using an ERTS-1 image p0023 N75-1425 Colour composites of ERTS multispectral imagery for applications in hydrology and oceanography Italian Mediterranean and North Sea areas p0045 N75-1421ian	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160 Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 Determination of the earth's aerosol albedo using Skylab data Lake Michigan [E75-10096] p0025 N75-15114 LAKE ONTARIO Lake Ontario water mass delineation from ERTS-1 p0042 A75-17184 Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State Lake Ontario
instrument S-193 Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological studies from space Nevada [E75-10088] p0038 N75-15106 INFRARED SCANNERS Some features of the urban environment of Tokyo by remote sensing infrared surface temperature p0012 A75-17102 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223 INFRARED SPECTRA Remote sensing of rock type in the visible and near-infrared Feasibility of using S-191 infrared spectra for geological studies and reserving specific processions.	detecting — Cameron and Starr Counties, Texas [E75-10109] ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-1723 Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcanoes [E75-10069] Evaluation of ERTS-1 data by analogue and digital techniques — Germany and Italy Digital land-use classification using an ERTS-1 image Italy Colour composites of ERTS multispectral imagery for applications in hydrology and oceanography — Italian	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160 Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 Determination of the earth's aerosol albedo using Skylab data Lake Michigan [E75-10096] p0025 N75-15114 LAKE ONTARIO Lake Ontario water mass delineation from ERTS-1 p0042 A75-17184 Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State Lake Ontario p0034 N75-10536
instrument S-193 ··· Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological studies from space ··· Nevada [E75-10088] p0038 N75-15106 INFRARED SCANNERS Some features of the urban environment of Tokyo by remote sensing ··· infrared surface temperature measurements p0012 A75-17102 Remote sensing techniques applied to the study of Italian volcanic areas · The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223 INFRARED SPECTRA Remote sensing of rock type in the visible and near-infrared p0033 A75-17220	detecting Cameron and Starr Counties. Texas [E75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0068 A75-17223 Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcanoes [E75-10069] p0037 N75-14188 Evaluation of ERTS-1 data by analogue and digital techniques Germany and Italy p0059 N75-14218 Digital land-use classification using an ERTS-1 image p0023 N75-14226 Colour composites of ERTS multispectral imagery for applications in hydrology and oceanography Italian Mediterranean and North Sea areas p0045 N75-14231 Geological features on ERTS-1 images of a test area in the west-central Alps Italy p0037 N75-14237 Geologic evaluation of central Italy from ERTS-1 and	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160 Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 Determination of the earth's aerosol albedo using Skylab data Lake Michigan [E75-10096] p0025 N75-15114 LAKE ONTARIO Lake Ontario water mass delineation from ERTS-1 p0042 A75-17184 Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State Lake Ontario [E74-10809] p0034 N75-10536 Dynamics of suspended sediment plumes in Lake Ontario
instrument S-193 ··· Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological studies from space ··· Nevada [E75-10088] p0038 N75-15106 INFRARED SCANNERS Some features of the urban environment of Tokyo by remote sensing ··· infrared surface temperature p0012 A75-17102 Remote sensing techniques applied to the study of Italian volcanic areas · The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223 INFRARED SPECTRA Remote sensing of rock type in the visible and pear-infrared p0033 A75-17220 Feasibility of using S-191 infrared spectra for geological studies from space ··· Nevada [E75-10089] p0038 N75-15107 INFRARED SPECTROMETERS	detecting Cameron and Starr Counties, Texas [E75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223 Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcanoes [E75-10069] p0037 N75-14188 Evaluation of ERTS-1 data by analogue and digital techniques Germany and Italy p0059 N75-14218 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Colour composites of ERTS multispectral imagery for applications in hydrology and oceanography Italian Mediterranean and North Sea areas p0045 N75-14231 Geological features on ERTS-1 images of a test area in the west-central Alps Italy p0037 N75-14237 Geological features on ERTS-1 images of a test area in the west-central Alps Italy p0037 N75-14230	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160 Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 Determination of the earth's aerosol albedo using Skylab data Lake Michigan [E75-10096] p0025 N75-15114 LAKE ONTARIO Lake Ontario water mass delineation from ERTS-1 p0042 A75-17184 Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State Lake Ontario [E74-10809] p0034 N75-10536 Dynamics of suspended sediment plumes in Lake Ontario [E74-10815] p0049 N75-10541
instrument S-193 Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological studies from space Nevada [E75-10088] p0038 N75-15106 INFRARED SCANNERS Some features of the urban environment of Tokyo by remote sensing infrared surface temperature measurements — infrared surface temperature measurements — p0012 A75-17102 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223 INFRARED SPECTRA Remote sensing of rock type in the visible and near-infrared p0033 A75-17220 Feasibility of using S-191 infrared spectra for geological studies from space Nevada [E75-10089] p0038 N75-15107 INFRARED SPECTROMETERS An orbiting visible/infrared spectrometer for terrestrial.	detecting Cameron and Starr Counties, Texas [E75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-1723 Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcanoes [E75-10069] p0037 N75-14188 Evaluation of ERTS-1 data by analogue and digital techniques Germany and Italy p0059 N75-14218 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14228 Colour composites of ERTS multispectral imagery for applications in hydrology and oceanography Italian Mediterranean and North Sea areas p0045 N75-14231 Geological features on ERTS-1 images of a test area in the west-central Alps Italy p0037 N75-14237 Geologic evaluation of central Italy from ERTS-1 and Skylab data p008 N75-14238 Agricultural resources investigations in northern Italy.	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160 Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 Determination of the earth's aerosol albedo using Skylab data Lake Michigan [E75-10096] p0025 N75-15114 LAKE ONTARIO Lake Ontario water mass delineation from ERTS-1 p0042 A75-17184 Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State Lake Ontario [E74-10809] p0034 N75-10536 Dynamics of suspended sediment plumes in Lake Ontario [E74-10815] p0049 N75-10541 A Skylab program for the International Hydrological
instrument S-193 ··· Utah and Texas Rad/Scat measurements [E75-10070] p0059 N75-14189 Feasibility of using S-191 infrared spectra for geological studies from space ··· Nevada [E75-10088] p0038 N75-15106 INFRARED SCANNERS Some features of the urban environment of Tokyo by remote sensing ··· infrared surface temperature p0012 A75-17102 Remote sensing techniques applied to the study of Italian volcanic areas · The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223 INFRARED SPECTRA Remote sensing of rock type in the visible and pear-infrared p0033 A75-17220 Feasibility of using S-191 infrared spectra for geological studies from space ··· Nevada [E75-10089] p0038 N75-15107 INFRARED SPECTROMETERS	detecting Cameron and Starr Counties, Texas [E75-10109] p0010 N75-15125 ITALY Agricultural resources investigations in Southern France and Northern Italy p0004 A75-17173 Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223 Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcanoes [E75-10069] p0037 N75-14188 Evaluation of ERTS-1 data by analogue and digital techniques Germany and Italy p0059 N75-14218 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Colour composites of ERTS multispectral imagery for applications in hydrology and oceanography Italian Mediterranean and North Sea areas p0045 N75-14231 Geological features on ERTS-1 images of a test area in the west-central Alps Italy p0037 N75-14237 Geological features on ERTS-1 images of a test area in the west-central Alps Italy p0037 N75-14230	Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 LAKE MICHIGAN Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160 Skylab water depth determination Lake Erie and Lake Michigan [E75-10048] p0050 N75-13348 Determination of the earth's aerosol albedo using Skylab data Lake Michigan [E75-10096] p0025 N75-15114 LAKE ONTARIO Lake Ontario water mass delineation from ERTS-1 p0042 A75-17184 Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State Lake Ontario [E74-10809] p0034 N75-10536 Dynamics of suspended sediment plumes in Lake Ontario [E74-10815] p0049 N75-10541

SUBJECT INDEX LASERS

A Skylab program for the International Hydrological Decade (IHD) Lake Ontario Basin [E75-10093] p0052 N75-15111	The application of ERTS-derived information to the	
[E75-10093] p0052 N75-15111		Evaluation of Skylab imagery as an information service
(2.0 .000)	regional land use planning process p0015 A75-18058 Land use classification with ERTS-1 data in the Houston	for investigating land use and natural resources New
		York
A Skylab program for the International Hydrological	area test site p0015 A75-18059	[NASA-CR-141372] p0024 N75-15100
Decade (IHD) Lake Ontario Basin	Geometronics-earth resource measurements and displays	Investigation of the environmental change pattern of
[E75-10094] p0052 N75-15112	for the Forest Service p0005 A75-18063	Japan
LAKE SUPERIOR	Land use mapping using ERTS-1 imagery for The Four	[E75-10086] p0024 N75-15104
Remote sensing of western Lake Superior	Rivers Resource Conservation and Development District in	Urban and regional land use analysis: CARETS and
p0048 A75-17161 a	Kansas p0016 A75-18070	census cities experiment package New ' Haven,
	Evaluation of Skylab imagery as an information service	Connecticut: Pontiac, Michigan: Phoenix, Arizona: District
A study of Minnesota forests and lakes using data from	for investigating land use and natural resources Long	of Columbia
Earth Resources Technology Satellites	Island, New York	[E75-10091] p0024 N75-15109
[NASA-CR-140865] p0007 N75-12421	[E74-10730] p0058 N75-10526	
LAKES		Study of recreational land and open space using Skylab
The use of remote sensing in limnological studies	ERTS-1 imagery interpretation techniques in the	imagery snow and ice hydrology of southeast
eutrophication of lakes p0048 A75-17163	Tennessee Valley land use and soil mapping	Michigan
Automatic classification of eutrophication of inland lakes	[E74-10811] p0016 N75-10538	[E75-10092] p0024 N75-15110
from spacecraft data p0048 A75-17164	Investigation of land use of northern megalopolis using	Urban and regional land use analysis: CARETS and
	ERTS-1 imagery	census cities experiment package
Monitoring trophic status of inland lakes - A quantitative	[NASA-CR-2459] p0017 N75-10545	[E75-10100] p0025 N75-15118
application of ERTS imagery p0049 A75-18067	Land use and mapping p0017 N75-10560	
A study of Minnesota forests and lakes using data from		Multidisciplinary analysis for highway engineering
Earth Resources Technology Satellites	Earth Resources Technology Satellite-1 (ERTS-1) data	purposes Maine
[NASA-CR-140865] p0007 N75-12421	and anthropology: Use of these data in carrying capacity	[E75-10103] p0025 N75-15121
Skylab study of water quality Kansas reservoirs and	estimates for sites in Upper Volta and Niger	LANDFORMS
	[E75-10001] p0078 N75-11406	Mineral resources, geological structures, and landform
lakes -0051 NZE 14194	Evaluation of Skylab imagery as an information service	surveys p0035 N75-10561
[E75-10075] p0051 N75-14194	for investigating land use and natural resources	Applications of Skylab EREP photographs to mapping
LAND MANAGEMENT	[E75-10018] p0020 N75-12397	
ERTS-I data for classifying native plant communities -		of landforms and environmental geology in the Great Plains
Central Colorado p0004 A75-17176	Urban and regional land use analysis: CARETS and	and Midwest Illinois, Iowa, Kansas, Missouri, Nebraska,
Resource analysis applications in Michigan NASA	census cities experiment package	and South Dakota
remote sensing p0069 A75-17231	[E75-10019] p0020 N75-12398	[E75-10055] p0037 N75-13355
	Study of recreational land and open space using Skylab	LANDSAT SATELLITES
Application of ERTS-1 data to the protection and	imagery	Forest defoliation assessment with satellite imagery
management of New Jersey's coastal environment	[E75-10020] p0020 N75-12399	p0003 A75-17171
p0042 A75-17233	Study of recreational land and open space using Skylab	A summary of ERTS data applications in Alaska
Application of ERTS data to environmental inventory and		
assessment p0016 A75-18069	imagery southeast Michigan	p0077 A75-17236
	[E75-10034] p0020 N75-12404	The economic value of remote sensing of earth resources
	Detection of moisture and moisture related phenomena	from space: An ERTS overview and the value of continuity
The role of ERTS in the establishment and of a nationwide	from Skylab Kansas, Oklahoma, and Arkansas	of service. Volume 3: Intensive use of living resources.
land cover information system	[E75-10035] p0007 N75-12405	agriculture. Part 2: Distribution effects
[NASA-CR-141045] p0020 N75-12425	The ERTS-1 investigation. Volume 7: Erts-1 land-use	[NASA-CR-141194] p0079 N75-14206
Evaluation of Skylab imagery as an information service		The economic value of remote sensing of earth resources
for investigating land use and natural resources New	analysis of the Houston area test site	
York	[E75-10009] p0021 N75-13334	from space: An ERTS overview and the value of continuity
[E75-10068] p0059 N75-14187	Use of feature extraction techniques for the texture and	of service. Volume 9: Oceans
The application of Skylab EREP data for land use	context information in ERTS imagery: Spectral and textural	[NASA-CR-141270] p0080 N75-14214
	processing of ERTS imagery classification of Kansas	
classification and mapping in the Cleveland and Columbus	land use	LANDSAT 1
areas	[E75-10011] p0059 N75-13335	Skylab and ERTS-1 investigations of coastal land-use
[E75-10083] p0023 N75-14201	- · · · · · · · · · · · · · · · · · · ·	and water properties in Delaware Bay
Evaluation of Skylab imagery as an information service	Evaluation of Skylab imagery as an information service	[AIAA PAPER 74-1220] p0041 A75-12034
for investigating land use and natural resources New	for investigating land use and natural resources Hudson	
York	River Valley and Long Island, New York	Evaluation of ERTS-1 and aircraft data for assessing
[NASA-CR-141372] p0024 N75-15100	[E75-10013]. p0059 N75-13336	internal drainage in irrigated agriculture
Usefulness of Skylab color photography and ERTS-1	Urban and regional land use analysis: CARETS and	p0002 A75-17143
multispectral imagery for mapping range vegetation types	census cities experiment package	ERTS-I data for classifying native plant communities -
in southwestern Wyoming	[E75-10056] p0022 N75-13356	Central Colorado p0004 A75-17176
[E75-10097] p0010 N75-15115	Utilization of EREP data in geological evaluation regional	ERTS 1 flight evaluation report, 23 October 1973 - 23
LAND USE		January 1974 orbits 6371 to 7651
Skylab and ERTS-1 investigations of coastal land-use	planning, forest management, and water management in	
	North Carolina	[NASA-CR-140712] p0077 N75-10546
and water properties in Delaware Bay [AIAA PAPER 74-1220] p0041 A75-12034	[E75-10063] p0008 N75-13363	Correlation and registration of ERTS multispectral
	Computer and photogrammetric general land use study	imagery by a digital processing technique
Automatic land use classification using Skylab S-192	of central north Alabama	[NASA-CR-134294] p0058 N75-12420
	[NASA-TR-R-431] p0022 N75-13372	The ERTS-1 investigation (ER-600). Volume 2: ERTS-1
multispectral data		
[AIAA PAPER 74-1224] p0064 A75-12036		
[AIAA PAPER 74-1224] p0064 A75-12036 Agricultural inventory techniques with orbital and	Evaluation of Skylab imagery as an information service	coastal/estuarine analysis Galveston Bay, Texas
[AIAA PAPER 74-1224] p0064 A75-12036	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371
[AIAA PAPER 74-1224] p0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image
[AIAA PAPER 74-1224] p0064 A75-12036 Agricultural inventory techniques with orbital and	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] p0059 N75-14187	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226
[AIAA PAPER 74-1224] p0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 ERTS-1 and data for national land use planning p0075 A75-13581	Evaluation of Skylab imagery as an information service for investigating land use and natural resources ··· New York [E75-10068] p0059 N75-14187 Develop techniques and procedures, using multispectral	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] p0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226
[AIAA PAPER 74-1224] p0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 ERTS-1 and data for national land use planning p0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p0047 A75-16240	Evaluation of Skylab imagery as an information service for investigating land use and natural resources ··· New York [E75-10068] p0059 N75-14187 Develop techniques and procedures, using multispectral	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 imagery of the
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] p0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] p0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p0008 N75-14196	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image p0023 N75-14226 Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] p0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p0008 N75-14196 The application of Skylab EREP data for land use	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing.	Evaluation of Skylab imagery as an information service for investigating land use and natural resources ··· New York [E75-10068] p0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p0008 N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image p0023 N75-14226 Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] p.0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p.0008 N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0078 A75-17118	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] p.0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p.0008 N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] p.0023 N75-14201	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] DOOB N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] DOO23 N75-14201 The economic value of remote sensing of earth resources	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] p.0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p.0008 N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] p.0023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource	Evaluation of Skylab imagery as an information service for investigating land use and natural resources ··· New York [E75-10068] p.0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p.0008 N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] p.0023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] p.0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p.0008 N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] p.0023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p0008 N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] p0080 N75-14208	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0078 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] p.0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p.0008 N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] p.0023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] p.0080 N75-14208 The economic value of remote sensing of earth resources	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 LASER DOPPLER VELOCIMETERS
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] D0023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service.	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERT5-1 image Italy p0023 N75-14226 Pedological interpretation of ERT5-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] p0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p0008 N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] p0023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydriogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p.0013 A75-17136 Land type analysis for regional land use planning from	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] DOOS N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] DOO23 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] December 120 Nasa-CR-141263 December 120 Nasa-CR-14	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 image y of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] p0021 N75-12462
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County.	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p0008 N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] p0080 N75-14210	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERT5-1 image Italy p0023 N75-14226 Pedological interpretation of ERT5-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] LASER RANGE FINDERS
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0078 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p.0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] p0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p0008 N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] p0023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] p0080 N75-14208 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 8: Land use. Part 1: Introduction and overview [NASA-CR-141266] p0080 N75-14210 The economic value of remote sensing of earth resources	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image p0023 N75-14226 Pedological interpretation of ERTS-1 image po023 N75-14226 Pedological interpretation of ERTS-1 image yo of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] p0021 N75-12462 LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0076 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use panalysis p.0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado p.0013 A75-17137 Implementation of an advanced table look-up classifier	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] DO023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] DO080 N75-14210 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] DO080 N75-14210 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service.	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERT5-1 image Italy p0023 N75-14226 Pedological interpretation of ERT5-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] p0021 N75-12462 LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data Design concepts for land use and natural resource inventories and information systems p.0013 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p.0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado p.0013 A75-17137 Implementation of an advanced table look-up classifier for large area land-use classification p.0014 A75-17153	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] p0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p0008 N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] p0023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] p0080 N75-14208 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 8: Land use. Part 1: Introduction and overview [NASA-CR-141266] p0080 N75-14210 The economic value of remote sensing of earth resources	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills Development of an experimental siborne system to the measurement of atmospheric winds [NASA-TM-X-64891] Development of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] Development of a single laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-14648
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado p.0013 A75-17137 Implementation of an advanced table look-up classifier for large area land-use classification of p.0014 A75-17153 Automatic classification of eutrophication of inland lakes	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] DO023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] DO080 N75-14210 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] DO080 N75-14210 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service.	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 image y of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] p0021 N75-12462 LASER RANGE FINDERS Geodatic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-14648 Possibilities of calculating the earth's local gravitational
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado Implementation of an advanced table look-up classifier for large area land-use classification p.0014 A75-17153 Automatic classification of eutrophication of inland lakes from spacecraft data	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] Develop The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] Develop The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] Develop The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] DO080 N75-14210 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of service. An ERTS overview and the value of continuity of service. An ERTS overview and the value of continuity of service. An ERTS overview and the value of continuity of service. An ERTS overview and the value of continuity of service. Noteme 6: Land use. Part 2: The role of service. Volume 6: Land use. Part 2: The role of	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERT5-1 image Italy p0023 N75-14226 Pedological interpretation of ERT5-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-14646 Possibilities of calculating the earth solical gravitational field from satellite observations p0029 A75-18524
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0078 A75-17118 Design concepts for land use and natural resources inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p.0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado table look-up classifier for large area land-use classification p.0013 A75-17133 Automatic classification of eutrophication of inland lakes from spacecraft data p.0048 A75-17164 Correlation of multispectral imagery with water analysis	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a nationwide land cover information system	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 image y of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] p0021 N75-12462 LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-14648 Possibilities of calculating the earth's local gravitational field from satellite observations p0029 A75-18524 The role of lidar in environmental sensing Isser
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado Implementation of an advanced table look-up classifier for large area land-use classification p.0014 A75-17153 Automatic classification of eutrophication of inland lakes from spacecraft data	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] p0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p0008 N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] p0023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] p0080 N75-14208 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] p0080 N75-14210 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of CRTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141267] p0080 N75-14211	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERT5-1 image Italy p0023 N75-14226 Pedological interpretation of ERT5-1 image yof the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] p0021 N75-12462 LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-14646 Possibilities of calculating the earth's local gravitational field from satellite observations p0029 A75-18524 The role of lidar in environmental sensing leser ceilometer and visual range measurement application
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0078 A75-17118 Design concepts for land use and natural resources inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p.0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado table look-up classifier for large area land-use classification p.0013 A75-17133 Automatic classification of eutrophication of inland lakes from spacecraft data p.0048 A75-17164 Correlation of multispectral imagery with water analysis	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] DO023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] DO080 N75-14210 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141267] DO080 N75-14211 Digital land-use classification using an ERTS-1 image	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 image y of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] p0021 N75-12462 LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-14648 Possibilities of calculating the earth's local gravitational field from satellite observations p0029 A75-18524 The role of lidar in environmental sensing Isser
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data Design concepts for land use and natural resource inventories and information systems p.0013 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p.0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado p.0013 A75-17137 Implementation of an advanced table look-up classifier for large area land-use classification p.0014 A75-17153 Automatic classification of eutrophication of inland lakes from spacecraft data p.0048 A75-17164 Correlation of multispectral imagery with water analysis - The Ross Barnett Reservoir Remote Sensing Project	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of RTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141267] Digital land-use classification using an ERTS-1 image Italy PO023 N75-14226	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERT5-1 image Italy p0023 N75-14226 Pedological interpretation of ERT5-1 image yof the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] p0021 N75-12462 LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-14646 Possibilities of calculating the earth's local gravitational field from satellite observations p0029 A75-18524 The role of lidar in environmental sensing leser ceilometer and visual range measurement application
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p.0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado p.0013 A75-17137 Implementation of an advanced table look-up classifier for large area land-use classification p.0014 A75-17155 Automatic classification of eutrophication of inland lakes from spacecraft data p.0048 A75-17165 Inventories of Delaware's coastal vegetation and land-use	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] DO023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] DO080 N75-14210 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141267] DO080 N75-14211 Digital land-use classification using an ERTS-1 image	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERT5-1 image Italy p0023 N75-14226 Pedological interpretation of ERT5-1 image y of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills Dougload Sample Sampl
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p.0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado p.0013 A75-17131 Implementation of an advanced table look-up classifier for large area land-use classification p.0014 A75-17153 Automatic classification of eutrophication of inland lakes from spacecraft data p.004 A75-17164 Correlation of multispectral imagery with water analysis - The Ross Barnett Reservoir Remote Sensing Project p.0014 A75-17165 Inventories of Delaware's coastal vegetation and land-use utilizing digital processing of ERTS-1 imagery	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of RTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141267] Digital land-use classification using an ERTS-1 image Italy PO023 N75-14226	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERT5-1 image Italy p0023 N75-14226 Pedological interpretation of ERT5-1 image yof the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-14646 Possibilities of calculating the earth socal gravitational field from satellite observations p0029 A75-18524 The role of lidar in environmental sensing laser ceilometer and visual range measurement application p0070 A75-19274 LASER RANGER/TRACKER Satellite techniques for the study of secular variations
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data Design concepts for land use and natural resource inventories and information systems p.0013 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p.0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado p.0013 A75-17137 Implementation of an advanced table look-up classifier for large area land-use classification p.0014 A75-17153 Automatic classification of eutrophication of inland lakes from spacecraft data p.0048 A75-17166 Inventories of Delaware's coastal vegetation and land-use utilizing digital processing of ERTS-1 imagery p.0004 A75-17186	Evaluation of Skylab imagery as an information service for investigating land use and natural resources ··· New York [E75-10068] p0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p0008 N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] p0023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] p0080 N75-14208 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] p0080 N75-14210 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of CRTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141267] p0080 N75-14211 Digital land-use classification using an ERTS-1 image p0023 N75-14226 A comparison of ERTS imagery with conventional aerial	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 image y of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] p0021 N75-12462 LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-14646 Possibilities of calculating the earth's local gravitational field from satellite observations p0029 A75-18524 The role of lidar in environmental sensing Isser ceilometer and visual range measurement application p0070 A75-19274 LASER RANGER/TRACKER Satellite techniques for the study of secular variations in position p0064 A75-12380
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p.0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado Implementation of an advanced table look-up classifier for large area land-use classification p.0014 A75-17153 Automatic classification of eutrophication of inland lakes from spacecraft data p.0048 A75-17164 Correlation of multispectral imagery with water analysis The Ross Barnett Reservoir Remote Sensing Project p.0014 A75-17166 Inventories of Delaware's coastal vegetation and land-use utilizing digital processing of ERTS-1 imagery p.0004 A75-17166 Land use classification and ground truth correlations from	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] p0023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a natiomwide land cover information system [NASA-CR-141267] Digital land-use classification using an ERTS-1 image Italy A comparison of ERTS imagery with conventional aerial photography for land-resource surveys in less developed	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERT5-1 image Italy p0023 N75-14226 Pedological interpretation of ERT5-1 image yof the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATION8 Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-14646 Possibilities of calculating the earth's local gravitational field from satellite observations p0029 A75-18524 The role of lidar in environmental sensing leser ceilometer and visual range measurement application p0070 A75-19274 LASER RANGER/TRACKER Satellite techniques for the study of secular variations in position p0064 A75-12980 LASERS
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data Design concepts for land use and natural resource inventories and information systems p.0013 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p.0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado p.0013 A75-17137 Implementation of an advanced tabble look-up classifier for large area land-use classification p.0014 A75-17153 Automatic classification of eutrophication of inland lakes from spacecraft data p.0048 A75-17164 Correlation of multispectral imagery with water analysis - The Ross Barnett Reservoir Remote Sensing Project p.0014 A75-17165 Inventories of Delaware's coastal vegetation and land-use utilizing digital processing of ERTS-1 imagery p.0004 A75-17186 Land use classification and ground truth correlations from simultaneously acquired aircraft and ERTS-1 MSS data	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] DO023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a natiomwide land cover information system [NASA-CR-141267] Digital land-use classification using an ERTS-1 image processing of earth resources with conventional aerial photography for land-resource surveys in less developed countries. Examples from the Rift Valley Lakes Basin, poos 1 p	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image relative policy
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-1718 Design concepts for land use and natural resource inventories and information systems p.0076 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use palanning p.0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado p.0013 A75-17136 Automatic classification of eutrophication of inland lakes from spacecraft data p.0048 A75-17166 Correlation of multispectral imagery with water analysis the Ross Barnett Reservoir Remote Sensing Project p.0014 A75-17165 Inventories of Delaware's coastal vegetation and land-use utilizing digital processing of ERTS-1 imagery p.004 A75-17186 Land use classification and ground truth correlations from simultaneously acquired aircraft and ERTS-1 MSS data p.0014 A75-17188	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] Down N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141267] Digital land-use classification using an ERTS-1 image	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERT5-1 image Italy p0023 N75-14226 Pedological interpretation of ERT5-1 image y of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] p0021 N75-12462 LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-14648 Possibilities of calculating the earth's local gravitational field from satellite observations p0029 A75-18524 The role of lidar in environmental sensing laser ceilometer and visual range measurement application p0070 A75-19274 LASER RANGER/TRACKER Satellite techniques for the study of secular variations in position LASERS Soviet satellite geodesy: Basis and status of Soviet laser geodesy and long-base line interferometry
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p.0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado p.0013 A75-17135 Implementation of an advanced table look-up classifier for large area land-use classification p.0014 A75-17153 Automatic classification of eutrophication of inland lakes from spacecraft data p.004 A75-17166 Correlation of multispectral imagery with water analysis - The Ross Barnett Reservoir Remote Sensing Project p.0014 A75-17166 Inventories of Delaware's coastal vegetation and land-use utilizing digital processing of ERTS-1 imagery with water for small processing of ERTS-1 imagery of the p.004 A75-17186 Land use classification and ground truth correlations from simultaneously acquired aircraft and ERTS-1 MSS data p.0014 A75-17188 Resource analysis applications in Michigan	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] p0059 N75-14187 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p0008 N75-14196 The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] p0023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] p0080 N75-14203 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview (NASA-CR-141266) p0080 N75-14210 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview (NASA-CR-141266) p0080 N75-14210 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141267] p0080 N75-14211 Digital land-use classification using an ERTS-1 image wold information system p0023 N75-1426 A comparison of ERTS imagery with conventional aerial photography for land-resource surveys in less developed countries. Examples from the Rift Valley Lakes Basin, Ethiopia Future US activities based on experience gained from ERTS-1 and EREP	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image p-0023 N75-14226 Pedological interpretation of ERTS-1 image pool of Pedological interpretation of ERTS-1 image pool of N75-14243 Pedological interpretation of Language pool of N75-14243 Digital-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17208 LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] p0021 N75-12462 LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-14648 Possibilities of calculating the earth's local gravitational field from satellite observations p0029 A75-18524 The role of lidar in environmental sensing laser ceilometer and visual range measurement application p0074 A75-19274 LASER RANGER/TRACKER Satellite techniques for the study of secular variations in position p0064 A75-12980 LASERS Soviet satellite geodesy: Basis and status of Soviet laser geodesy and long-base line interferometry [AD-781138/3] p0029 N75-10524
[AIAA PAPER 74-1224] p0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 ERTS-1 and data for national land use planning p0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p0007 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p0013 A75-17133 Interactive computer processing for land use planning p0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado p0013 A75-17175 Implementation of an advanced table look-up classifier for large area land-use classification of p004 A75-17165 Automatic classification of eutrophication of inland lakes from spacecraft data p0048 A75-17165 Correlation of multispectral imagery with water analysis The Ross Barnett Reservoir Remote Sensing Project p004 A75-17165 Inventories of Delaware's coastal vegetation and land-use utilizing digital processing of ERTS-1 imagery p004 A75-17186 Land use classification and ground truth correlations from simultaneously acquired aircraft and ERTS-1 MSS data p0014 A75-17188 Resource analysis applications in Michigan NASA remote sensing p0069 A75-17231	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] DO023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] DO080 N75-14210 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of an antiomwide land cover information system [NASA-CR-141267] Digital land-use classification using an ERTS-1 image Italy A comparison of ERTS imagery with conventional serial photography for land-resource surveys in less developed countries. Examples from the Rift Valley Lakes Basin, Ethiopia FULLIP Italy	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 image y of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] p0021 N75-12462 LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-14646 Possibilities of calculating the earth's local gravitational field from satellite observations p0029 A75-18524 The role of lidar in environmental sensing Isser ceilometer and visual range measurement application p0070 A75-19274 LASER RANGER/TRACKER Satellite techniques for the study of secular variations in position p0064 A75-12980 LASERS Soviet satellite geodesy: Basis and status of Soviet laser geodesy and long-base line interferometry [AD-781138/3] p0029 N75-10524 A study of a dual polarization laser backscatter system
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado Douglas A75-17135 Automatic classification of eutrophication of inland lakes from spacecraft data Correlation of multispectral imagery with water analysis - The Ross Barnett Reservoir Remote Sensing Project DOUA A75-17166 Inventories of Delaware's coastal vegetation and land-use utilizing digital processing of ERTS-1 imagery with water analysis - The Ross Barnett Reservoir Remote Sensing Project DOUA A75-17186 Land use classification and ground truth correlations from simultaneously acquired aircraft and ERTS-1 MSS data p.004 A75-17188 Resource analysis applications in Michigan	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141267] Digital tand-use classification using an ERTS-1 image in that the pool of ERTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141267] Digital tand-use classification using an ERTS-1 image ool of ERTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141267] Digital tand-use classification using an ERTS-1 image services the pool of ERTS imagery with conventional aerial photography for land-resource surveys in less developed countries. Examples from the Rift Valley Lakes Basin, Ethiopia POOS N75-14253 Work dealing with densification of altitimetric-planometric	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERT5-1 image Italy p0023 N75-14226 Pedological interpretation of ERT5-1 image; of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATION8 Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] p0021 N75-12462 LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-14646 Possibilities of calculating the earth's local gravitational field from satellite observations p0029 A75-18524 The role of lidar in environmental sensing leser ceilometer and visual range measurement application p0070 A75-19274 LASER RANGER/TRACKER Satellite techniques for the study of secular variations in position LASERS Soviet satellite geodesy: Basis and status of Soviet laser geodesy and long-base line interferometry [AD-781138/3] p0029 N75-10524 A study of a dual polarization laser backscatter system for remote identification and measurement of water
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data Design concepts for land use and natural resource inventories and information systems p.0013 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p.0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado p.0013 A75-17137 Implementation of an advanced table look-up classifier for large area land-use classification p.0014 A75-17153 Automatic classification of eutrophication of inland lakes from spacecraft data p.0048 A75-17164 Correlation of multispectral imagery with water analysis - The Ross Barnett Reservoir Remote Sensing Project p.0014 A75-17165 Inventories of Delaware's coastal vegetation and land-use utilizing digital processing of ERTS-1 imagery p.0004 A75-17186 Land use classification and ground truth correlations from simultaneously acquired aircraft and ERTS-1 MSS data p.0014 A75-17188 Resource analysis applications in Michigan	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] DO023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] DO080 N75-14210 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a natiomwide land cover information system [NASA-CR-141267] Digital land-use classification using an ERTS-1 image rully pools of the stablishment and updating of a natiomwide land cover information system [NASA-CR-141267] Digital land-use classification using an ERTS-1 image rully pools not pools of the stablishment and updating of a natiomwide land cover information system [NASA-CR-141267] DO080 N75-142167 Digital land-use classification using an ERTS-1 image rully pools not pools n	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 image y of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] p0021 N75-12462 LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-1468A Possibilities of calculating the earth's local gravitational field from satellite observations p0029 A75-18524 The role of lidar in environmental sensing laser ceilometer and visual range measurement application p0070 A75-19274 LASER RANGER/TRACKER Satellite techniques for the study of secular variations in position p0064 A75-12280 LASERS Soviet satellite geodesy: Basis and status of Soviet laser geodesy and long-base line interferometry [AD-781138/3] p0029 N75-10524 A study of a dual polarization laser backscatter system for remote identification and measurement of water pollution
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed synthetic aperture radar imagery p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data natural resources and information systems p.0076 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p.0013 A75-17134 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado p.0013 A75-17133 Implementation of an advanced table look-up classifier for large area land-use classification p.0014 A75-17134 Correlation of multispectral imagery with water analysis 7the Ross Barnett Reservoir Remote Sensing Project p.0014 A75-17165 Inventories of Delaware's coastal vegetation and land-use utilizing digital processing of ERTS-1 imagery p.0004 A75-17186 Land use classification and ground truth correlations from simultaneously acquired aircraft and ERTS-1 MSS data p.0014 A75-17188 Resource analysis applications in Michigan NASA remote sensing p.0069 A75-17231 An intensive natural resources inventory aids land use planning and resources management in Puerto Rico p.0005 A75-18056	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141267] Digital tand-use classification using an ERTS-1 image in that the pool of ERTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141267] Digital tand-use classification using an ERTS-1 image ool of ERTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141267] Digital tand-use classification using an ERTS-1 image services the pool of ERTS imagery with conventional aerial photography for land-resource surveys in less developed countries. Examples from the Rift Valley Lakes Basin, Ethiopia POOS N75-14253 Work dealing with densification of altitimetric-planometric	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERT5-1 image Italy p0023 N75-14226 Pedological interpretation of ERT5-1 image yof the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TH-74646 Possibilities of calculating the earth's local gravitational field from satellite observations p0027 A75-14646 Possibilities of calculating the earth's local gravitational field from satellite observations p0029 A75-18524 The role of lidar in environmental sensing loser ceilometer and visual range measurement application p0070 A75-19274 LASER RANGER/TRACKER Satellite techniques for the study of secular variations in position LASERS Soviet satellite geodesy: Basis and status of Soviet laser geodesy and long-base line interferometry [AD-781138/3] p0029 N75-10563 A study of a dual polarization laser backscatter system for remote identification and measurement of water pollution [NASA-CR-140665]
[AIAA PAPER 74-1224] p.0064 A75-12036 Agricultural inventory techniques with orbital and high-altitude imagery p.0001 A75-12872 ERTS-1 and data for national land use planning p.0075 A75-13581 Coastal and estuarine studies with ERTS-1 and Skylab p.0047 A75-16240 Extraction of urban land cover data from multiplexed p.0054 A75-17117 Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing, natural resources, social and economic data Design concepts for land use and natural resource inventories and information systems p.0013 A75-17118 Design concepts for land use and natural resource inventories and information systems p.0013 A75-17133 Interactive computer processing for land use planning p.0054 A75-17134 Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p.0013 A75-17136 Land type analysis for regional land use planning from photomorphic mapping - An example for Boulder County, Colorado p.0013 A75-17137 Implementation of an advanced table look-up classifier for large area land-use classification p.0014 A75-17153 Automatic classification of eutrophication of inland lakes from spacecraft data p.0048 A75-17164 Correlation of multispectral imagery with water analysis - The Ross Barnett Reservoir Remote Sensing Project p.0014 A75-17165 Inventories of Delaware's coastal vegetation and land-use utilizing digital processing of ERTS-1 imagery p.0004 A75-17186 Land use classification and ground truth correlations from simultaneously acquired aircraft and ERTS-1 MSS data p.0014 A75-17188 Resource analysis applications in Michigan	Evaluation of Skylab imagery as an information service for investigating land use and natural resources New York [E75-10068] Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] DO023 N75-14201 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction and overview [NASA-CR-141266] DO080 N75-14210 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a natiomwide land cover information system [NASA-CR-141267] Digital land-use classification using an ERTS-1 image rully pools of the stablishment and updating of a natiomwide land cover information system [NASA-CR-141267] Digital land-use classification using an ERTS-1 image rully pools not pools of the stablishment and updating of a natiomwide land cover information system [NASA-CR-141267] DO080 N75-142167 Digital land-use classification using an ERTS-1 image rully pools not pools n	coastal/estuarine analysis Galveston Bay, Texas [NASA-TM-X-58118] p0051 N75-13371 Digital land-use classification using an ERTS-1 image Italy p0023 N75-14226 Pedological interpretation of ERTS-1 image y of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 LASER APPLICATIONS Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 LASER DOPPLER VELOCIMETERS Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] p0021 N75-12462 LASER RANGE FINDERS Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-1468A Possibilities of calculating the earth's local gravitational field from satellite observations p0029 A75-18524 The role of lidar in environmental sensing laser ceilometer and visual range measurement application p0070 A75-19274 LASER RANGER/TRACKER Satellite techniques for the study of secular variations in position p0064 A75-12280 LASERS Soviet satellite geodesy: Basis and status of Soviet laser geodesy and long-base line interferometry [AD-781138/3] p0029 N75-10524 A study of a dual polarization laser backscatter system for remote identification and measurement of water pollution

LAW (JURISPRUDENCE) MAGNETIC MEASUREMENT The Coho Project: Living resources prediction feasibility Oil pollution detection, monitoring and law enforcement - Louisiana Gulf Coast waters Report of the geomagnetic and geoelectric observations, 1972 (rapid variations) · · · meteorological forecasting of fish study, volume 1 concentrations [PB-234057/8] [F75-10027] n0022 N75-13343 [REPT-13] n0032 N75-15228 00044 N75-10681 LIFE SCIENCES The Coho Project: Living resources prediction feasibility MAGNETIC SIGNATURES The Proceedings of the Skylab Life Sciences Symposium. Magnetopause rotational forms study. Volume 2: Environmental report p0011 A75-11221 volume 1 [NASA-TM-X-58154-VOL-1] p0044 N75-10682 [PB-234058/6] MAGNETIC STORMS p0081 N75-14358 Investigation of the environmental change pattern of On the longitudinal localization of the substorm active LIGHT (VISIBLE RADIATION) Janan region and its changes during the substorm [E75-10086] Remote sensing of rock type in in the visible and p0033 A75-17220 p0027 A75-11861 Geophysical variations at Sodankyla during a geomagnetic storm, 17-18 December 1971 [REPT-9] p0030 N75-13432 MARINE TECHNOLOGY near-infrared Analysis of methods for the geodetic support of complex sea geophysical surveys

[JPRS-63621] p0031 N75-13498 Radiative transfer of visible radiation in turbid p0019 N75-11552 atmosphere p0030 N75-13432 An examination of whiteouts and transmission through MAGNETIC TAPES ice fog in the visible and near infrared [DREO-TN-74-18] MARKET RESEARCH Mapping of water bodies in northern Germany from ERTS pes p0051 N75-14227 p0021 N75-12467 Improved (ERTS) information and its impact on U.S. tapes markets for agricultural commodities: A quantitiative economic investigation of production, distribution and net LIGHT SCATTERING Lectures on the scattering of light ... by dielectric pool N75-11556 MAGNETOPAUSE Magnetopause rotational forms p0011 A75-11221 enhere vnort effects LIMESTONE MAGNETOSPHERIC INSTABILITY NASA-CR-141044] Determination of points of entry for potential On the longitudinal localization of the substorm active MARSHLANDS Evaluation of satellite remote sensing and automatic data techniques for characterization of wetlands and coastal contaminants into limestone aquifers using thermal infrared region and its changes during the substo p0027 A75-11861 imagery [NASA-CR-120540] p0021 N75-12546 marshlands --- detection and measurement of accretion in estuary system of Atchafalaya River in southern Solar wind and geomagnetic activity LIMNOLOGY DO027 A75-11975 The use of remote sensing in timnological studies ---Louisiana MAINE phication of lakes p0048 A75-17163 [E75-10025] Multidisciplinary analysis for highway engineering purposes --- Maine [E75-10103] p0025 N75-15121 p0044 N75-13341 LITHOLOGY Evaluation of satellite remote sensing and automatic data techniques for characterization of wetlands and coastal marshlands --- Atchafalaya River Basin, Louisiana [E75-10028] p0050 N75-13344 The Great Basin investigation [E75-10076] p0037 N75-14195 MALAGASY REPUBLIC The Great Basin investigation --- photomapping lithology ERTS image contribution to geological knowledge and geological structures (E75-10102) The use of color infrared imagery for the study of marsh iggy tracks p0061 N75-15127 covering different types of climates (France, Togo, Dahomey, Madagascar) p0038 N75-14240 p0038 N75-15120 buggy tracks Madagascar) LLANOS ORIENTALES (COLOMBIA) MARYLAND Agricultural resources investigations in northern Italy. ERTS-1 contributions to the ecoger ical knowledge of Urban and regional land use analysis: CARETS and southern France and Madagascar --- ERTS-B and aerial rice growing and forestry surveys p0009 N75-14252 the Llanos of the Orinoco (Colombia and Venezuela) census cities experiment package p0023 N75-14245 [£75-10019] p0020 N75-12398 MAN MACHINE SYSTEMS Remote sensing geophysics from Skylab --- rationing-type LONG ISLAND (NY) Interactive computer processing for land use planning p0054 A75-17134 Evaluation of Skylab imagery as an information service for investigating land use and natural resources --- Long algorithms to enhance multispectral data algorithms to employed process and process and Urban and regional land use analysis: CARETS and Urban and regional land use analysis: CARETS and New Haven, Petrick Interactive machine assessment of critical land resources using ERTS-1 data p0013 A75-17149 Island, New York census cities experiment package --- New Haven, Connecticut: Pontiac, Michigan; Phoenix, Arizona; District [E74-10730] p0056 N75-10526 MANAGEMENT PLANNING Evaluation of Skylab imagery as an information service for investigating land use and natural resources --- Hudson Planning and management of science programs on of Columbia Skylab [E75-10091] River Valley and Long Island, New York [AIAA PAPER 74-1215] p0075 A75-12030 Urban and regional land use analysis: CARETS and An overview of Texas activities in remote sensing p0076 A75-17235
Third Earth Resources Technology Satellite Symposium. [E75-10013] p0059 N75-13336 census cities experiment package LONGITUDE MEASUREMENT [E75-10100] p0025 N75-15118 On the longitudinal localization of the substorm active A demonstration of ERTS-1 analog and digital techniques applied to strip mining in Maryland and West Virginia [NASA-TM-X-70798] p0039 N75-15133 region and its changes during the substorm Volume 2: Summary of results p0077 N75-10549 p0027 A75-11861 [NASA-SP-356] Evaluation of Skylab imagery as an information service LOUISIANA MASS SPECTROSCOPY Mapping of the 1973 Mississippi River floods by the DAA-2 satellite p0028 A75-17139 for investigating land use and natural resources --- New Airborne vapor surveillance system [AD-785585] NOAA-2 satellite n0021 N75-12459 [E75-10068] p0059 N75-14187 Application of ERTS data to environmental inventory and SSACHUSETTS The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus assessment p0016 A75-18069 Investigation of land use of northern megalopolis using A review of initial investigations to utilize ERTS-1 data ERTS-1 imagery [NASA-CR-2459] areas in determining the availability and distribution of living marine resources --- harvest and management of fisheries n0017 N75-10545 [E75-10083] MANNED SPACE FLIGHT p0023 N75-14201 MATERIALS SCIENCE resources in Mississippi Sound and Gulf waters p0043 N75-10555 Skylab contributions to the future The progeny of Skylab - The pioneering phase of our expansion into space p0075 A75-13618 Land use and mapping p0017 N75-10560 MATHEMATICAL MODELS [AIAA PAPER 74-1216] p0063 A75-12031 Mathematical modeling for the detection of fish by an Evaluation of satellite remote sensing and automatic data techniques for characterization of wetlands and coastal marshlands --- detection and measurement of accretion MSFC integrated experiments preliminary report --- for airborne laser p0065 A75-15953 the Skylab program [NASA-TM-X-64881] MAXIMUM LIKELIHOOD ESTIMATES p0078 N75-10967 Planning applications in east central Florida ··· maximum likelihood classification programming estuary system of Atchafalaya River in southern Louisiana Coherent optics potential applications to mapping p0021 N75-13337 [E75-10025] p0044 N75-13341 p0027 A75-14683 Color terrain map of Yellowstone National Park, MEASURING INSTRUMENTS Oil pollution detection, monitoring and law enforcement Modern trends in environmental testing [SLA-73-5931] p0016 N75-10469 siana Gulf Coast waters computer-derived from ERTS MSS data [E75-10027] p0004 A75-17185 MEDITERRANEAN SEA Orbital geology --- an aid to mineral exploration and Evaluation of satellite remote sensing and automatic data Recent and present sedimentation complexity of the mapping the earth's surface
[NASA-TT-F-16000] techniques for characterization of wetlands and coastal Rhone delta and the French mediterranean coast (Golion p0035 N75-10887 marshlands --- Atchafalaya River Basin, Louisiana p0045 N75-14228 project) --- ERTS-1 imagery p0045 N75-14228
Colour composites of ERTS multispectral imagery for Use of a digital computer to construct isoline maps based on a stereomodel of terrain [AD-784170] p0057 N75-11453 [E75-10028] p0050 N75-13344 LOW FREQUENCIES applications in hydrology and oceanography -- Italian Mediterranean and North Sea areas p0045 N75-14231 MEGALOPOLISES The earth as a radio source - Terrestrial kilometric Computer and photogrammetric general land use study of central north Alabama radiation p0011 A75-11211 Investigation of land use of northern megalopolis using ERTS-1 imagery
[NASA-CR-2459] p0017 N75-10545 LUMINESCENCE [NASA-TR-R-431] p0022 N75-13372 Crude and refined petroleum oil structured luminescence Cartographic evaluation of Skylab-A S-192 scanner signatures induced by UV laser or lamp and their remote images --- Chesapeake Bay, Phoenix, and San Francisco sensing applications p0033 A75-17207 MESOMETEOROLOGY Prediction of the Fraunhofer line detectivity of minescent materials --- crude and refined oils, effluents Observing and forecasting local effects from satellite data mesoscale cloud patterns p0016 A75-18153 [E75-10078] p0031 N75-14197 The potential of small-scale mapping from ERTS-1 nagery p0060 N75-14221 Mapping of water bodies in northern Germany from ERTS p0016 A75-18153 and vegetation p0068 A75-17221 MESOSPHERE LUNAR RANGEFINDING Ice particulates in the mesosphere p0012 A75-16752 p0051 N75-14227 Geodetic aspects of laser distance measurements to the METEOROLOGICAL CHARTS moon and radio-interference measurement ments to quasars p0027 A75-14646 Natural-resources mapping of Ethiopia from ERTS-1 Meteorological data. Volume 145 no. 1: The European climatology 1974 meteorological satellite data from the imagery p0031 N75-14235 Scientific research and experimental design planning and Berlin zone of reception, part 1, 1st quarter coordination for 1973 DO023 N75-13479 p0081 N75-14259 M [AD-785998] METEOROLOGICAL FLIGHT The next problems in the area of the implementation of Automated remote measurement system in agricultural meteorology --- for crop survey by aerial photometry planning for cartographical-geodetic work MAGNETIC DIPOLES p0032 N75-15143 p0064 A75-12947 A generator of high-power current pulses of semisinusoidal shape for an aerial variant of the method Fully automated compilation of gravity and geophysical Airborne survey of major air basins in California maps [AD-785838] p0011 A75-14360 of transient processes --- for aerial prospecting p0053 A75-13310 p0032 N75-15147 The wind vector in a layer over water p0041 A75-15179 MAGNETIC DISTURBANCES Catalog of earthquake photographs --

A remote sensing study of Pacific hurricane Ava

Electronics in meteorological instrumentation

METEOROLOGICAL INSTRUMENTS

p0012 A75-17112

p0065 A75-14726

p0072 N75-13455

p0043 N75-10566

a geomagnetic disturbance

electrons and

disturbance

Morning-evening asymmetry of the thermosphere during

Detection of synchronous changes in fluxes of energetic

during

gamma-quanta

p0011 A75-11103

p0064 A75-12317

MARINE ENVIRONMENTS

[NASA-CR-137475]

at VIMS

Remote sensing applications in marine science programs

SUBJECT INDEX MOUNTAINS

MICROWAVE IMAGERY Remote sensing geophysics from Skylab --- Southern The point visibility meter - A lurward scattle for the measurement of aerosol extinction coefficient p0070 A75-19266 The point visibility meter - A forward scatter instrument Skylab earth resources program p0011 A75-13577 A radio picture of the earth --- microwave remote sensing d Nevada minerals and rock p0030 N75-12413 [E75-10045] Systems study of an automated fire weather data from satellites n0066 A75-16191 Geologic and mineral and water re sources investigations Four channel simultaneous X-L band imaging SAR dar p0054 A75-17116 in western Colorado, using Skylab EREP data [NASA-TM-X-62402] p0037 N75-14190 p0006 N75-12277 radar [E75-10071] Emissivities and forward scattering of natural and The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 7: Nonreplenishable natural resources: Minerals, fossil fuels and geothermal energy sources [NASA-CR-141264] METEOROLOGICAL PARAMETERS man-made material at three millimeter wavelength Measurements of meteorological parameters and air DO067 A75-17197 pollution spread over Stockholm, Sweden, by means of emote sensing and airborne instrumentation p0018 N75-11531 The usefulness of imaging passive microwave for rural p0055 A75-17199 and urban terrain analysis
Microwave radiometric characteristic METEOROLOGICAL RADAR INNESOTA p0049 A75-17200 terrain The role of lidar in environmental sensing 🏎 laser A study of Minnesota forests and lakes using data from MICROWAVE OSCILLATORS ceilometer and visual range measurement application Earth Resources Technology Satellites [NASA-CR-140865] Microwave rotational spectroscopy: A physical technique p0018 N75-11529 p0007 N75-12421 for specific pollutant monitoring **METEOROLOGICAL SATELLITES** Design data collection with Skylab/EREP microwave MICROWAVE RADIOMETERS Satellites and computer in the service of meteorology: WWW - The global weather observation system instrument S-193 --- Texas, Utah, Illinois, and Minnesota Simultaneous active and passive microwave response [E75-10080] p0072 N75-14199 of the earth - The Skylab radscat experiment DO053 A75-11274 p0012 A75-17114 MISSILE RANGES Technology today for tomorrow: Proceedings of the Eleventh Space Congress, Cocoa Beach, Fla., April 17-19, 1974, Volumes 1 & 2 p0011 A75-13576 Pacific range electromagnetic signature studies /Project On the detectability of atmospheric carbon monoxide by microwave remote sensing Moisture detection from Skylab p0013 A75-17129 PRESS/ MISSILE SIGNATURES soil studies for Texas p0002 A75-17145 Rainfall estimations from geosynchronous satellite Pacific range electromagnetic signature studies / Project PRESS/ p0042 A75-17479 and Kansas sites Deteriorating effects on 3 mm way e passive imagery Tagery Observing and forecasting local effects from satellite data p0055 A75-17198 MISSION PLANNING Microwave radiometric characteristics of snow-cover mesoscale cloud patterns p0016 A75-18153 Planning and management of science programs on p0049 A75-17200 Space iconics Skylah Electrically scanning microwave radiometers [AIAA PAPER 74-1215] p0075 A75-12030 [NASA-TT-F-798] p0057 N75-11419 Multi-frequency radiometric measurements of foam and MISSISSIPPI Geographic correlation of television pictures obtained Correlation of multispectral imagery with water analysis from weather satellites p0071 N75-11432 mono-molecular slick --- microwave sensing of ocean process pooles A75-17208 - The Ross Barnett Reservoir Remote Sensing Project p0014 A75-17165 Geographical correlation of TV and IR images obtained weather satellites --- cloud cover p photography p0071 N75-11433 Application and sensitivity investigation of Fourier transforms for microwave radiometric inversions [NASA-CR-140754] p0070 N75-11310 A review of initial investigations to utilize ERTS-1 data in determining the availability and distribution of living marine resources --- harvest and management of fisheries resources in Mississippi Sound and Gulf waters System for electronic transformation and geographic Design data collection with Skylab/EREP microwave instrument S-193 --- Texas, Utah, Illinois, and Minnesota correlation of satellite television information --p0071 N75-11434 p0043 N75-10555 p0072 N75-14199 Use of TV pictures obtained from weather satellites to [E75-10080] MICROWAVE REFLECTOMETERS Application of remote sensing to state and regional study the geological structure of the earth problems --- for Mississippi [NASA-CR-140805] p0036 N75-11438 Microwave profiling of snowpack free-water content DO078 N75-11414 p0049 A75-17950 **METEOROLOGICAL SERVICES** MICROWAVE SCATTERING MISSISSIPPI DELTA (LA) Satellites and computer in the service of meteorology: WWW - The global weather observation system Mapping of the 1973 Mississippi River floods by the NOAA-2 satellite p0028 A75-17139 measurement of soil water conten [NASA-CR-141685] p0001 A75-16242 p0053 A75-11274 Radar response to vegetation soil moisture manning A review of initial investigations to utilize FRTS-1 data METEOROLOGY via microwave backscattering
MICROWAVE SENSORS p0005 A75-18556 in determining the availability and distribution of living marine resources --- harvest and management of fisheries Conference on Weather Forecasting and Analysis, 5th, St. Louis, Mo., March 4-7, 1974. Preprints resources in Mississippi Sound and Gulf waters p0043 N75-10555 On the feasibility of remote monitoring of soil moisture p0016 A75-18151 Meteorological interpretation of high resolution images p0067 A75-17147 MICROWAVE SPECTRA p0017 N75-10560 for remote sensing of earth resources p0060 N75-14229 Land use and mapping A microwave correlation spectron MEXICO MISSISSIPPI RIVER (US) p0063 A75-11732 onitorina Mapping of the 1973 Mississippi River floods by the A simulation of the San Andreas fault experiment MICROWAVES p0033 A75-11068 DO028 A75-17139 NOAA-2 satellite Fault tectonics and earthquake hazards in the Peninsular Ranges, Southern California --- Gulf of California [E75-10074] p0031 N75-14193 Preliminary analysis of Skylab RADSCAT results over Geographic applications of ERTS-1 imagery to landscape the ocean --- using radar backscatter and microwave change --- Mississippi River and Great Smoky Mountains of Tennessee and North Carolina [NASA-CR-141454] DO046 N75-14793 MICHIGAN p0034 N75-10537 [E74-10810] MIE SCATTERING Resource analysis applications in Michigan --- NASA MISSOURI Lectures on the scattering of light --- by dielectric p0069 A75-17231 Mapping of the 1973 Mississippi River floods by the NOAA-2 satellite p0028 A75-17139 Study of recreational land and open space using Skylab p0019 N75-11556 p0028 A75-17139 - southeast Michigan MIGRATION Applications of Skylab EREP photographs to mapping DO020 N75-12404 Utilization of Skylab (EREP) system for appraising of landforms and environmental geology in the Great Plains Developing processing techniques for Skylab data --changes in continental migratory bird habitat and Midwest --- Illinois, Iowa, Kansas, Missouri, Nebraska, and South Dakota p0006 N75-10543 [E74-10817] Utility of ERTS for monitoring the breeding habit of migratory waterfowl --- United States and Canada p0006 N75-10557 [E75-10040] p0058 N75-12409 p0037 N75-13355 [E75-10055] Urban and regional land use analysis: CARETS and ensus cities experiment package --- New Haven. MOISTURE CONTENT Census cities Microwave emission characteristics of moist soils Connecticut; Pontiac, Michigan; Phoenix, Arizona; District Utilization of Skylab (EREP) system for appraising p0001 A75-15180 changes in continental migratory bird habitat [E75-10061] p0008 Passive microwave sensing of moist soils [E75-10091] p0024 N75-15109 p0008 N75-13361 p0066 A75-17124 Study of recreational land and open space using Skylab magery --- snow and ice hydrology of southeast Utilization of Skylab (EREP) system for appraising Detection of moisture and moisture related phenomena changes in continental migratory bird habitat [E75-10087] p0009 from Skylab --- Kansas, Oklahoma, and Arkansas p0009 N75-15105 E75-10092 [E75-10035] p0007 N75-12405 DO024 N75-15110 MILLIMETER WAVES MICROBIOLOGY MOISTURE METERS Emissivities and forward scattering of natural and man-made material at three millimeter wavelength Skylab environmental and crew microbiology studies p0024 N75-14366 Microwave profiling of snowpack free-water content p0049 A75-17950 DO067 A75-17197 MICROORGANISMS MOLECULAR ABSORPTION Microbial abundance and thermantarctic dry valley soils MINERAL DEPOSITS The effect of atmospheric water classification of ERTS data Geologic and mineral and water resources investigations in Western Colorado, using Skylab EREP data p0005 A75-17973 p0013 A75-17131 Skylab environmental and crew robiology studies p0024 N75-14366 MONOCHROMATIC RADIATION p0034 N75-10530 [E74-10802] Some analytical results for radiative transfer in thick MICROWAVE EMISSION An interdisciplinary analysis of multispectral satellite data atmospheres --- of monochromatic radiation p0019 N75-11555 ficrowave emission characteristics of moist soils for selected cover types in the Colorado Mountains, using automatic data processing techniques --- hydrology, ecology, geology, vegetation, and mineral deposits [E75-10002] p0078 N75-12390 n0001 A75-15180 MONTANA The radio emission of the earth as Color terrain map of Yellowstone National Park, computer-derived from ERTS MSS data book p0076 A75-15319 A study of microwave emission p Geologic and mineral and water resources investigations p0004 A75-17185 p0041 A75-17120 in Western Colorado, using Skylab EREP data --- Utah Experiment to evaluate feasibility of utilizing Skylab-EREP MICROWAVE EQUIPMENT [E75-10043] p0058 N75-12412 Signatures of various earth surfaces measured by the imbus-5 microwave spectrometer p0066 A75-17119 remote sensing data of tectonic analysis of the Bighorn Mountains region, Wyoming-Montana --- Black Hills, South Geologic and mineral and water resources investigations western Colorado, using Skylab EREP data Nimbus-5 microwave spectrometer Passive microwave sensing of moist p0025 N75-15123 Dakota/Wyoming [E75-10105] p0066 A75-17124 [E75-10014] p0030 N75-12395 MINERAL EXPLORATION MICROWAVE FREQUENCIES Water resource and hazard planning report for the Clark Fork River Valley above Missoula, Missoula County, A technology to renovate the search for no A Shuttle/SpaceLab RF environm nt survey facility p0016 A75-19457 p0063 A75-10434 deposits

Remote sensing to detect the toxic effects of metals

Orbital geology --- an aid to mineral exploration and

Mineral resources, geological structures, and landform

p0035 N75-10887

p0035 N75-10561

on vegetation for mineral exploration p0034 A75-17226

mapping the earth's surface [NASA-TT-F-16000]

MINERALS

Theoretical investigations on the determination of the

assuming multispectral thermal radiation measurements at

MICROWAVE HOLOGRAPHY
Investigation of microwave hologram techniques for

in the atmosphere

p0024 N75-14331

p0067 A75-17195

vertical temperature distribution

60 GHz

[BMFT-FB-W-74-04]

application to earth resources

[PB-236041/0] MONTE CARLO METHOD Solution of the radiative transfer theory problems by the Monte Carlo method p0058 N75-11551

Montana

MOUNTAINS Fault tectonics and earthquake hazards in the Peninsular Ranges, Southern California --- Gulf of California [E75-10074] p0031 N75-14193

n0051 N75-14280

An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104] p0073 N75-15122 MULTISPECTRAL BAND SCANNERS Automatic land use classification using Skylab S-192 multispectral data p0064 A75-12036 [AIAA PAPER 74-1224] Mapping soil features from multispectral scanner data p0027 A75-15435 A method of specifying remotely sensed units for soil sample points sample points
The effect of atmospheric water vapor on automatic classification of ERTS data p0013 A75-17131 Adaptive processing of MSS data using a decision-directed Kalman filter p0055 A75-17151 SICLOPS - A system of computer programs for rectified apping of airborne scanner imagery p0055 A75-17156 Effect of atmospheric haze and sun angle on automatic classification of ERTS-1 data nOO14 A75-17157 Skylab S-192 ratio codes of soil, mineral, and rock spectra for ratio image selection and interpretation p0003 A75-17158
Coastal wetlands analysis from ERTS MSS digital data p0048 A75-17178 Color terrain map of Yellowstone computer-derived from ERTS MSS data vstone National Park. p0004 A75-17185 Land use classification and ground truth correlations from simultaneously acquired aircraft and ERTS-1 MSS data p0014 A75-17188 A motion-compensated spatial scanner --- MSS design p0068 A75-17213 Statistical separability of agricultural cover types in subsets of one to twelve spectral channels p0005 A75-17216 Multi-aspect techniques in remote sensing --- for MSS p0068 A75-17217 imagery Unsupervised mapping of geologic features and soils in California p0034 A75-17225 Geological and geophysical studies from space and ospects for their development p0035 N75-11438 prospects for their development Quantitative determination of stratospheric aerosol [F75-10010] p0020 N75-12393 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil --- Southern Texas p0006 N75-12394 [E75-10012] A cloud physics investigation utilizing Skylab data [E75-10021] p0020 N75-12400 Experimental evaluation of atmospheric effects on radiometric measurements using the EREP of Skylab ... Bonneville Salt Flats, Utah [E75-10036] o0020 N75-12406 Developing processing techniques for Skylab data ---Michigan test site [E75-10040] p0058 N75-12409 New theoretical models and ratio imaging techniques associated with the NASA earth resources spectral nformation system [NASA-CR-140357] n0058 N75-12417 Correlation and registration of ERTS multispectral imagery --- by a digital processing technique [NASA-CR-134294] p0058 N75-12420 The ERTS-1 investigation. Volume 7: Erts-1 land-use analysis of the Houston area test site [E75-10009] p0021 N75-13334 Application of Skylab EREP photographs to study of the modern episode of accelerated erosion in southern [F75-10054] p0036 N75-13354 Spectroradiometric and multispectral scanner over Lake [E75-10066] n0072 N75-13366 Automated selection and location of terrain features on the basis of spectral characteristics p0030 N75-13375 Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcances [E75-10069] p0037 N75-14188 p0037 N75-14188 The Great Basin investigation [E75-10076] p0037 N75-14195 Cartographic evaluation of Skylab-A S-192 scanner images --- Chesapeake Bay, Phoenix, and San Francisco [E75-10078] DO031 N75-14197 Quantitative determination of stratospheric aerosol characteristics [E75-10079] p0023 N75-14198 Evaluation of ERTS-1 data by analogue and digital chaigues --- Germany and Italy p0059 N75-14218 techniques --- Germany and Italy Optical processing of ERTS-1 (MSS) images: Examples for the Arcachon basin and Rhone delta and Arcachon Basin, France p00 Rhone Delta p0060 N75-14223 Colour composites of ERTS multispectral imagery for applications in hydrology and oceanography --- Italian Mediterranean and North Sea areas p0045 N75-14231

Comparison of various numerical techniques for ssification of multispectral data. Application to geology
- ERTS-1 imagery of Rhone Valley (France)
p0038 N75-14244

Use of ERTS-1 imagery to interpret wind-erosion hazard Reformatting Skylab and MSS data over northern in the Sandhills of Nebraska [E75-10044] [E75-10016] p0059 N75-13347 p0061 N75-15101 Developing processing techniques for Skylab data [E75-10082] p0061 N75-15102 Skylab water depth determination --- Lake Erie and Lake [E75-10048] Analysis of Skylab MSS data, and calibration of thermal p0050 N75-13348 Remote sensing geophysics from Skylab --- rationing-type algorithms to enhance multispectral data data over Lake Monroe, Indiana p0073 N75-15113 (E75-10095) 75-10052] p0030 N75-13352 The ERTS-1 investigation (ER-600). Volume 2: ERTS-1 [E75-10052] A cloud physics investigation utilizing Skylab data [E75-10099] p0025 N75-15117 coastal/estuarine analysis --- Galveston Bay, Texas [NASA-TM-X-58118] n0051 N75-MULTISPECTRAL PHOTOGRAPHY p0051 N75-13371 Interactive computer processing for land use planning Remote sensing of ocean current boundary layer --- Gulf p0054 A75-17134 of Mexico Correlation of multispectral imagery with water analysis [E75-10031] p0045 N75-14185 - The Ross Barnett Reservoir Remote Se Sensing Project p0014 A75-17165 Remote sensing of ocean current boundary layer --- Gulf of Mexico Grass canopy bidirectional spectral reflectance p0003 A75-17170 Geologic and mineral and water resources investigations western Colorado, using Skylab EREP data Evaluation of Skylab imagery as an information service western Colorado, using Skylab Ener data [75-10071] p0037 N75-14190 Skylab study of water quality --- Kansas reservoirs and [E75-10071] for investigating land use and natural resources --- Long Island, New York p0056 N75-10526 [E74-10730] [E75-10075] p0051 N75-14194 Developing processing techniques for Skylab data ---Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E74-10799] p0057 N75-10527 Overall evaluation of ERTS-1 imagery for cartographic application --- Australia, Antarctica, U.S., and U.S.S.R. [E74-10800] p0029 N75-10528 p0008 N75-14196 Study to develop improved spacecraft snow survey methods using Skylab/EREP data --- Sierra Nevada, Cascades, and Great Plains An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10081] 75-10081] p0073 N75-14200 The application of Skylab EREP data for land use [E74-10803] o0057 N75-10531 classification and mapping in the Cleveland and Columbus Skylab a proposal aerotriangulation with very small scale --- North Carolina and Virginia photography [E74-10804] [E75-10083] p0023 N75-14201 p0070 N75-10532 European earth-resources satellite ex [ESRO-SP-100] Thailand national programme of the Earth Resources p0081 N75-14216 SRO-SP-100]
Preliminary evaluation of ERTS-1 imagery of Central p0060 N75-14242 Technology Satellite p0077 N75-10533 [E74-10806] A comparison of the capabilities of ERTS-1 and the The Thailand programme of the Earth Resources Technology Satellite [E74-10807] Skylark earth-resources rocket for resources surveying in central Argentina p0023 N75-14250 Central Argentina p0023 N75-14250
The remote identification of terrain features and materials p0077 N75-10534 Assessment of ERTS-1 imagery as a tool for regional at Pennsylvania test sites: An investigative study of geological analysis in New York State --- Lake Ontario [E74-10809] p0034 N75-10536 [PB-235992/5] n0031 N75-14263 ERTS-1 imagery interpretation techniques in the Tennessee Valley --- land use and soil mapping [E74-10811] p0016 N75-10538 [PB-235992] 5] pouds in restricted Applicability of Skylab remote sensing for detection and monitoring of surface mining activities — evironmental effects and reclamation in Tennessee and Kentucky Skylab-EREP investigations of wetlands ecology --- New [E75-10085] p0038 N75-15103 and Chesapeake Bay A Skylab program for the International Hydrological [E74-10816] p0049 N75-10542 ecade (IHD) --- Lake Ontario Basin Utilization of Skylab (EREP) system for appraising changes in continental migratory bird habitat p0052 N75-15111 [E75-10093] A Skylab program for the International Hydrological Decade (IHD) --- Lake Ontario Basin p0006 N75-10543 [E74-10817] Geologic evaluation and applications of ERTS-1 imagery [E75-10094] n0052 N75-15112 p0034 N75-10552 Usefulness of Skylab color photography and ERTS-1 An analysis of applications development systems for remotely sensed, multispectral data for the Earth Observations Division of the NASA Lyndon B. Johnson multispectral imagery for mapping range vegetation types n southwestern Wyoming p0010 N75-15115 Urban and regional land use analysis: CARETS and p0071 N75-11447 census cities experiment package [E75-10100] Study of atmospheric effects in Skylab data [E75-10006] p0020 N75-12391 Experiment to evaluate feasibility of utilizing Skylab-EREP remote sensing data of tectonic analysis of the Bighorn p0025 N75-15118 Skylab: A proposal aerotriangulation with very small scale photography --- North Carolina and Virginia [E75-10101] p0073 N7 p0073 N75-15119 The Great Basin investigation --- photomapping lithology Mountains region, Wyoming-Montana --- Black Hills, South Dakota/Wyoming and geological structures [E75-10102] p0038 N75-15120 p0030 N75-12395 [E75-10014] Evaluation of Skylab imagery as an information service for investigating land use and natural resources Multidisciplinary analysis for highway engineering purposes --- Maine [E75-10103] n0020 N75-12397 [E75-10018] n0025 N75-15121 Multispectral photography for earth resources
[NASA-CR-141283] p0074 N Study of recreational land and open space using Skylab p0074 N75-15138 [F75-10020] n0020 N75-12399 A Skylab program for the International Hydrological Decade (IHD) --- Lake Ontario Basin N [E75-10039] p0050 N75-12408 Index of Skylab data available at Colorado School of Mines --- multispectral photography of Colorado NASA PROGRAMS NASA-KSC's earth resources benefits from space poorts A75-13592 [E75-10041] p0072 N75-12410 In situ spectroradiometric calibration of EREP imagery exploration and oceanography of Block Island Sound --- Arizona NATURAL GAS geology [E75-10042] The use of space photos for search of oil and gas fields n0072 N75-12411 A study of the usefulness of Skylab EREP data for earth Offshore investigation: Producible shut-in leases as of resources studies in Australia --- Australia geology p0036 N75-12415

p0034 A75-17229

January 1974, phase 2 --- an estimation of natural gas reserves in offshore wells p0036 N75-11458

NEAR INFRARED RADIATION

Use of feature extraction techniques for the texture and

Evaluation of Skylab imagery as an information service

for investigating land use and natural resources --- Hudson River Valley and Long Island, New York

- Evaluation of satellite remote sensing and automatic data techniques for characterization of wetlands and coastal

marshlands --- Atchafalaya River Basin, Louisiana [E75-10028] p0050 N75-13344
Utilization of ERTS-1 imagery in cultivation and settlement site identification and carrying capacity estimates

p0059 N75-13335

p0059 N75-13336

n0008 N75-13345

context information in ERTS imagery: Spectral and textural processing of ERTS imagery --- classification of Kansas

[E75-10011]

[E75-10013]

[F75-10029]

n Upper Volta and Niger

An examination of whiteouts and transmission through ice fog in the visible and near infrared [DREO-TN-74-18] p0021 N75-12487 Remote sensing geophysics from Skylab --- near infrared reflectance anomalies of andesites and basalts in Southern California and Nevada [275-10051] p0036 N75-13351

NEAR ULTRAVIOLET RADIATION

Measurements of the extraterrestrial solar radiant flux from 2981 to 4000 A and its transmission through the earth's atmosphere as it is affected by dust and ozor n0070 A75-19547

NEARSHORE WATER California nearshore processes -currents and sediments p0048 A75-17179

Surface currents along the California coast observed on Remote sensing of ocean current boundary layer --- Gulf NIMBUS 4 SATELLITE p0048 A75-17180 Nimbus 4 observations of changes in total ozone and **ERTS** imagery of Mexico stratospheric temperatures during a sudden warming [F75.10098] NEBRASKA p0046 N75-15116 p0012 A75-14841 OCEAN DATA ACQUISITIONS SYSTEMS Applications of remote sensing by the state Remote sounding of stratospheric temperatures using p0069 A75-17232 Preliminary results of fisheries investigated with Skylab-3 high resolution radiance measurements from the IRIS-D [NASA-CR-141130] p0022 N75-13452 p0003 A75-17167 Satellite pictures can help quide Sandhills irrigation --ot irrigation in Sand Hills Region, Nebras Report of the meeting on drifting buoys for the first GARP NIMBUS 5 SATELLITE [E75-10023] p0021 N75-13339 global experiment Signatures of various earth surfaces measured I IGARP-SPEC-REPT-131 Use of ERTS-1 imagery to interpret wind-erosion hazard p0043 N75-10674 p0066 A75-17119 Nimbus-5 microwave spectrometer OCEAN SURFACE in the Sandhills of Nebraska NITROUS OXIDES [E75-10044] p0059 N75-13347 The detection by ERTS-1 of wind-induced ocean surface features in the lee of the Antilles Islands Optical-acoustic detection of low concentrations of Applications of Skylab EREP photographs to mapping landforms and environmental geology in the Great Plains hydrogen fluoride, nitrous oxide, and carbon dioxide in gases p0041 A75-10644 using pulsed hydrogen-fluoride laser radiation A satellite-aircraft thermal study of the upwelled waters f Spanish Sahara p0041 A75-10645 and Midwest --- Illinois, Iowa, Kansas, Missouri, Nebraska, p0063 A75-11443 off Spanish Sahara South Dakota NOAA 2 SATELLITE [875-10055] p0037 N75-13355 Mapping of the 1973 Mississippi River floods by the NOAA-2 satellite p0028 A75-17139 Monitoring and predicting El Nino invasions --- equatorial NEVADA Pacific Ocean-atmosphere interactions Remote sensing geophysics from Skylab --- Southern Environmental satellite imagery, September 1974 p0041 A75-12719 California and Nevada minerals and rocks p0058 N75-11571 Estimation of sea surface temperature from remote [£75-10045] p0030 N75-12413 NOAA 3 SATELLITE sensing in the 11to 13-micron window reg Images from the NOAA-3 very high resolution radiometer p0065 A75-14803 Remote sensing geophysics from Skylab --- near infrared over the North Sea and adjoining countries Skylab radar altimeter - Short-wavelength perturbations tected in ocean surface profiles p0066 A75-16096 reflectance anomalies of andesites and basalts in Southern p0066 A75-16188 California and Nevada detected in ocean surface profiles [£75-10051] p0036 N75-13351 NORTH AMERICA Measurement of sea surface currents using airborne Feasibility of using S-191 infrared spectra for geological A satellite view of weather systems over the North Doppler radar and inertial navigation systems p0063 A75-10790 p0042 A75-17203 studies from space --- Nevada Observing and forecasting local effects from satellite data [£75-10088] p0038 N75-15108 Multi-frequency radiometric measurements of foam and mesoscale cloud patterns p0016 A75-18153 mono-molecular slick --- microwave sensing of ocean p0068 A75-17208 Feasibility of using S-191 infrared spectra for geological udies from space --- Nevada Study to develop improved spacecraft snow survey surface methods using Skylab/EREP data --- Sierra Nevada. Cascades, and Great Plains [E75-10089] p0038 N75-15107 The gravity field and the definition of stationary sea NEW ENGLAND (US) surface topography n0042 A75-18124 p0073 N75-14200 [E75-10081] Land use and mapping p0017 N75-10560 AIDJEX bulletin number 25 --- study of pack ice, sea A radar altimeter for satellite application p0056 N75-10425 ce, and Beaufort Sea ice [AD-784373] New England reservoir management p0045 N75-14262 [PB-235298/7] [E75-10065] Report of the meeting on drifting buoys for the first GARP o0050 N75-13365 global experiment [GARP-SPEC-REPT-13] NORTH CAROLINA NEW GUINEA (ISLAND) Skylab a proposal aerotriangulation with very small scale p0043 N75-10674 Australian and Soviet gravity surveys along the Australian · · · North Carolina and Virginia The Coho Project: Living resources prediction feasibility study. Volume 3: System evaluation report calibration line p0070 N75-10532 [E74-10804] BULL- 161 p0031 N75-13454 IPB-234059/41 Geographic applications of ERTS-1 imagery to landscape change --- Mississippi River and Great Smoky Mountains p0044 N75-10683 NEW HAVEN (CT) Urban and regional land use analysis: CARETS and census cities experiment package --- New Haven, Connecticut; Pontiac, Michigan; Phoenix, Arizona; District OCEANOGRAPHY of Tennessee and North Carolina Skylab altimeter applications and scientific results [AIAA PAPER 74-1221] p0063 A75-1 [E74-10810] p0034 N75-10537 AAA PAPER 74-1221] p0063 A75-12035 Interaction of marine geodesy, satellite technology and of Columbia [E75-10091] Utilization of EREP data in geological evaluation regional p0024 N75-15109 planning, forest management, and water management in ocean physics North Carolina [NASA-CR-137470] NEW JERSEY p0043 N75-10671 Application of ERTS-1 data to the protection and p0008 N75-13363 Ocean color, a three component system [NASA-CR-139145] p00p0044 N75-11411 Skylab: A proposal aerotriangulation with very small scale photography --- North Carolina and Virginia management of New Jersey's coastal environment p0042 A75-17233 A simulation of synthetic aperture radar imaging of ocean Skylab-EREP investigations of wetlands ecology ... New [E75-10101] p0073 N75-15119 Jersey and Chesapeake Bay [NASA-TM-X-72629] p0044 N75-12172 NORTH SEA An assessment of the potential contributions to oceanography from Skylab visual observations and hand-held photography --- Pacific Ocean, Atlantic Ocean p0049 N75-10542 Images from the NOAA-3 very high resolution radiometer over the North Sea and adjoining countries Urban and regional land use analysis: CARETS and sus cities experiment package p0066 A75-16188 p0020 N75-12398 and Gulf of Mexico Colour composites of ERTS multispectral imagery for Remote sensing geophysics from Skylab --- rationing-type [E75-10017] n0044 N75-12396 applications in hydrology and oceanography --- Italian Mediterranean and North Sea areas p0045 N75-14231 In situ spectroradiometric calibration of EREP imagery and oceanography of Block Island Sound --- Arizona Igorithms to enhance multispectral data p0030 N75-13352 [E75-10052] NORTHERN HEMISPHERE Urban and regional land use analysis: CARETS and census cities experiment package --- New Haven, Connecticut; Pontiac, Michigan; Phoenix, Arizona; District Comparison of the Southern and Northern Hemisphere p0072 N75-12411 eneral circulation characteristics as determined by satellite The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity ozone data [UCRL-75556] o0017 N75-10638 [E75-10091] p0024 N75-15109 of service. Volume 9: Oceans Urban and regional land use analysis: CARETS and INASA-CR-141270 NORWAY o0080 N75-14214 ERTS-1 images in glaciology and hydrology: Norwegian Colour composites of ERTS multispectral imagery for consus cities experiment package p0051 N75-14233 applications in hydrology and oceanography --- Italian Mediterranean and North Sea areas p0045 N75-14231 OFFSHORE ENERGY SOURCES [E75-10100] p0025 N75-15118 experience Exploration for fossil and nuclear fuels from orbital Computer derived coastal water classifications via ectral signatures p0042 A75-17177
Evaluation of Skylab imagery as an information service altitudes --- results of ERTS program for oil exploration [NASA-TM-X-70781] p0017 N75-11413 Offshore investigation: Producible shut-in leases as of January 1974, phase 2 --- an estimation of natural gas for investigating land use and natural resources --- Long reserves in offshore wells p0036 N75-11458 NUMERICAL ANALYSIS York Comparison of various numerical techniques for classification of multispectral data. Application to geology --- ERTS-1 imagery of Rhone Valley (France) Multidisciplinary applications of ERTS and Skylab data [E74-10730] p0056 N75-10526 Ohio p0015 A75-17234
Automatic mapping of strip mine operations from pacecraft data p0034 A75-18065 Assessment of ERTS-1 imagery as a tool for regional eological analysis in New York State --- Lake Ontario p0038 N75-14244 [E74-10809] p0034 N75-10536 spacecraft data Mineral resources, geological structures, and landform riveys p0035 N75-10581 Evaluation of Skylab imagery as an information service Automated strip mine and reclamation mapping from RTS --- Ohio coal p0035 N7S-10556 0 ERTS --- Ohio coal The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus for investigating land use and natural resources --- Hudson River Valley and Long Island, New York OCEAN CURRENTS areas p0059 N75-13336 [E75-10083] [£75-10013] Satellite photography of eddies in the gulf loop current p0023 N75-14201 Evaluation of Skylab imagery as an information service OIL EXPLORATION p0041 A75-11255 for investigating land use and natural resources --- New Oil exploration needs for digital processing of imagery p0033 A75-10437 Measurement of sea surface currents using airborne Doppler radar and inertial navigation systems p0059 N75-14187 [E75-10068] p0042 A75-17203 Plutonic structure of the earth's crust on space images p0027 A75-17125
An evaluation of the suitability of ERTS data for the Evaluation of Skylab imagery as an information service Remote sensing of ocean current boundary layer -for investigating land use and natural resources --- New purposes of petroleum exploration --- Anadarko Basin in Oklahoma and Texas p0035 N75-10553 [E74-10801] p0042 N75-10529 [NASA-CR-141372] p0024 N75-15100 An assessment of the potential contributions to oceanography from Skylab visual observations and hand-held photography --- Pacific Ocean, Atlantic Ocean Exploration for fossil and nuclear fuels from orbital NIGER altitudes --- results of ERTS program for oil exploration
[NASA-TM-X-70781] p0017 N75-11413 Earth Resources Technology Satellite-1 (ERTS-1) data and anthropology: Use of these data in carrying capacity estimates for sites in Upper.Volta and Niger [E75-10001] p0078 N75-11406 and Gulf of Mexico OIL FIELDS [E75-10017] p0044 N75-12396 [E75-10001] p0078 N75-11406
Utilization of ERTS-1 imagery in cultivation and settlement site identification and carrying capacity estimates The use of space photos for search of oi and gas fields p0034 A75-17229 Remote sensing of ocean current boundary layer --- Gulf of Mexico p0045 N75-14185 [E75-10031] in Upper Volta and Niger Development of an experimental airborne laser remote ensing system for the detection and classification of oil pills p0068 A75-17206 [E75-10029] Remote sensing of ocean current boundary layer --- Gulf p0008 N75-13345 NIGERIA [E75-10067] An examination of the extent of fire in the grassland Multi-frequency radiometric measurements of foam and and savanna of Africa along the southern side of the Sahara p0002 A75-17104

Pedological interpretation of ERTS-1 imagery of the Preliminary analysis of Skylab RADSCAT results over a mono-molecular slick --- microwave sensing of ocea p0068 A75-17208 Sahara the ocean --- using radar backscatter and microwave Oil slick detection by X-band synthetic aperture radar p0068 A75-17209

INASA-CR-141454)

Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243

p0046 N75-14793

The application of spectrometric and polarization techniques for remote sensing of oil on sea water p0068 A75-17210 Oil-on-Water sensor --- for detecting oil by emitting videband IR pulse [AD-784104] Oil pollution detection, monitoring and law enforcement --- Louisiana Gulf Coast waters [E75-10027] p0022 N75-13343

OKLAHOMA An evaluation of the suitability of ERTS data for the purposes of petroleum exploration --- Anadarko Basin in p0035 N75-10553 Detection of moisture and moisture related phenomena

from Skylab --- Kansas, Oklahoma, and Arkansas p0007 N75-12405

OPTICAL DATA PROCESSING

Oil exploration needs for digital processing of imagery p0033 A75-10437

Coherent optics potential applications to mapping p0027 A75-14683 Coherent optics in mapping; Proceedings of the Tutorial

Seminar and Technology Utilization Program, Rochester, N.Y., March 27-29, 1974 p0027 A75-16676 Cloud-screening from aerial photography applying coherent optical pattern recognition tech p0054 A75-16690

Classification and mapping of coal refuse, vegetative cover types, and forest types by digital processing ERTS-1 p0054 A75-17110

unsupervised crop p0002 A75-17113 clustering algorithm for classification Adaptive processing of decision-directed Kalman filter MSS data using a p0055 A75-17151

Implementation of an advanced table look-up classifier for large area land-use classification p0014 A75-17153 Modifications in a computer-implemented method for the

detection and extraction of objects in aerial photographs p0055 A75-17154

SICLOPS - A system of computer programs for rectified mapping of airborne scanner imagery p0055 A75-17156 Boundaries of ERTS and aircraft data within which useful water quality information can be obtained

p0014 A75-17182

Optical data processing analysis of stream patterns exhibited on ERTS-1 imagery p0056 A75-17219 Optical processing of ERTS-1 (MSS) images: Examples for the Arcachon basin and Rhone delta p0060 N75-14223 and Arcachon Basin, France

ARNICA/ERTS-1 experiment. Results and perspectives --- image processing for southwest France and Pyre p0081 N75-14246

OPTICAL DENSITY

Optical film density values from color IR photography for wetland soils mapping p0005 A75-18073

OPTICAL EQUIPMENT

Optical quantum generators in geodetic measurements
[AD-787281] p0032 N75-15237 p0032 N75-15237

TICAL MEASURING INSTRUMENTS

Prediction of the Fraunhofer line detection luminescent materials --- crude and refined oils, detectivity of p0068 A75-17221 and vegetation

OPTICAL POLARIZATION Remote detection of soil surface moisture

DOOO3 A75-17146

OPTICAL PROPERTIES

Boundaries of ERTS and aircraft data within which useful water quality information can be obtained p0014 A75-17182

OPTICAL RADAR

Remote measurements of water pollution with a lidar polarimeter p0014 A75-17205 The role of lidar in environmental sensing · · · laser

ceilometer and visual range measurement application p0070 A75-19274

Lidar observations of atmospheric particulate content p0019 N75-11559

OPTICAL REFLECTION

Vegetation canopy reflectance D0001 A75-16241 Grass canopy bidirectional spectral reflectance p0003 A75-17170

OPTICAL SCANNERS

A satellite view of weather systems over the North p0063 A75-10790 American Arctic OPTICAL THICKNESS

Determination of the aerosol content from ERTS-1 data p0013 A75-17130

OPTICAL TRACKING

A simulation of the San Andreas fault experiment p0033 A75-11068

ORBIT CALCULATION

Transformation of terrestrial survey data to Doppler stellite datum p0054 A75-16569 satellite datum

ORBITAL POSITION ESTIMATION

Analytical estimates of precision in determining parameters of motion of artificial earth satellite when analyzing imprecise measurements with correlated errors p0053 A75-11419

ORBITAL WORKSHOPS

The final Skylab mission: Man at home and at work in space [NASA-TM-X-70394] p0078 N75-12019 OREGON

Evaluate ERTS imagery for mapping and detection of changes of snowcover on land and on glaciers --- Alaska and Washington

(E74-10812)

Study to develop improved spacecraft snow survey methods using Skylab/EREP data --- Sierra Nevada. Cascades, and Great Plains

[E75-10081] p0073 N75-14200

ORTHOPHOTOGRAPHY

An electronic drawing aid for map compiling --- from ereo-orthophotos p0027 A75-10076 stereo-orthophotos Recent progress in urban mapping p0015 A75-18054 Automatic contouring on the Gestalt Photomapper 2-1 p0028 A75-18055

Skylab induced environment --- spacecraft-originated contaminant effects [AIAA PAPER 74-1225]

Nimbus 4 observations of changes in total ozone and stratospheric temperatures during a sudde p0012 A75-14841

Measurements of the extraterrestrial solar radiant flux from 2981 to 4000 A and its transmission through the earth's atmosphere as it is affected by dust and ozone

Comparison of the Southern and Northern Hemisphere general circulation characteristics as determined by satellite [UCRL-75556] p0017 N75-10638

P

PACIFIC OCEAN

A simulation of the San Andreas fault experiment p0033 A75-11068

Monitoring and predicting El Nino invasions --- equatorial Pacific Ocean-atmosphere interactions

p0041 A75-12719

A remote sensing study of Pacific hurricane Ava p0012 A75-17112

Pacific range electromagnetic signature studies /Project PRESS/ p0042 A75-17479

Comparison of the Southern and Northern Hemisphere general circulation characteristics as determined by satellite ozone data [UCRL-75556] n0017 N75-10638

An assessment of the potential contributions to and Gulf of Mexico [E75-10017]

p0044 N75-12396

PANORAMIC CAMERAS

Geometric distortions of opticom television systems p0057 N75-11425 Use of a panoramic aerial camera in aerosurveying [AD-784162] n0071 N75-11450

PARTICLE DENSITY (CONCENTRATION)

Methods for the automatic recording of atmospheric dust concentrations and some of the results obtained p0057 N75-11502

PARTICLE SIZE DISTRIBUTION

A camera for photography of atmospheric particles from p0053 A75-13526 Aircraft measurements of the concentration and size distribution of atmospheric sulfate particles

p0057 N75-11504
Particulate sizes from polarization measurements --measured by ground based polarimeter

Lidar observations of atmospheric particulate content p0019 N75-11559

PARTICULATE SAMPLING

Observation and measurement of atmosph [WMO-368] VMO-368] p0018 N75-11476 A technique for measurement of sulphur dioxide and sulphate concentrations from an aircraft

p0018 N75-11498

PASSIVE L-BAND RADIOMETERS Detection of moisture and moisture related phenomena from Skylab --- radiometer and scatterometer analysis of

Texas site p0007 N75-13333 [E75-10007] Detection of moisture and moisture related phenomena from Skylab --- Texas and Kansas

p0008 N75-14192 PATTERN RECOGNITION

Mapping soil features from multispectral scanner data

p0027 A75-15435

Cloud screening from aerial photography applying coherent optical pattern recognition techn p0054 A75-16690

Implementation of an advanced table look-up classifier for large area land-use classification p0014 A75-17153

_Texture analysis with Fourier series --- for crop identification from color aerial photographs

p0004 A75-17187 Multi-aspect techniques in remote sensing --- for MSS agery p0068 A75-17217 Automatic pattern recognition and mapping utilizing ermal infrared imagery p0070 A75-18074 The application of pattern Gressons Skylark earth resources rocket imagery p0056 A75-19268

Pattern recognition relevant to reconnaissance and remote sensing --- noting classification methods
[PHL-1974-23] p0070 N

p0070 N75-10573 An extension of the combinatorial geometry technique for modeling vegetation and terrain features
[AD-782883] p000

p0006 N75-10577 Use of feature extraction techniques for the texture and context information in ERTS imagery: Spectral and textural processing of ERTS imagery --- classification of Kansas p0059 N75-13335

Planning applications in east central Florida --- maximum likelihood classification programming

[E75-10015] n0021 N75-13337 PENNSYLVANIA

Forest defoliation assessment with satellite imagen p0003 A75-17171 Urban and regional land use analysis: CARETS and

census cities experiment package p0020 N75-12398 [E75-10019] Remote sens ing geophysics from Skylab --- rationing-type

algorithms to enhance multispectral data p0030 N75-13352 The remote identification of terrain features and materials

at Pennsylvania test sites: An investigative study of

[PB-235992/5] p0031 N75-14263 Urban and regional land use analysis: CARETS and ensus cities experiment package --- New Haven. census cities experiment package --- New Haven, Connecticut; Pontiac, Michigan; Phoenix, Arizona; District

[E75-10091] p0024 N75-15109 Urban and regional land use analysis: CARETS and census cities experiment package

[F75-10100] p0025 N75-15118 PERFORMANCE PREDICTION

Single flight stereo radar capabilities --- for terrain

p0028 A75-17201 PERFORMANCE TESTS

Water depth penetration film test p0047 A75-12873 PHASED ARRAYS

Performance of a hologram matrix radar

p0065 A75-14792

Cartographic evaluation of Skylab-A S-192 scanner images --- Chesapeake Bay, Phoenix, and San Francisco

[E75-10078] Urban and regional land use analysis: CARETS and census cities experiment package --- New Haven, Connecticut; Pontiac, Michigan; Phoenix, Arizona; District

of Columbia [E75-10091] p0024 N75-15109 **PHOTOGEOLÓGY**

Plutonic structure of the earth's crust on space images p0027 A75-17125 Skylab S-192 ratio codes of soil, mineral, and rock spectra

for ratio image selection and interpretation p0003 A75-17158

Optical data processing analysis of stream patterns chibited on ERTS-1 imagery p0056 A75-17219 exhibited on ERTS-1 imagery Geological and geophysical studies from space and prospects for their development p0035 N75-11436

Experience in integrated geological and geographical interpretation of space photographs obtained in the USSR p0036 N75-11437

Geologic and mineral and water resources investigations in Western Colorado, using Skylab EREP data --[E75-10043] p0058 N75-12412 Geologic and mineral and water resources investigations

western Colorado, using Skylab EREP data [E75-10071] p0037 N75-14190 Geologic evaluation of central Italy from ERTS-1 and

Skylab data p0038 N75-14238 The Great Basin investigation --- photomapping lithology

and geological structures [E75-10102] p0038 N75-15120 Geologic and mineral and water resources investigations

p0025 N75-15123 [E75-10105]

in western Colorado, using Skylab EREP data

PHOTOGRAMMETRY

Coherent optics potential applications to mapping p0027 A75-14683

Coherent optics in mapping: Proceedings of the Tutorial Seminar and Technology Utilization Program, Rochester, N.Y., March 27-29, 1974 p0027 A75-16676

American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings p0056 A75-18052

Geographic correlation of television pictures obtained p0071 N75-11432 from weather satellites Computer and photogrammetric general land use study

of central north Alabama [NASA-TR-R-431] p0022 N75-13372

Work dealing with densification of altimetric-planometric data points employing universal instruments and radio altimeter and statoscopic indications --- for land improvement

[AD-785877] PHOTOGRAPHIC FILM

netration film test p0047 A75-12873

p0061 N75-14257

Recent progress in urban mapping p0015 A75-18054 PHOTOSENSITIVITY Optical film density values from color IR photography p0005 A75-18073 Water depth penetration film test p0047 A75-12873 stland soils mapping Automatic contouring on the Gestalt Photomapper 2 p0028 A75-18055 PHOTOGRAPHIC MEASUREMENT PHYSICAL PROPERTIES Terrain measurement from holographi Remote sensing and land use inventory and mapping Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical p0015 A75-18057 The application of ERTS-derived information to the p0001 A75-16682 The application or convergence pools A75-1800b geometronics-earth resource measurements and displays pools A75-18063 pools A75-18063 nd thermal characteristics of plants and soil PHOTOGRAPHIC PROCESSING p0007 N75-12403 Evaluation of Skylab imagery as an information service for investigating land use and natural resources --- Long PHYSIOLOGICAL EFFECTS The final Skylab mission: Man at home and at work in Island, New York p0034 A75-18065 Automatic mapping of strip mine [E74-10730] n0056 N75-10526 spacecraft data [NASA-TM-X-70394] Developing processing techniques for Skylab data --regetation and soil identification Trends in the use of remote sensing for state resources at environmental management p0005 A75-18068 PLANETARY ATMOSPHERES and environmental management [E74-10799] p0057 N75-10527 Methods of calculating infrared transfer: A review Application of ERTS data to environmental inventory and p0019 N75-11560 in planetary atmospheres PHOTOGRAPHIC RECORDING p0016 A75-18069 Water depth penetration film test p0047 A75-12873 PLANETARY GRAVITATION Land use mapping using ERTS-1 imagery for The Four Use of terrestrial gravity data in combination with satellite **PHOTOGRAPHS** Rivers Resource Conservation and Development District Catalog of earthquake photographs --- isoseismal maps results for determining good heights in Central Europe p0029 A75-18128 p0072 N75-13455 Overall evaluation of ERTS-1 imagery for cartographic PHOTOINTERPRETATION application --- Australia, Antarctica, U.S., and U.S.S.R. [E74-10800] p0029 N75-10528 PLANETARY RADIATION The radio emission of the earth as a A comparison of Skylab and ERTS data for agricultural Geographic applications of ERTS-1 imagery to landscape change --- Mississippi River and Great Smoky Mountains p0076 A75-15319 crop and natural vegetation interpretation
[AIAA PAPER 74-1219] p0001 A75-12033 book PLANETARY SURFACES of Tennessee and North Carolina Space iconics Special color analysis techniques --- for reconnaissand [E74-10810] p0034 N75-10537 INASA-TT-F-798 p0057 N75-11419 sensor imagery p0064 A75-12874 Evaluate ERTS imagery for mapping and detection of Environmental studies of Iceland with ERTS-1 imagery p0012 A75-17105
Classification and mapping of coal refuse, vegetative PLANTS (BOTANY) changes of snowcover on land and on glaciers --- Alaska A study to explore the use of orbital remote sensing to and Washington determine native arid plant distribution --- Arizona Regional [E74-10812] p0049 N75-10539 cological Test Site cover types, and forest types by digital processing ERTS-1 p0054 A75-17110 Skylab-EREP investigations of wetlands ecology --- New p0006 N75-10544 [E74-10818] Jersey and Chesapeake Bay Areal extent of snow in forested regions - A practical Develop techniques and procedures, using multispectral [E74-10816] n0049 N75-10542 estimation technique using ERTS-1 data systems, to identify from remotely sensed data the physical Utilization of Skylab (EREP) system for appraising p0002 A75-17122 nd thermal characteristics of plants and soil --- Southern changes in continental migratory bird habitat Moisture detection from Skylab --- soil studies for Texas [E74-10817] p0002 A75-17145 p0006 N75-10543 p0006 N75-12394 and Kansas sites [E75-10012] Automated strip mine and reclamation mapping from p0035 N75-10556 Modifications in a computer-implemented method for the Develop techniques and procedures, using multispectral ERTS --- Ohio coal detection and extraction of objects in aerial photographs systems, to identify from remotely sensed data the physical Land use and mapping p0017 N75-10560 nd thermal characteristics of plants and soil p0055 A75 Correlation of missions 191, 51M and helicopter [E75-10033] p0007 N75-12403 Lake Ontario water mass delineation from ERTS-1 photography --- aerial photography and mapping of three areas in Test Site 175 in Texas p0042 A75-17184
The interpretation and use of false-colour infra-red and A study of the early detection of insect infestations and density/distribution of host plants p0058 N75-12279 [NASA-CR-134289] p0008 N75-13362 true colour photography of part of Argentina obtained by [E75-10062] Mapping exposed silicate rock types and exposed ferric Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical Skylark earth resources rockets An intensive natural resources inventory aids land use planning and resources management in Puerto Rico and ferrous compounds from a space platform p0036 N75-12392 [E75-10008] and thermal characteristics of plants and soil
[E75-10077] p0008 Hydrologic significance of lineaments in central p0005 A75-18056 p0008 N75-14196 Early results from the experimental Automated Image Data Extraction System --- for aerial terrain sensing PLASMA INTERACTIONS SMA INTERACTIONS
Solar wind and geomagnetic activity
p0027 A75-11975 p0050 N75-12401 [E75-10030] p0050 N75-12401 The ERTS-1 investigation. Volume 7: Erts-1 land-use p0069 A75-18072 alysis of the Houston area test site ERTS-1 imagery interpretation techniques in the Tennessee Valley --- land use and soil mapping [E74-10811] p0016 N75-10538 PLOTTERS [E75-10009] p0021 N75-13334 An electronic drawing aid for map compiling --- from erec-orthophotos p0027 A75-10076
The GLOBAL computer program and GLOBEMAP: A Mapping of the major structures of the African rift system [E74-10811] stereo-orthophotos Ethiopia, Somalia, and Yemen Experience in integrated geological and geographical p0030 N75-13340 [E75-10024] interpretation of space photographs obtained in special purpose three-dimensional plotter subroutine [AD-785869] p0032 N75-1 Applications of Skylab EREP photographs to mapping of landforms and environmental geology in the Great Plains p0036 N75-11437 p0032 N75-15141 Geologic and mineral and water resources investigations in Western Colorado, using Skylab EREP data --- Utah and Midwest --- Illinois, Iowa, Kansas, Missouri, Nebraska, from power plant p0014 A75-17162 Scanning thermal plumes --- from and South Dakota 75-10043] p0058 N75-12412 Meteorological interpretation of high resolution images p0037 N75-13355 [E75-10055] Dynamics of suspended sediment plumes in Lake Preliminary results in the comparison of Skylab, ERTS and RB-57 images for the detection of lineaments and for remote sensing of earth resources p0060 N75-14229 Interpretation of ERTS-1 data for investigating the French [E74-10815] p0049 N75-10541 fractures in Precambrian, Paleozoic, and late Tertiary rocks on and near the Colorado Plateau, North-Central Arizona POLAR CAPS western coastal zones (Fralit program) p0051 N75-14230 On the determination of secular variation in the earth's [E75-10060] p0037 N75-13360 Multispectral photography for earth regravity field and of secular polar motion Study to develop improved spacecraft snow survey methods using Skylab/EREP data --- Sierra Nevada, p0074 N75-15138 DO029 A75-18125 [NASA-CR-141283] PHOTOMAPPING POLAR SUBSTORMS Cascades, and Great Plains [E75-10081] On the longitudinal localization of the substorm active An electronic drawing aid for map compiling ---75-10081] p0073 N75-14200 The application of Skylab EREP data for land use stereo-orthophotos p0027 A75-10076 region and its changes during the substo p0027 A75-11861 Hologrammetry - Concepts and applications classification and mapping in the Cleveland and Columbus p0065 A75-15434
Coherent optics in mapping; Proceedings of the Tutorial POLARIMETERS Remote measurements of water pollution with a lider plarimeter p0014 A75-17205 [E75-10083] p0023 N75-14201 Seminar and Technology Utilization Program, Rochester, N.Y., March 27-29, 1974 P0027 A75-18676 Methodological aspects and regional examples of snow-cover mapping from ERTS-1 and EREP imagery of POLARIMETRY Environmental studies of Iceland with ERTS-1 imagery Remote detection of soil surface moisture Classification and mapping of coal refuse, vegetative cover types, and forest types by digital processing ERTS-1 data p0051 N75-14232 p0003 A75-17146
The application of spectrometric and polarization Investigation of accuracy of determination of plane coordinates by means of the Central Scientific Research techniques for remote sensing of oil on sea water p0068 A75-17210 p0054 A75-17110 Institute of Geodesy, aerial photography and cartography POLARIZATION (WAVES) Transference of ERTS-1 spectral signatures in time and p0061 N75-14255 ace p0068 A75-17111
Interactive computer processing for land use planning [AD-785886] A study of a dual polarization laser backscatter system for remote identification and measurement of water Morphographic and morphometric analysis of vegetative DO054 A75-17134 cover photographs in aerial photography for the mapping pollution POLARIZATION CHARACTERISTICS
Particulate circulates Land type analysis for regional land use planning from p0009 N75-14258 photomorphic mapping - An example for Boulder County [AD-785932] p0013 A75-17137 Particulate sizes from polarization measurements --measured by ground based polarimeter Usefulness of Skylab color photography and ERTS-1 Colorado
Application of remote sensing to the location of hydrologically active /source/ areas p0066 A75-17141 SICLOPS - A system of computer programs for rectified mapping of airborne scanner imagery p0055 A75-17156 Microwave radiometric characteristics of snow-covered terrain multispectral imagery for mapping range vegetation types tern Wyoming nO019 N75-11558 POLLUTION CONTROL p0010 N75-15115 [E75-10097] Research in earth resources utilization, physical The Great Basin investigation --- photomapping lithology ineering, and pollution and geological structures [AR-29] p0022 N75-13369 p0038 N75-15120 POLLUTION MONITORING Summary of flight performance of the Skylab Earth An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using A microwave correlation spectrometer for air pollution politoring p0063 A75-11732 Resources Experiment Package / EREP/ p0076 A75-17211 monitoring p0063 A75-11732
On evaluating compliance with air pollution levels 'not automatic data processing techniques Statistical separability of agricultural cover types in [E75-10104] p0073 N75-15122 to be exceeded more than once a year subsets of one to twelve spectral channels Use of forest vegetation as an indicator in the interpretation and large scale mapping of soils in a forest p0011 A75-12803 Trace gas analysis of power plant plumes via aircraft p0005 A75-17216 rface compositional mapping in the Wind River Range measurement - 03, NOx, and SO2 chemistry p0064 A75-13148 and Basin, Wyoming by multispectral techniques applied [AD-785967] p0010 N75-15146 to ERTS-1 data p0028 A75-17224 Applications of remote sensing by the state of p0069 A75-17232 **PHOTOMETRY** Airborne survey of major air basins in California p0011 A75-14380 Special color analysis techniques --- for reconnaissan p0064 A75-12874 attidisciplinary applications of ERTS and Skylab data sensor imagery Determination of the aerosol content in

Methods for the automatic recording of atmospheric dust concentrations and some of the results obtained

p0057 N75-11502

in Ohio

p0015 A75-17234

A summary of ERTS data applications in Alaska p0077 A75-17236 p0013 A75-17130

p0048 A75-17161

from ERTS-1 data

Remote sensing of western Lake Se

Scanning thermal plumes --- from PROJECT PLANNING power plant Investigation of radar discrimination of sea ice p0014 A75-17162 The next problems in the area of the implementation of p0042 A75-17121 The economic impact of remote sensing data as the On the feasibility of remote monitoring of soil moisture ing for cartographical-geodetic work p0032 N75-15143 p0067 A75-17147 source of nonpoint pollution monitoring and control [AD-785882] microwave sensors Radar response to vegetation --- soil moisture mapping a microwave backscattering p0005 A75-18556 p0076 A75-17166 PROJECTORS Remote measurements of water pollution with a lidar polarimeter p0014 A75-17205 The ITC 8-image comparator and its possible use in ERTS and EREP investigations p0060 N75-14222 via microwave backscattering p0005 A75-18556
Preliminary analysis of Skylab RADSCAT results over polarimeter Development of an experimental airborne laser remote the ocean --- using radar backscatter and microwave **PUERTO RICO** sensing system for the detection and classification of oil An intensive natural resources inventory aids land use planning and resources management in Puerto Rico p0068 A75-17206 p0046 N75-14793 Oil slick detection by X-band synthetic aperture radar p0005 A75-18056 RADAR SIGNATURES p0068 A75-17209 Pacific range electromagnetic signature studies /Project RESS/ p0042 A75-17479 PULSE GENERATORS Conference on Environmental Sensors and Applications, PRESS/ A generator of high-power current pulses of semisinusoidal shape for an aerial variant of the method of transient processes --- for aerial prospecting London, England, November 18, 19, 1974, Proceedings RADAR TRANSMISSION p0070 A75-19265 A study of a dual polarization laser backscatter system Bistatic rader sea state monitoring [NASA-CR-137469] p0043 N75-10673 p0053 A75-13310 for remote identification and measurement of water RADIANCE **PYRANOMETERS** Ground-based determination of atmospheric radiance for Ground-based measurements of solar radiation by [NASA-CR-140665] n0017 N75-10563 correction of ERTS-1 data p0054 A75-16476
Measurement of trace stratospheric gases by inverting geophysical monitoring for climatic change (USA) --- using Application of remote sensing for prediction and detection oheliometers, pyranometers, and radiometers of thermal pollution thermal infrared limb radiance profiles p0018 N75-11527 [NASA-CR-139182] p0017 N75-10572 Feasibility study of a Gunn-diode microwave cavity PYRENEES MOUNTAINS (EUROPE) RADIATION DOSAGE ARNICA/ERTS-1 experiment. Results and perspectives ctrometer as a formaldehyde monitor ARNICA/ER15-1 experiment. Results of the Pyrenees pools 175-14246 Atmospheric release advisory capability: Research and p0071 N75-11311 rogress [UCRL-75644-REV-2] Observation and measurement of atmospheric pollution n0025 N75, 15202 p0018 N75-11476 [WMO-368] **PYROHELIOMETERS** RADIATIVE TRANSFER Aircraft measurements of the concentration and size Proceedings of the UCLA International Conference on Ground-based measurements of solar radiation by geophysical monitoring for climatic change (USA) --- using pyroheliometers, pyranometers, and radiometers distribution of atmospheric sulfate particles Radiation and Remote Probing of the Atmosphere emphasizing radiative transfer [NASA-CR-140816] p0057 N75-11504 diometers p0018 N75-11527 Correlation spectroscopy techniques for the regional and global measurements of air pollution p0018 N75-11505
Global measurements of air pollution from satellites ... p0078 N75-11549 Solution of the radiative transfer pooss N75-11551 Monte Carlo method p0018 N75-11521 nploying radiometer techniques Radiative transfer of visible radiation in turbid p0019 N75-11552 hploying radiometer techniques
Microwave rotational spectroscopy: A physical technique
receific collutant monitoring p0018 N75-11529 for specific pollutant monitoring Some analytical results for radiative transfer in thick Measurements of meteorological parameters and air pollution spread over Stockholm, Sweden, by means of QUALITY CONTROL atmospheres --- of monochromatic radiation Quality control in global and regional air chemistry p0019 N75-11555 remote sensing and airborne instrumentati monitoring networks: Lessons of the European Air Methods of calculating infrared tra nsfer: A review ---p0019 N75-11560 p0018 N75-11540 p0018 N75-11531 Chemistry Network n planetary atmospheres Airborne vapor surveillance system QUANTITATIVE ANALYSIS RADIO ALTIMETERS [AD-785585] p0021 N75-12459 Quantitative determination of stratospheric aerosol Skylab altimeter applications and scientific results Oil pollution detection, monitoring and law enforcement p0063 A75-12035 characteristics [AIAA PAPER 74-1221] Skylab radar altimeter - Short-wavelength perturbations etected in ocean surface profiles p0066 A75-16096 isiana Gulf Coast waters [E74-10808] n0016 N75-10535 p0022 N75-13343 QUASARS detected in ocean surface profiles The economic value of remote sensing of earth resources Geodetic aspects of laser distance measurements to the A radar altimeter for satellite application from space: An ERTS overview and the value of continuity of service. Volume 8: Atmosphere [NASA-CR-141191] p0080 N75-14213 moon and radio-interference measurements to quasars [AD-784373] p0056 N75-10425 DO027 A75-14646 Work dealing with densification of altimetric-planometric data points employing universal instruments and radio PONTIAC (MI) altimeter and statoscopic indications --- for Urban and regional land use analysis: CARETS and R census cities experiment package --- New Haven. Connecticut; Pontiac, Michigan; Phoenix, Arizona; District [AD-785877] p0061 N75-14257 RADIO EMISSION of Columbia RADAR CROSS SECTIONS [E75-10091] p0024 N75-15109 Skylab altimeter applica [AIAA PAPER 74-1221] lications and scientific res The earth as a radio source - Terrestrial kilometric p0063 A75-12035 p0011 A75-11211 POPULATIONS planet --- Russian Estimation of population density in Tokyo districts from The radio emission of the earth Radar cross sections of vegetation canopies determined ERTS-1 data p0076 A75-17103 p0076 A75-15319 by monostatic and bistatic scatterometry p0005 A75-17190 POSITION (LOCATION) RADIO FREQUENCY INTERFERENCE Transformation of terrestrial survey data to Doppler BADAR FOUIDMENT Geodetic aspects of laser distance measurements to the p0054 A75-16569 satellite datum Skylab radar altimeter - Short-wavelength perturba moon and radio-interference measure ments to quasars p0027 A75-14646 p0066 A75-16096 POSITION ERRORS detected in ocean surface profiles RADAR IMAGERY A simulation of the San Andreas fault RADIO METEOROLOGY Simultaneous active and passive microwave response of the earth - The Skylab radscat experiment p0033 A75-11068 Performance comparison of techniques for obtaini stereo radar images p0063 A75-100
NASA-KSC's earth resources benefits from spa n0063 A75-10075 Satellite techniques for the study p0012 A75-17114 p0064 A75-12980 in position RADIO PROBING exploration p0075 A75-13592 POTABLE WATER Performance of a hologram matrix radar Microwave emission characteristics of moist soils Can spinoffs from space provide water for a thirsty p0065 A75-14792 p0047 A75-13578 n0001 A75-15180 Extraction of urban land cover data from multiplexed inthetic aperture radar imagery p0054 A75-17117

Development of research in the USSR on the use of RADIO TRANSMISSION POWER SPECTRA synthetic aperture radar imagery Cloud screening from aerial photography applying vestigations of the ionosphere by space technic p0065 A75-13868 coherent optical pattern recognition technique radar images for geological purposes p0033 A75-17126 Investigation of microwave hologram techniques for p0054 A75-16690 PRECAMBRIAN PERIOD Very high resolution radiometer p0064 A75-12109 application to earth resources n0067 A75-17195 Global measurements of air pollution from satellites --nploying radiometer techniques p0018 N75-11521 minary results in the comparison of Skylab, ERTS Single flight stereo radar capabilities --- for terrain ralysis p0028 A75-17201 and RB-57 images for the detection of lineaments and employing radiometer techniques fractures in Precambrian, Paleozoic, and late Tertiary rocks on and near the Colorado Plateau. North-Central Arizona Measurement of sea surface currents using airborne Doppler radar and inertial navigation systems p0042 A75-17203 Ground-based measurements of solar radiation by ysical monitoring for climatic change (USA) --- using [E75-10060] p0037 N75-13360 pyroheliometers, pyranometers, and radiometers PRECIPITATION (METEOROLOGY) Oil slick detection by X-band synthetic aperture radar p0068 A75-17209 n0018 N75-11527 A camera for photography of atmosph Recordings of the spectral-radio-meter SIGNE during the D0053 A75-13526 aircraft Application of radar imagery to e ummer of 1972 p0034 A75-17222 Geographic applications of ERTS-1 imagery to landscape [FOA-2-C-2577-E1] p0072 N75-12283 change --- Mississippi River and Great Smoky Mountains of Tennessee and North Carolina [E74-10810] p0034 N75-10537 Small scale thematic mapping - A case for radar Experimental evaluation of atmospheric effects on p0028 A75-18062 radiometric measurements using the EREP of Skylab ... A simulation of synthetic aperture radar imaging of ocean Bonneville Salt Flats, Utah PREDICTION ANALYSIS TECHNIQUES [E75-10036] p0020 N75-12406 [NASA-TM-X-72629] p0044 N75-12172 Application of remote sensing for prediction and detection Meteorological data. Volume 145 no. 1: The European climatology 1974 meteorological satellite data from the of thermal pollution [NASA-CR-139182] RADAR MAPS Feasibility of using multiplexed SLAR imagery for water p0017 N75-10572 Berlin zone of reception, part 1, 1st quarter resources management and mapping vegetation communities n0054 A75-17115 p0023 N75-13479 From aerospace photographs to forecastin, and wetness, conducted in the USSR, in 1974
[NASA-TT-F-16151] calculating flows
[NASA-TT-F-16006] Small scale thematic mapping p0059 N75-13367 p0028 A75-18062 Atmospheric release advisory capability: Research and RADAR MEASUREMENT RAIN progress [UCRL-75644-REV-2] The wind vector in a layer over water p0025 N75-15202 Rainfall estimations from geosynchronous satellite p0041 A75-15179 p0047 A75-13596 PROJECT MANAGEMENT of sea ice p0042 A75-17121 imagery Investigation of radar discrimination of RAMAN SPECTROSCOPY Planning_and_management, of_science programs on-The remote Raman spectrometer is a viable instrument for remote sensing of the environment The role of lidar in environmental sensing --- laser ceilometer and visual range measurement application [AIAA PAPER 74-1215] p0075 A75-12030 p0068 A75-17214

n0070 A75-19274

p0001 A75-16242

Laser Raman spectroscopy of solutes dissolved in water

p0056 N75-10202

from a remote platform [PB-233432/4]

An overview of Texas activities in remote sensing

Pacific range electromagnetic signature studies / Project

p0076 A75-17235

p0042 A75-17479

RADAR SCATTERING

Active microwave measurement of soil water content [NASA-CR-141685] p0001 A75-16242

control of terrestrial resources

p0053 A75-13758

[IAF PAPER 74-140]

RANGE FINDERS The potential benefits for development derived from space Remote detection of soil surface moisture Investigation of accuracy of determination of plane coordinates by means of the Central Scientific Research Institute of Geodesy, aerial photography and cartography p0003 A75-17146 [IAF PAPER 74-147] n0075 A75-13761 On the feasibility of remote monitoring of soil moisture ith microwave sensors p0067 A75-17147 Spectrophotometric observations 'Soyuz-13' earth with microwave sensors airborne rangefinder Operational use of satellite and high altitude remote IJAF PAPER 74-156) n0065 A75-13765 [AD.785886] o0061 N75-14255 sensing for the generation of input data for water demand models p0067 A75-17148 Economic effectiveness of using the space systems of RANGE RESOURCES Oetermining range condition and forage production potential in California from ERTS-1 imagery distant indication of natural resources Interactive machine assessment of critical land resources n0075 A75-13770 [IAF PAPER 74-165] using ERTS-1 data Mathematical modeling for the detection of fish by an p0013 A75-17149 n0003 A75-17169 Skylab S-192 ratio codes of soil, mineral, and rock spectra airborne laser Agriculture, forestry, range resources for ratio image selection and interpretation A radio picture of the earth --- microv wave remote sensing p0066 A75-16191 p0006 N75-10558 p0003 A75-17158 from satellites Satellite pictures can help quide Sandhills irrigation ---Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites Coastal and estuarine studies with ERTS-1 and Skylab p0047 A75-16240
Active microwave measurement of soil water content p0001 A75-16242 center-pivot irrigation in Sand Hills Region, Nebraska [E75-10023] p0021 N75-13 p0021 N75-13339 p0048 A75-17160 Usefulness of Skylab color photography and ERTS-1 [NASA-CR-141685] Remote sensing of western Lake Superio multispectral imagery for mapping range vegetation types in southwestern Wyoming Ground-based determination of atmospheric radiance for correction of ERTS-1 data p0054 A75-16476 p0048 A75-17161 Scanning thermal plumes --from power plant p0014 A75-17162 [E75-10097] n0010 N75-15115 p0054 A75-16476 ng with an airborne laser p0066 A75-16485 on Remote Sensing of condensers RANGELANDS absorption spectrometer The use of remote sensing in limnological studies Inventory of forest and rangeland resources, including International Symposium eutrophication of lakes p0048 A75-17163 Environment, 9th, University of Michigan, Ann Arbor, Mich., April 15-19, 1974, Proceedings, Volumes 1, 2 & 3 forest stress --- Colorado, Georgia, and Black Hills, South Correlation of multispectral imagery with water analysis
The Ross Barnett Reservoir Remote Sensing Project
p0014 A75-17165 [E75-10032] n0007 N75-12402 p0066 A75-17101 nment of Tokyo by Use of ERTS-1 imagery to interpret wind-erosion hazard Some features of the urban enviro The economic impact of remote sensing data as the remote sensing ... infrared surface temperature reasurements pool 2 A75-17102
Estimation of population density in Tokyo districts from pO076 A75-17103
An examination of the extent of fire in the grassland and savanna of Africa along the southern side of the Sahara pool A75-17104
A method of specifying remotely sensed units for soil in the Sandhills of Nebraska source of nonpoint pollution monitoring and control [E75-10044] o0059 N75-13347 p0076 A75-17166 Preliminary results of fisheries investigation associated p. Obulah-3 p.0003 A75-17167 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland with Skylab-3 MID Skylab-3
Rapid stock assessment of pilchard populations by aircraft-borne remote sensors p0003 A75-17168
Determining range condition and forage production potential in California from ERTS-1 imagery [NASA-CR-141263] DO080 N75-14208 REAL TIME OPERATION A method of specifying remotely sensed units for soil Atmospheric release advisory capability: Research and sample points mple points
Geologic interpretation of ERTS-1 satellite images for p0033 A75-17108 n0003 A75-17169 progress Grass canopy bidirectional spectral reflectance p0003 A75-17170 UCRL-75644-REV-2] p0025 N75-15202 East Aswan area, Egypt RECLAMATION Geologic interpretation of ERTS-1 satellite images for lest Aswan area, Egypt p0033 A75-17109 Forest defoliation assessment with satellite imagery p0003 A75-17171 tromated strip mine and reclamation mapping from 5 --- Ohio coal p0035 N75-10556 West Aswan area, Egypt ERTS -- Ohio coal Classification and mapping of coal refuse, vegetative Applicability of Skylab remote sensing for detection and Spectral reflectance studies on mineral deficiency in corn cover types, and forest types by digital processing ERTS-1
data p0054 A75-17110 p0003 A75-17172 monitoring of surface mining activities --- strip mining and reclamation in Kentucky and Tennessee data Agricultural resources investigations in Southern France Transference of ERTS-1 spectral signatures in time and pace p0066 A75-17111 Agricultural resources investigations in Souriem related and Northern litaly p0004 A75-17173
Thermal behaviour of some rice fields affected by a yellows-type disease p0004 A75-17174
First results from the crop identification technology D0044 N75-12407 [E75-10038] Applicability of Skylab remote sensing for detection and A remote sensing study of Pacific hurricane Ava p0012 A75-17112 monitoring of surface mining activities — evironmental effects and reclamation in Tennessee and Kentucky Simultaneous active and passive microwave response assessment for remote sensing / CITARS/ [E75-10085] p0038 N75-15103 of the earth - The Skylab radscat experiment sessment for remote sensing /CITARS/ p0004 A75-17175 ERTS-I data for classifying native plant communities -entral Colorado p0004 A75-17176 p0012 A75-17114 RECREATION Feasibility of using multiplexed SLAR imagery for water resources management and mapping vegetation communities p0054 A75-17115 Study of recreational land and open space using Skylab Central Colorado imagery [575-10020] Computer derived coastal water classifications n0020 N75-12399 communities p0042 A75-17177 spectral signatures Four channel simultaneous X-L band imaging SAR p0054 A75-17116 Study of recreational land and open space using Skylab California nearshore processes - ERTS 1 --- coastal - southeast Michigan p0048 A75-17179 currents and sediments [E75-10034] Land-use planning aided by computer cellular DO020 N75-12404 Surface currents along the California coast observed on Study of recreational land and open space using Skylab imagery --- snow and ice hydrology of southeast Michigan modelling/mapping system to combine remote sensing, p0048 A75-17180 ERTS imagery natural resources, social and economic data Ocean internal waves off the North American and African n0076 A75-17118 passis from ERTS-1 p0042 A75-17183 Investigation into the spectral signature of agricultural ops during their state of growth p0004 A75-17189 Signatures of various earth surfaces measured by the mbus-5 microwave spectrometer p0066 A75-17119 coasts from ERTS-1 Signatures of various earm successions properties of sea ice - p0041 A75-17120 [E75-10092] p0024 N75-15110 REDUNDANCY crops during their state of growth Photographs from balloons - Their use in agronomy and management of environment p0055 A75-17191 Reduction of information redundancy in ERTS-1 and EREP data. A proposal for an experimental study Investigation of radar discrimination The interpretation and use of false-colour infra-red and p0060 N75-14225 p0042 A75-17121 e colour photography of part of Argentina obtained by ylark earth resources rockets p0067 A75-17192 REGIONAL PLANNING Areal extent of snow in forested regions - A practical Skylark earth resources rockets ERTS-1 and data for national land use planning estimation technique using ERTS-1 data A preliminary feasibility study on a synthetic aperture radar /SARSAT/ for earth resources surveys p0067 A75-17194 p0075 A75-13581 p0002 A75-17122 The application of ERTS-derived information to the ERTS applications in Thailand - A p regional land use planning process p0015 A75-18058 p0076 A75-17123 Investigation of microwave hologram application to earth resources am techniques for p0067 A75;17195 Application of remote sensing to state and regional problems --- for Mississippi [NASA-CR-140805] p0078 N75-11414 · Passive microwave sensing of moist soils p0066 A75-17124 Deteriorating effects on 3 mm wave passive imagen Development of research in the USSR on the use of p0055 A75-17198 radar images for geological purposes p0033 A75-17126 Urban and regional land use analysis: CARETS and The usefulness of imaging passive microwave for rural p0055 A75-17199 On the detectability of atmospheric carbon monoxide by icrowave remote sensing p0013 A75-17129 census cities experiment package nd urban terrain analysis [E75-10019] Microwave radiometric characteristics of snow-covered p0020 N75-12398 microwave remote sensing Utilization of EREP data in geological evaluation regional On the natural limitations of target differentiation by terrain planning, forest management, and water management in North Carolina Single flight stereo radar capabilities --- for terrain alysis p0028 A75-17201 means of spectral discrimination techniques --- ERTS remote p0054 A75-17132 analysis [E75-10063] Electrically scanning microwave radiometers -- for tellite-borne remote sensing p0056 A75-17202 p0008 N75-13363 Interactive computer processing for land use planning RELIEF MAPS p0054 A75-17134 satellite-borne remote sensing The calculation of cloud shadows in modelling of the orth's surface survey from space p0014 A75-17204 Coherent optics potential applications to mapping Use of ERTS-1 imagery for land evaluation in Pennington punty, South Dakota p0055 A75-17135 p0027 A75-14683 earth's surface survey from space County, South Dakota Remote measurements of water pollution with a lidar plarimeter p0014 A75-17205

Development of an experimental airborne laser remote Automatic contouring on the Gestalt Photomapper 2-1 Land type analysis for regional land use planning from p0028 A75-18055 photomorphic mapping - An example for Boulder County, REMOTE SENSORS p0013 A75-17137 sensing system for the detection and classification of oil A technology to renovate the search for new mineral Flood inundation in the southeastern United States from p0063 A75-10434 deposits p0047 A75-17138 Crude and refined petroleum oil structured luminescence Automated image processing in forestry aircraft and satellite imagery p0053 A75-10435 Mapping of the 1973 Mississippi River floods by the OAA-2 satellite p0028 A75-17139 signatures induced by UV laser or lamp and their remote sensing applications p0033 A75-17207 NOAA-2 satellite sensing applications More information and less data from space --- data Multi-frequency radiometric measurements of foam and a mono-molecular slick --- microwave sensing of ocean surface p0068 A75-17208 p0063 A75-11730 reduction in remote sensing systems p0063 A75-11730
Automatic land use classification using Skylab S-192 A hydrogeomorphic approach to evaluating flood potential in central Texas from orbital and suborbital remote sensing multispectral data p0047 A75-17140 imagery p0064 A75-12036 p0064 A75-12109 Oil slick detection by X-band synthetic aperture radar p0068 A75-17209 [AIAA PAPER 74-1224] Application of remote sensing to the location of drologically active /source/ areas p0066 A75-17141 Very high resolution radiometer hydrologically active /source/ areas The application of spectrometric and polarization Special color analysis techniques -for reconnaissance The use of remote sensing and natural indicators to p0064 A75-12874 techniques for remote sensing of oil on sea water sensor imagery delineate floodolains - Preliminary findings Automated remote measurement system in agricultural meteorology --- for crop survey by aerial photometry p0064 A75-12947 p0068 A75-17210 p0067 A75-17142 Summary of flight performance of the Skylab Earth Resources Experiment Package / EREP/ Evaluation of ERTS-1 and aircraft data for assessing internal drainage in irrigated agriculture p0076 A75-17211 Skylab earth resources program p0011 A75-13577 p0002 A75-17143 nner --- MSS design p0068 A75-17213 A motion-compensated spatial scanner Geoscientific research with rockets a d satellites Use of visible, near-infrared, and thermal infrared remote (IAF PAPER 74-116) p0053 A75-13741 sensing to study soil moisture p0002 A75-17144 Orientation of methods of information processing for the The remote Raman spectrometer is a viable instrument

soil studies for Texas

p0002 A75-17145

Moisture detection from Skylab

p0068 A75-17214

for remote sensing of the environment

Geologic interpretation of infrared thermal images in East Qatrani area, Western Desert, Egypt p0033 A75-17215 ... for MSS Multi-aspect techniques in remote se sensing --- for MSS p0068 A75-17217 Geothermal reconnaissance from quantitative analysis of ermal infrared images p0056 A75-17218 thermal infrared images Remote sensing of rock type in the visible and p0033 A75-17220 Prediction of the Fraunhofer line detectivity of luminescent materials --- crude and refined oils, effluents p0068 A75-17221 and vegetation Application of radar imagery to environmental geologic soning of Taxas p0034 A75-17222 ng of Texas mapping of lexas pound from the transport of the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0089 A75-17223 Surface compositional mapping in the Wind River Range lasin, Wyoming by multispectral techniques techniques applied p0028 A75-17224 to ERTS-1 data Unsupervised mapping of geologic features features and soils in p0034 A75-17225 Remote sensing to detect the toxic effects of metals on vegetation for mineral exploration p0034 A75-17226
Elements of the deep structure of the earth's crust on the space images of the East Caucasus p0028 A75-17228 The use of space photos for search of oil and gas fields p0034 A75-17229 The remote-sensing program of the Geological Survey Alabama p0069 A75-17230 p0069 A75-17230 in Michigan --- NASA p0069 A75-17231 of Alabama remote sensing
Applications of remote sensing by the state of p0069 A75-17232 Nebraska Application of ERTS-1 data to the protection and management of New Jersey's coastal environme p0042 A75-17233 Multidisciplinary applications of ERTS and Skylab data p0015 A75-17234 An overview of Texas activities in remote sensing p0076 A75-17235 American Society of Photogrammetry, Fall Meeting, Washington, D.C., September 10-13, 1974, Proceedings p0056 A75-18052 Remote sensing and land use inventory and mapping p0015 A75-18057
Land use classification with ERTS-1 data in the Houston ea test site p0015 A75-18059 Remote sensing for regional resource analysis p0015 A75-18061 Small scale thematic mapping - A case for radar p0028 A75-18062 imagery The application of ERTS-1 data to the Corps of Engineers Civil Works Mission --- for ecological monitoring p0015 A75-18064 Automatic mapping of strip mine operations from pacecraft data p0034 A75-18065 spacecraft data Trends in the use of remote sensing for state resources p0005 A75-18068 and environmental management p0005 A75-18068
Application of ERTS data to environmental inventory and assessment
The Skylab S-1908 Earth Terrain Camera amera p0069 A75-18071 Automatic pattern recognition and mapping utilizing ermal infrared imagery p0070 A75-18074 thermal infrared imagery Earth sciences in the age of the satellite --- remote sensing principles and techniques p0070 A75-18693 Conference on Environmental S sors and Applications, London, England, November 18, 19, 1974, Proceedings p0070 A75-19265 A radar altimeter for satellite applicati [AD-784373] AD-784373] p0056 N75-10425 A study of a dual polarization laser backscatter system for remote identification and measurement of water [NASA-CR-140665] p0017 N75-10563 The application of remote sensing techniques: Technical and methodological issues
[NASA-CR-140674] p0070 N75-10564 Remote sensing applications in marine science programs of VIMS [NASA-CR-137475] p0043 N75-10566 A survey for the use of remote sensing in the Chesapeake ay region INASA-CR-1374761 n0043 N75-10571 Application of remote sensing for prediction and detection of thermal pollution [NASA-CR-139182] DO017 N75-10572 Pattern recognition relevant to reconnaissance and remote sensing --- noting classification metho [PHL-1974-23] DOO70 p0070 N75-10573 The Coho Project: Living resources prediction feasibility study, volume 1 --- meteorological forecasting of fish concentrations [PB-234057/8] n0044 N75-10681 The Coho Project: Living resources prediction feasibility study. Volume 2: Environmental report [PB-234058/6] p0044 N75-10682 The Coho Project: Living resources prediction feasibility Volume 3: System evaluation report p0044 N75-10683 [PB-234059/4] Application of remote sensing to state and regional

An analysis of applications development systems for The remote identification of terrain features and materials remotely sensed multispectral data for the Earth at a California test site: An investigative study of Observations Division of the NASA Lyndon B. Johnson [PB-235991/7] p0031 N75-14261 Space Center [NASA-CR-140345] p0071 N75-11447 The remote identification of terrain features and materials at Pennsylvania test sites: An investigative study of Oil-on-Water sensor --- for detecting oil by emitting techniques [AD-784104] [PB-235992/5] p0031 N75-14263 p0072 N75-11454 Investigation of transient earth resources phenomena: Correlation spectroscopy techniques for the regional and global measurements of air pollution p0018 N75-11505 Continuation study [NASA-CR-139163] p0073 N75-15131 Measurements of meteorological parameters and air Usefulness of remote sensing techniques for the environmental monitoring of the Sacramento-San Josquin pollution spread over Stockholm, Sweden, by means of remote sensing and airborne instrumentation p0025 N75-15135 p0018 N75-11531 Fundamentals of the space indication of earth Proceedings of the UCLA International Conference on resources [NASA-TT-F-16108] Radiation and Remote Probing of the Atmosphere --emphasizing radiative transfer
[NASA-CR-140816] p0078 N75-11549 p0074 N75-15136 Feasibility of remote sensing for detecting thermal pollution. Part 1: Feasibility study. Part 2: Implementation Infrared remote sounding p0072 N75-11562 plan --- coastal ecology [NASA-CR-134453] Systems study of an automated fire weather data p0025 N75-15199 RESEARCH AND DEVELOPMENT [NASA-TM-X-62402] p0006 N75-12277 Research in earth resources engineering, and pollution [AR-29] utilization, physical Recordings of the spectral-radio-meter SIGNE during the mmer of 1972 p0022 N75-13369 RESEARCH MANAGEMENT [FOA-2-C-2577-E1] DO072 N75-12283 Scientific research and experimental design planning and Feasibility study ASCS remote sensing/compliance coordination for 1973 determination syste [AD-785998] p0081 N75-14259 INASA-CR-1342RRI p0007 N75-12419 measuring system in p0007 N75-12518 RESERVOIRS Automated remote Correlation of multispectral imagery with water analysis agrometeorology ERTS-B (Earth Resources Technology Satellite) - The Ross Barnett Reservoir Remote Sensing Project p0014 A75-17165 spacecraft design remote sensor description, and technology rilization New England reservoir manageme [E75-10065] [NASA-NEWS-RELEASE-74-329] p0079 N75-13370 p0050 N75-13365 Computer and photogrammetric general land use study Skylab study of water quality ---Kansas reservoirs and central north Alabama o0022 N75-13372 [NASA-TR-R-431] [F75-10075] n0051 N75-14194 An application of infrared remote sensing techniques to RESOURCES MANAGEMENT U.S. energy resources - Outlook for the future p0022 N75-13419 [AD-786028] p0075 A75-12999 The econo nic value of remote sensi Feasibility of using multiplexed SLAR imagery for water from space: An ERTS overview and the value of continuity of service. Volume 1: Summary resources management and mapping p0054 A75-17115 communities [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Design concepts for land use and natural resource inventories and information systems p0013 A75-17133 Rapid stock assessment of pilchard populations by reraft-borne remote sensors p0003 A75-17168 Volume 2: Source document [NASA-CR-141193] aircraft-borne remote sensors p0079 N75-14204 [NASA-UR-141193]
The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources: Agriculture. Part 1: Overview Resource analysis applications in Michigan --- NASA mote sensing p0069 A75-17231 remote sensing Multidisciplinary applications of ERTS and Skylab data in Ohio p0015 A75-17234 p0079 N75-14205 [NASA-CR-141265] A summary of ERTS data applications in Alaska p0077 A75-17236 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources, agriculture. Part 2: Distribution effects An intensive natural resources inventory aids land use planning and resources management in Puerto Rico Distribution effects p0005 A75-18056 p0079 N75-14206 [NASA-CR-141194] [NASA-CR-141194]
The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources, agriculture. Part 3: The integrated impact of improved (ERS) information on US agricultural commodities. Remote sensing for regional resource analysis p0015 A75-18061 Trends in the use of remote sensing for state resources of environmental management p0005 A75-18068 and environmental management Land use mapping using ERTS-1 imagery for The Four [NASA-CR-141262] p0079 N75-14207
The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity Rivers Resource Conservation and Development District in p0016 A75-18070 Thailand national programme of the Earth Resources Volume 4: Forestry, wildlife and rangeland Technology Satellite [E74-10806] [NASA-CR-141263] p0080 N75-14208 p0077 N75-10533 The economic value of remote sensing of earth resources The Thailand programme of the Earth Resources from space: An ERTS overview and the value of continuity of service. Volume 5: Inland water resources Technology Satellite [E74-10807] p0077 N75-10534 [NASA-CR-141269] p0080 N75-14209 Third Earth Resources Technology Satellite Symposium. ue of remote sensing of earth resources The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity Volume 2: Summary of results [NASA-SP-356] nO077 N75-10549 of service. Volume 6: Land use. Part 1: Introduction Utility of ERTS for monitoring the breeding habit of migratory waterfowl --- United States and Canada [NASA-CR-141266] p0080 N75-14210 p0006 N75-10557 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity Agriculture, forestry, range resources p0006 N75-10558 of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a nationwide Evaluation of Skylab imagery as an information service land cover information system for investigating land use and natural resources [NASA-CR-141267] p0080 N75-14211 [E75-10018] p0020 N75-12397 The economic value of remote sensing of earth resources Research in earth resources utilization, physical engineering, and pollution [AR-29] from space: An ERTS overview and the value of continuity of service. Volume 7: Nonreplenishable natural resources: p0022 N75-13369 Minerals, fossil fuels and geothermal energy sources
[NASA-CR-141264] p0080 N75-14212 The ERTS cost-benefit study [NASA-CR-141264] [PB-234065/1] p0079 N75-13373 The economic value of remote sensing of earth resources Evaluation of Skylab imagery as an information service from space: An ERTS overview and the value of continuity for investigating land use and natural resources --- New Volume 8: Atmosphere [NASA-CR-141191] p0080 N75-14213 [E75-10068] p0059 N75-14187 The economic value of remote sensing of earth resources Evaluation of Skylab imagery as an information service m space: An ERTS overview and the value of continuity for investigating land use and natural resources --- New of service. Volume 9: Oceans [NASA-CR-141270] p0080 N75-14214 [NASA-CR-141372] p0024 N75-15100 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity_ RHODE ISLAND Investigation of land use of northern megalopolis using ERTS-1 imagery of service. Volume 10: Industry [NASA-CR-141192] p0080 N75-14215

INASA-CR-24591

[E75-10042]

DO017 N75-10545

p0072 N75-12411

In situ spectroradiometric calibration of EREP imagery

and oceanography of Block Island Sound --- Arizona

Meteorological interpretation of high resolution images for remote sensing of earth resources p0060 N75-14229

remote-sensing programme over active Italian volcano

Contribution of space platforms to a ground and airborne

p0073 N75-14238

problems --- for Mississippi [NASA-CR-140805]

p0078 N75-11414

RHOME DELTA (ERANCE)

Optical processing of ERTS-1 (MS	
for the Arcachon basin and Rhone of and Arcachon Basin, France	p0060 N75-1422
Recent and present sedimentation Rhone delta and the French mediter	rranges coast (Coline
project) ERTS-1 imagery	p0045 N75-14221
Comparison of various numer	
classification of multispectral data. A	application to geolog
ERTS-1 imagery of Rhone Valley	(France)
-, -, -, -, -, -, -, -, -, -, -, -, -, -	p0038 N75-1424
RICE	
Thermal behaviour of some rice	fields affected by
yellows-type disease	p0004 A75-17174
Agricultural resources investigation southern France and Madagascar -	ons in northern Italy
southern France and Madagascar -	ERTS-B and seria
rice growing and forestry surveys	p0009 N75-1425
RIVER BASINS	***
Pedológical interpretation of ER Rima-Sokoto river basin (N. Nigeria)	
ROCKET SOUNDING	pood 1173-1424
Geoscientific research with rockets	e and establitae
[IAF PAPER 74-116]	p0053 A75-1374
ROCKET-BORNE PHOTOGRAPHY	
The interpretation and use of false	e-colour infra-red and
true colour photography of part of	Argentina obtained b
Skylark earth resources rockets	p0067 A75-1719
The application of pattern classif Skylark earth resources rocket image	ication techniques to
valur resources rocket image	роо56 A75-1926
A comparison of the capabilities	
Skylark earth-resources rocket for r	esources surveying it
central Argentina	p0023 N75-1425
ROCKS	
Mapping exposed silicate rock typ	es and exposed ferri
and ferrous compounds from a space [E75-10008]	e platform
Remote sensing geophysics from	p0036 N75-1239
California and Nevada minerals and	meks
[E75-10045]	p0030 N75-1241
Remote sensing geophysics from S	kylab solar heating
of ground surface and infrared	reflectance of roc
samples	
[E75-10057]	p0037 N75-1335
Remote sensing geophysics from S analysis of weathered rocks	skylab stratigraphi
[E75-10058]	p0037 N75-1335
ROTATING PLASMAS	pood / 1175-1555
Magnetopause rotational forms	p0011 A75-1122
RURAL AREAS	
Some findings on the applications	o OI EN IS BIIU SKYIBI
imagery for metropolitan land use ar	nalysis
imagery for metropolitan land use ar The usefulness of imaging passive	nalysis p0013 A75-17136 e microwave for rura
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis	nalysis p0013 A75-17139 e microwave for rure p0055 A75-17199
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis Study of recreational land and ope	nalysis p0013 A75-17139 e microwave for rure p0055 A75-17199
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis Study of recreational land and ope imagery southeast Michigan	nalysis p0013 A75-17139 e microwave for rurs p0055 A75-17199 en space using Skytal
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis Study of recreational land and op- imagery southeast Michigan [E75-10034]	nalysis p0013 A75-17139 e microwave for rure p0055 A75-17199
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis Study of recreational land and op- imagery southeast Michigan [275-10038] RURAL LAND USE	nalysis p0013 A75-1713; e microwave for rure p0055 A75-1719; en space using Skytal p0020 N75-1240;
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis Study of recreational land and op- imagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in re-	nalysis p0013 A75-1713: e microwave for rurs p0055 A75-1719: en space using Skylai p0020 N75-1240: efation to aircraft and
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis Study of recreational land and op- imagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in re- satellite studies of agricultural	nelysis p0013 A75-1713/s p0013 A75-1713/s p0055 A75-1719/s n space using Skylal p0020 N75-1240- efation to aircraft and land use and land
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis Study of recreational land and op- imagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in re-	nelysis p0013 A75-1713/s p0013 A75-1713/s p0055 A75-1719/s n space using Skylal p0020 N75-1240- efation to aircraft and land use and land
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis Study of recreational land and ope imagery southeast Michigan [275-10034] RURAL LAND USE Ground-truth measurements in re satellite studies of agricultural classification in Britain ERTS-1 and	nalysis p0013 A75-1713/s microwave for rure p0055 A75-1719/s an space using Skytal p0020 N75-1240/slation to aircraft and land use and land d serial remote sensing
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis Study of recreational land and ope imagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in restabilitie studies of agricultural classification in Britain ERTS-1 and in Great Britain	nalysis p0013 A75-1713/s microwave for rure p0055 A75-1719/s an space using Skytal p0020 N75-1240/slation to aircraft and land use and land d serial remote sensing
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis Study of recreational land and ope imagery southeast Michigan [275-10034] RURAL LAND USE Ground-truth measurements in re satellite studies of agricultural classification in Britain ERTS-1 and	nalysis p0013 A75-1713/s microwave for rure p0055 A75-1719/s an space using Skytal p0020 N75-1240/slation to aircraft and land use and land d serial remote sensing
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis. Study of recreational land and ope imagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in resatellite studies of agricultural classification in Britain ERTS-1 and in Great Britain	nalysis p0013 A75-1713/s microwave for rure p0055 A75-1719/s an space using Skytal p0020 N75-1240/slation to aircraft and land use and land d serial remote sensing
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis Study of recreational land and ope imagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in resatellite studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA)	nelysis pools A75-1713; e microways for rure p0055 A75-1719; e microways for rure p0055 A75-1719; en space using Skyla p0020 N75-1240; fation to eircraft and use and land serial remote sensin p0009 N75-1425
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis. Study of recreational land and optimagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in restabilitie studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA) Usefulness of remote sensing environmental monitoring of the Sac	nelysis p0013 A75-1713(e microwave for rure p0055 A75-1713(en space using Skyte p0020 N75-1240- piation to eircraft an land use and land d serial remote sensin- p0009 N75-1425
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis Study of recreational land and ope imagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in resatellite studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA) Usefulness of remote sensing environmental monitoring of the Sac Deta	nelysis p0013 A75-1713/ e microwave for rure p0055 A75-1719/ en space using Skyla p0020 N75-1240- efstion to eircraft and land use and land d serial remote sensin p0009 N75-1425
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis. Study of recreational land and optimagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in resatellite studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA) Usefulness of remote sensing environmental monitoring of the Sac Deta SAUNITY	nelysis p0013 A75-1713(e microwave for rure p0055 A75-1719(en space using Skyta p0020 N75-1240- po020 N75-1240- pool series and land d serial remote sensin- p0009 N75-1425 techniques for the cramento-San Josquii p0025 N75-1513;
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis and urban terrain analysis Study of recreational land and optimagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in restabilitie studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA) Usefulness of remote sensing environmental monitoring of the Sac Delta SALUNITY Irrigation scheduling, freeze war	nelysis p0013 A75-1713; e microwave for rure p0055 A75-1719; e microwave for rure p0055 A75-1719; e microwave for rure p0020 N75-1240- efation to aircraft and land use and land d serial remote sensin p0009 N75-1425 techniques for th ramento-San Josqui p0025 N75-1513; ming and soil saliniti
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis. Study of recreational land and ope imagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in resatellite studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA) Usefulness of remote sensing environmental monitoring of the Sac Delta SALINITY Irrigation scheduling, freeze war detecting Cameron County, Texas	nelysis p0013 A75-1713/ e microwave for rure p0055 A75-1713/ e microwave for rure p0055 A75-1713/ e microwave for rure p0055 A75-1713/ p0020 N75-1240- p10020 N75-1425 techniques for the ramento-San Josquii p0025 N75-1513/ ming and soil salinitis
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis. Study of recreational land and optimagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in resatellite studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA) Usefulness of remote sensing environmental monitoring of the Sac Delta SAUNITY Irrigation scheduling, freeze war detecting Cameron County, Texas [E75-10046]	nelysis p0013 A75-1713; e microwave for rure p0055 A75-1719; e microwave for rure p0055 A75-1719; e microwave for rure p0020 N75-1240; efation to eircraft and land use and land d serial remote sensin p0009 N75-1425; techniques for the rramento-San Josquii p0025 N75-1513; ming and soil selinits p0007 N75-1241;
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis and urban terrain analysis Study of recreational land and optimagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in restellite studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA) Usefulness of remote sensing environmental monitoring of the Sac Delta BAUNITY Irrigation scheduling, freeze war detecting Cameron County, Texas [E75-10048] Irrigation scheduling, freeze warr	nelysis pO013 A75-1713; e microwave for rure pO055 A75-1719; e microwave for rure pO055 A75-1719; e microwave for rure pO055 A75-1719; pO009 N75-1240- techniques and land aerial remote sensin pO009 N75-1425 techniques for th ramento-San Joaqui pO025 N75-1513; ming and soil salinit s pO007 N75-1241- ting and soil salinit godor N75-1124- ting and soil salinit
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis and urban terrain analysis Study of recreational land and optimagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in restabilitie studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA) Usefulness of remote sensing environmental monitoring of the Sac Delta SALUNITY Irrigation scheduling, freeze war detecting Cameron County, Texas [E75-10046] Irrigation scheduling, freeze war detecting Cameron County, Texas (E75-10050)	nelysis pO013 A75-1713; e microwave for rure pO055 A75-1719; e microwave for rure pO055 A75-1719; e microwave for rure pO050 N75-1240; fation to aircraft and land use and land d serial remote sensin pO009 N75-1425; techniques for th ramento-San Josqui pO025 N75-1513; ming and soil salinitis pO007 N75-1241; ining and soil salinitis pO007 N75-1241; ining and soil salinitis pO005 N75-1335;
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis and urban terrain analysis Study of recreational land and optimagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in restabilitie studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA) Usefulness of remote sensing environmental monitoring of the Sac Delta SALUNITY Irrigation scheduling, freeze war detecting Cameron County, Texas [E75-10046] Irrigation scheduling, freeze war detecting Cameron County, Texas (E75-10050)	nelysis pO013 A75-1713; e microwave for rure pO055 A75-1719; e microwave for rure pO055 A75-1719; e microwave for rure pO050 N75-1240; fation to aircraft and land use and land d serial remote sensin pO009 N75-1425; techniques for th ramento-San Josqui pO025 N75-1513; ming and soil salinitis pO007 N75-1241; ining and soil salinitis pO007 N75-1241; ining and soil salinitis pO005 N75-1335;
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis, Study of recreational land and optimagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in resatellite studies of agricultural classification in Britain ERTS-1 and in Great Britain soft agricultural leassification in Britain ERTS-1 and in Great Britain Cameron County, Texast (E75-10046) Irrigation scheduling, freeze warr detecting Cameron County, Texast (E75-10050) Irrigation scheduling, freeze war detecting Cameron County, Texast (E75-10050) Irrigation scheduling, freeze war detecting Cameron County, Texast (E75-10050)	nelysis p0013 A75-1713; e microwave for rure p0055 A75-1719; e microwave for rure p0055 A75-1719; e microwave for rure p0050 N75-1240; station to sircraft and and use and land a serial remote sensin p0009 N75-1425; techniques for the tramento-San Josquii p0025 N75-1513; ming and soil salinit p0007 N75-1241; ing and soil salinit p0050 N75-1335; ming and soil salinit p0050 N75-1335;
The usefulness of imaging passive and urban terrain analysis. Study of recreational land and operating of the studies of the studies of the studies of the studies of agricultural liclassification in Britain ERTS-1 and in Great Britain in Great Britain in Great Britain of the studies of agricultural liclassification in Britain ERTS-1 and in Great Britain in Great Britain ERTS-1 and in Great Britain in Great Britain ERTS-1 and i	techniques for the ramento-San Joegus po005 N75-1240- po015 A75-1713 e microwave for rure po055 A75-1713 e microwave for rure po055 A75-1713 po009 N75-1240- techniques and land aerial remote sensin po009 N75-1425 techniques for the ramento-San Joegui po025 N75-1513 ming and soil salinitis po007 N75-1241 po005 N75-1335- ming and soil salinitis po005 N75-1420
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis at the property of recreational land and operating the property of the prop	nelysis p0013 A75-1713; e microwave for rure p0055 A75-1719; e microwave for rure p0055 A75-1719; e microwave for rure p0055 A75-1719; p0020 N75-1240; station to eircraft and land use and land d serial remote sensin p0009 N75-1425; techniques for the tramento-San Joequi p0025 N75-1513; ming and soil salinit s p0007 N75-1241; ining and soil salinit p0005 N75-1335; minty Texas p0008 N75-1420; minty Texas p0008 N75-1420; minty Texas p0008 N75-1420; minty Texas
The usefulness of imaging passive and urban terrain analysis. Study of recreational land and operating of the studies of the studies of the studies of the studies of agricultural liclassification in Britain ERTS-1 and in Great Britain in Great Britain in Great Britain of the studies of agricultural liclassification in Britain ERTS-1 and in Great Britain in Great Britain ERTS-1 and in Great Britain in Great Britain ERTS-1 and i	nelysis pools and solid semicrowave for rure p0055 A75-1713; a microwave for rure p0055 A75-1718; a microwave for rure p0055 A75-1718; p0000 N75-1240; station to aircraft and land use and land derial remote sensing p0009 N75-1425; techniques for the ramento-San Josqui p0025 N75-1513; ming and soil salinities p0007 N75-1241; ning and soil salinities, Texas p0008 N75-1420; ning and soil salinities, Texas p0008 N75-1420; ning and soil salinities.
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis and urban terrain analysis Study of recreational land and optimagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in restellite studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA) Usefulness of remote sensing environmental monitoring of the Sac Delta SALUNITY Irrigation scheduling, freeze war detecting Cameron County, Texas [E75-10046] Irrigation scheduling, freeze war detecting Cameron and Starr Cou [E75-10084] Irrigation scheduling, freeze war detecting Cameron and Starr Cou [E75-10084] Irrigation scheduling, freeze war detecting Cameron and Starr Cou [E75-10084] Irrigation scheduling, freeze war detecting Cameron and Starr Cou [E75-10084] SALYUT SPACE STATION	nelysis p0013 A75-1713; e microwave for rure p0055 A75-1719; e microwave for rure p0055 A75-1719; e microwave for rure p0055 A75-1719; p0020 N75-1240; station to eircraft and land use and land d serial remote sensin p0009 N75-1425; techniques for the tramento-San Joequi p0025 N75-1513; ming and soil salinit s p0007 N75-1241; ining and soil salinit p0005 N75-1335; minty Texas p0008 N75-1420; minty Texas p0008 N75-1420; minty Texas p0008 N75-1420; minty Texas
The usefulness of imaging passive and urban terrain analysis. Study of recreational land and operating of recreational land and operating of recreational land and operating of recreations of the land of the land of the land of the land of	nelysis pools a75-1713/e microwave for rure p0055 A75-1713/e microwave for rure p0055 A75-1718/e microwave for rure p0055 A75-171240-p0020 N75-1240-p0020 N75-1425 techniques for the ramento-San Joaqui p0025 N75-1513/ming and soil salinitiating and soil salinitianty. Texas p0008 N75-1420 ing and soil salinitianties. Texas p0001 N75-1512 indies. Texas p0010 N75-1512
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis and urban terrain analysis Study of recreational land and opi imagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in restellite studies of agricultural classification in Britain ERTS-1 and in Great Britain soft agricultural least life in Great Britain in Great Britain ERTS-1 and in Great Britain ERTS-1 and in Great Britain soft remote sensing environmental monitoring of the Sec Delta SAUNITY Irrigation scheduling, freeze warr detecting Cameron County, Texas [E75-10048] Irrigation scheduling, freeze warr detecting Cameron County, Texas detecting Cameron and Starr Co [E75-10084] Irrigation scheduling, freeze warr detecting Cameron and Starr Co [E75-10084] Irrigation scheduling, freeze warr detecting Cameron and Starr Co [E75-10084] ALYUT SPACE STATION The earth in the lens of Salyut-3 [NASA-TT-F-18028]	nelysis pools and solid semicrowave for rure p0055 A75-1713; a microwave for rure p0055 A75-1718; a microwave for rure p0055 A75-1718; p0000 N75-1240; station to aircraft and land use and land derial remote sensing p0009 N75-1425; techniques for the ramento-San Josqui p0025 N75-1513; ming and soil salinities p0007 N75-1241; ning and soil salinities, Texas p0008 N75-1420; ning and soil salinities, Texas p0008 N75-1420; ning and soil salinities.
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis and urban terrain analysis Study of recreational land and optimagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in restellite studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA) Usefulness of remote sensing environmental monitoring of the Sac Delta SALUNITY Irrigation scheduling, freeze war detecting Cameron County, Texas [E75-10048] Irrigation scheduling, freeze war detecting Cameron and Starr Co. [E75-10050] Irrigation scheduling, freeze war detecting Cameron and Starr Co. [E75-10109] SALYUT SACE STATION The earth in the lens of Salyut-3 [NASA-TT-F-18028] SAN FRANCISCO (CA)	nelysis pOO13 A75-1713; e microwave for rure pOO55 A75-1713; e microwave for rure pOO55 A75-1718; en space using Skyla pOO20 N75-1240- sfation to aircraft and and use and land a erist remote sensin pOO09 N75-1425 techniques for the strain pool of the strain p
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis Study of recreational land and optimagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in restellite studies of agricultural classification in Britain ERTS-1 and in Great Britain and in Great Britain and in Great Britain are in Great Britain and Information and Sacramental Monitoring of the Secondary Infragation scheduling, freeze ware detecting Cameron County, Texat [E75-10056] Irrigation scheduling, freeze ware detecting Cameron and Starr Cot [E75-10084] Irrigation scheduling, freeze ware detecting Cameron and Starr Cot [E75-10109] SALYUT SPACE STATION The sarth in the lens of Salyut-3 [RNSA-TT-F-18028] SAN FRANCISCO (CA) Carrigraphic evaluation of Sky	nelysis pO013 A75-1713; e microwave for rure pO055 A75-1719; e microwave for rure pO055 A75-1719; e microwave for rure pO055 A75-1719; e microwave for rure pO020 N75-1240; station to eircraft and land use and land d serial remote sensin pO009 N75-1425; techniques for th tramento-San Joequi pO025 N75-1513; ming and soil salinit pO007 N75-1241; ming and soil salinit pO008 N75-1335 ming and soil salinit pO008 N75-1335 ming and soil salinit ping and soil salinit ping and soil salinit mittes, Texas pO010 N75-1512 p0070 N75-1512
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis and urban terrain analysis Study of recreational land and optimagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in restellite studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA) Usefulness of remote sensing environmental monitoring of the Sac Delta SALUNITY Irrigation scheduling, freeze war detecting Cameron County, Texas [E75-10048] Irrigation scheduling, freeze war detecting Cameron and Starr Co. [E75-10050] Irrigation scheduling, freeze war detecting Cameron and Starr Co. [E75-10109] SALYUT SACE STATION The earth in the lens of Salyut-3 [NASA-TT-F-18028] SAN FRANCISCO (CA)	nelysis pO013 A75-1713; e microwave for rure pO055 A75-1719; e microwave for rure pO055 A75-1719; e microwave for rure pO055 A75-1719; e microwave for rure pO020 N75-1240; station to eircraft and land use and land d serial remote sensin pO009 N75-1425; techniques for th tramento-San Joequi pO025 N75-1513; ming and soil salinit pO007 N75-1241; ming and soil salinit pO008 N75-1335 ming and soil salinit pO008 N75-1335 ming and soil salinit ping and soil salinit ping and soil salinit mittes, Texas pO010 N75-1512 p0070 N75-1512
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis and urban terrain analysis Study of recreational land and opi imagery	nelysis pO013 A75-1713; e microwave for rure pO055 A75-1719; e microwave for rure pO055 A75-1719; e microwave for rure pO055 A75-1719; e microwave for rure pO020 N75-1240; station to eircraft and land use and land d serial remote sensin pO009 N75-1425; techniques for th tramento-San Joequi pO025 N75-1513; ming and soil salinit pO007 N75-1241; ming and soil salinit pO008 N75-1335 ming and soil salinit pO008 N75-1335 ming and soil salinit ping and soil salinit ping and soil salinit mittes, Texas pO010 N75-1512 p0070 N75-1512
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis Study of recreational land and opimagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in restellite studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA) Usefulness of remote sensing environmental monitoring of the Sac Delta BALINITY Irrigation scheduling, freeze war detecting Cameron County, Texas [E75-10048] Irrigation scheduling, freeze war detecting Cameron and Starr Co. [E75-10050] Irrigation scheduling, freeze war detecting Cameron and Starr Co. [E75-10109] SALYUT SPACE STATION The earth in the lens of Salyut-3 [NASA-TT-F-18028] SAN FRANCISCO (CA) Cartographic evaluation of Sky images Chesapeaks Bay, Phoen test sites [E75-10078] SAN FRANCISCO BAY (CA)	nelysis pOO13 A75-1713; e microwave for rure pOO55 A75-1713; e microwave for rure pOO55 A75-1713; e microwave for rure pOO55 A75-1712, pOO20 N75-1240; fation to aircraft and use and land serial remote sensin pOO09 N75-1425; techniques for the ramento-San Joepon pOO25 N75-1513; ming and soil salinities, poo15 N75-1420; poo15 N75-1096; poo15 N75-1096
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis at the property of recreational land and operating the property of the prop	nelysis pO013 A75-1713; e microwave for rure pO055 A75-1719; e microwave for rure pO055 A75-1719; e microwave for rure pO055 A75-1719; e microwave for rure pO020 N75-1240; station to eircraft and land use and land d serial remote sensini pO009 N75-1425; techniques for th tramento-San Joequi pO025 N75-1513; ming and soil salinit pO007 N75-1241; ming and soil salinit pO008 N75-1335 ming and soil salinit pO008 N75-1335 ming and soil salinit ping and soil salinit mittes, Texas pO010 N75-1512 p0070 N75-1098 stat-A S-192 scanne ix, and San Francisc pO031 N75-1419
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis and urban terrain analysis Study of recreational land and optimagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in restelline studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA) Usefulness of remote sensing environmental monitoring of the Sac Delta SALUNITY Irrigation scheduling, freeze war detecting Cameron County, Texas [E75-10048] Irrigation scheduling, freeze war detecting Cameron and Starr Co. [E75-10050] Irrigation scheduling, freeze war detecting Cameron and Starr Co. [E75-10084] Irrigation scheduling, freeze war detecting Cameron and Starr Co. [E75-10109] SALYUT SPACE STATION The earth in the lens of Salyut-3 [NASA-TT-F-18028] SAN FRANCISCO (CA) Cartographic evaluation of Sky images Chesapeake Bay, Phoen test sites [E75-10078] SAN FRANCISCO BAY (CA) Surface currents along the Califoners.	nelysis pOO13 A75-1713; e microwave for rure pOO55 A75-1713; e microwave for rure pOO55 A75-1713; e microwave for rure pOO55 A75-1712, pOO20 N75-1240; fation to aircraft and use and land serial remote sensin pOO09 N75-1425; techniques for the ramento-San Joepon pOO25 N75-1513; ming and soil salinities, poo15 N75-1420; poo15 N75-1096; poo15 N75-1096
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis at the study of recreational land and operations of the study of recreational land and operation of the study of recreational land and operation of the study of recreational land and operation of the study	nelysis pO013 A75-1713 e microwave for rure pO055 A75-1719 e microwave for rure pO055 A75-1718 e microwave for rure pO055 A75-1718 e microwave for rure pO055 A75-1718 pO020 N75-1240 techniques for the ramento-San Joequi pO025 N75-1513 ming and soil salinit pO007 N75-1241- ming and soil salinit pO008 N75-1335 ming and soil salinit pO008 N75-1335 ming and soil salinit pO008 N75-1335 ming and soil salinit pO008 N75-1419 po070 N75-1512 pO070 N75-1512 po070 N75-1512 po071 N75-1513
imagery for metropolitan land use ar The usefulness of imaging passive and urban terrain analysis and urban terrain analysis Study of recreational land and optimagery southeast Michigan [E75-10034] RURAL LAND USE Ground-truth measurements in restelline studies of agricultural classification in Britain ERTS-1 and in Great Britain S SACRAMENTO VALLEY (CA) Usefulness of remote sensing environmental monitoring of the Sac Delta SALUNITY Irrigation scheduling, freeze war detecting Cameron County, Texas [E75-10048] Irrigation scheduling, freeze war detecting Cameron and Starr Co. [E75-10050] Irrigation scheduling, freeze war detecting Cameron and Starr Co. [E75-10084] Irrigation scheduling, freeze war detecting Cameron and Starr Co. [E75-10109] SALYUT SPACE STATION The earth in the lens of Salyut-3 [NASA-TT-F-18028] SAN FRANCISCO (CA) Cartographic evaluation of Sky images Chesapeake Bay, Phoen test sites [E75-10078] SAN FRANCISCO BAY (CA) Surface currents along the Califoners.	nelysis pO013 A75-1713 e microwave for rure pO055 A75-1719 e microwave for rure pO055 A75-1718 e microwave for rure pO055 A75-1718 e microwave for rure pO055 A75-1718 pO020 N75-1240 techniques for the ramento-San Joequi pO025 N75-1513 ming and soil salinit pO007 N75-1241- ming and soil salinit pO008 N75-1335 ming and soil salinit pO008 N75-1335 ming and soil salinit pO008 N75-1335 ming and soil salinit pO008 N75-1419 po070 N75-1512 pO070 N75-1512 po070 N75-1512 po071 N75-1513

SAND HILLS REGION (NE)

p0021 N75-13339

IUCID-18585

[E75.10044] radiation imagen achievements [IAF PAPER 74-116] (IAF PAPER 74-165) County, South Dakota NOAA-2 satellite using ERTS-1 data currents and sediments [AD-781138/3] rone data [UCRL-75556] Status and p [JPRS-63231] environmental monitoring of the Sacramento-San Jos p0025 N75-15135 Satellite pictures can help guide Sandhills irrigation center-pivot irrigation in Sand Hills Region, Nebraska [E75-10023] P0021 N75-133:

Use of ERTS-1 imagery to interpret wind-erosion hazard in the Sandhills of Nebraska Fundamentals of the space indication of earth p0059 N75-13347 [NASA-TT.F. 16108] n0074 N75-15136 SATELLITE ATTITUDE CONTROL SATELLITE ORBITS Skylab. II - The three month vigil SATELLITE OBSERVATION p0065 A75-13963 ERTS 1 flight evaluation report, January 23, 1973 to April 23, 1973 --- analysis of ERTS-1 systems during orbits The detection by ERTS-1 of wind-induced ocean surface 2600-3810 features in the lee of the Antilles Islands [NASA-CR-140809] p0077 N75-10548 p0041 A75-10844 SATELLITE TRACKING A satellite view of weather systems over the North merican Arctic p0063 A75-10790 Analytical estimates of precision in determining parameters of motion of artificial earth satellite when merican Arctic
A simulation of the San Andreas fault experiment p0033 A75-11068 analyzing imprecise measurements w p0053 A75-11419 The earth as a radio source - Terrestrial kilometric diation p0011 A75-11211 Satellite techniques for the study of secular variations position p0064 A75-12980 in position Magnetopause rotational forms p0011 A75-11221 Earth parameters from global satellite triangulation and Solar wind and geomagnetic activity trilateration p0064 A75-12985 DO027 A75-11975 Rainfall estimations from geosynchronous satellite Transformation of terrestrial su vey data to Doppler pO054 A75-16569 satellite datum Skylab contributions to the future --- summary of chievements p0075 A75-13618 Applications of satellite geodesy the internal structure of the earth the exploration of p0029 A75-18123 Geoscientific research with rockets and satellites Possibilities of calculating the earth's local gravitational dd from satellite observations p0029 A75-18524 p0053 A75-13741 field from satellite observations Economic effectiveness of using the space systems of ATELLITE-BORNE INSTRUMENTS distant indication of natural resou The community satellite. III p0053 A75-11822 p0075 A75-13770 Skylab altimeter applications and scientific results Estimation of sea surface temperature from remote p0063 A75-12035 [AIAA PAPER 74-1221] sensing in the 11to 13-micron window region p0065 A75-14803 Very high resolution radiometer DO064 A75-12109 Orientation of methods of information processing for the The radio emission of the earth as a a planet --- Russian p0076 A75-15319 control of terrestrial resources [IAF PAPER 74-140] p0053 A75-13758 A radio picture of the earth --- microwave remote sensing A ratio picture of the earth --- microwave failure sensing poofs A75-16191 Coastal and estuarine studies with ERTS-1 and Skylab p0047 A75-16240 space techniques p0065 A75-13868 Investigations of the ionosphere by Skylab radar altimeter - Short-wa length perturbations atmospheric radiance for p0054 A75-18478 detected in ocean surface profiles p0066 A75-16096 correction of ERTS-1 data Investigation of microwave hologram techniques for International Symposium on Remote Sensing of Environment, 9th, University of Michigan, Ann Arbor, Mich., application to earth resources p0067 A75-17195 ave radiometers --- for Electrically scanning microw April 15-19, 1974, Proceedings. Volu mes 1, 2 & 3 satellite-borne remote sensing p0058 A75-17202
An orbiting visible/infrared spectrometer for terrestrial, p0066 A75-17101 A remote sensing study of Pacific hurricane Ava atmospheric and oceanographic applications --- on Skylab p0012 A75-17112 p0068 A75-17212 Determination of the aerosol content in the atmosphere Investigations of diffuse cosmic gamma radiation in the from ERTS-1 data
Use of ERTS-1 imagery for land p0013 A75-17130 range 28 keV-4.1 MeV p0015 A75-17667
The potentialities for determining content of minor gas constituents in the atmosphere by satellites uation in Pennington o0055 A75-17135 Flood inundation in the southeastern United States for p0015 A75-17932 p0047 A75-17138 aircraft and satellite imagery rcraft and satellite imagery
Mapping of the 1973 Mississippi River floods by the p0028 A75-17139 SATELLITE-BORNE PHOTOGRAPHY A satellite-aircraft thermal study of the upwelled waters p0041 A75-10645 off Spanish Sahara A satellite view of weather systems over the North Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models p0067 A75-17148 merican Arctic
Satellite photography of eddies in the gulf loop current p0041 A75-11255 Interactive machine asset of critical land resources p0013 A75-17149 Agricultural inventory techniques s with orbital and p0001 A75-12872 Agricultural integral power and integral from the first fine and fine first fi Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 sate p0048 A75-17160 Thermal emission of applomerations as recorded on Remote sensing of western Lake Supe environmental satellite imagery --- of European urban p0048 A75-17161 Determining range condition and forage production potential in California from ERTS-1 imagery p0003 A75-17169 [IAF PAPER 74-139] n0011 A75-13757 Images from the NOAA-3 very high resolution radiometer over the North Sea and adjoining countries ERTS 1 --- coastal California nearshore processes p0066 A75-16188 The detection of dust storms over land and water with in modelling of the p0014 A75-17204 The calculation of cloud shadows satellite visible and infrared measure earth's surface survey from space Surface compositional mapping in the Wind River Range Environmental studies of Iceland with ERTS-1 imagery and Basin, Wyoming by multispectral techniques applied to ERTS-1 data p0028 A75-17224 ERTS applications in Thailand - A progress report p0076 A75-17123 Some findings on the applications of ERTS and Skylab The application of ERTS-1 data to the Corps of Engineers Civil Works Mission --- for ecological monitoring p0015 A75-18064 imagery for metropolitan land use analysis mental inventory and p0016 A75-18069 Application of ERTS data to environ Moisture detection from Skylab -and Kansas sites
Automatic classification of eutrop ination with satellite Use of terrestrial gravity data in com results for determining goold heights in Central Europe from spacecraft data p0029 A75-18128 Preliminary results of fisheries in Observing and forecasting local effects from satellite data with Skylab-3 mesoscale cloud patterns p0016 A75-18153 atellite --- remote sensing Earth sciences in the age of the p0070 A75-18893 First results from the crop identification technology principles and techniques Soviet satellite geodesy: Basis and status of Soviet laser assessment for remote sensing / CITARS/ odesy and long-base line interferometry p0029 N75-10524 Computer derived coastal water classifications via Comparison of the Southern and Northern Hemisphere spectral signatures general circulation characteristics as determined by satellite Lake Ontario water mass delineation from ERTS-1 Evaluation of ERTS data utilization in developing puntries p0055 A75-17193
The use of space photos for search of oil and gas fields p0034 A75-17229 p0017 N75-10838 rospects of development of space geode p0029 N75-10869 Global measurements of air pollution from satellites --ruploying radiometer techniques p0018 N75-11521 The application of ERTS-derived information to the gional land use planning process p0015 A75-18058 emploving radiometer techniques regional land use planning process
The Skylab S-1908 Earth Terrain Ca Environmental satellite imagery, September 1974 p0058 N75-11571 The role of ERTS in the establishment and of a nationwide land cover information system [NASA-CR-141045] [NASA-TT-F-798] DO020 N75-12425 Geometric distortions of opticommercial panoramic Environmental definition program satellite monitoring sion systems Geographic correlation of televis of the stratosphere and troposphere

a0022 N75-13458

Geographical correlation of earth cloud cover pictures obtained from unoriented satellites p0071 N75-11435 Geological and geophysical studies from space and ospects for their development p0035 N75-11436
Experience in integrated geological and geographical interpretation of space photographs obtained in the USSR An essessment of the potential contributions to earnography from Skylab visual observations in the depletors of the potential contributions to earnography from Skylab visual observations. oceanography from Skylab visual observations and hand-held photography --- Pacific Ocean, Atlantic Ocean nd Gulf of Mavico Application of Skylab EREP photographs to study of the oldern episode of accelerated erosion in southern nodern episode of accelerated [E75-10054] p0036 N75-13354

Meteorological data. Volume 145 no. 1: The European climatology 1974 meteorological satellite data from the Berlin zone of reception, part 1, 1st quarter p0023 N75-13479 Meteorological data. Volume 139, no. 4: The European climatology, 1973. Meteorological satellite-data from the Berlin zone of reception, part 4, fourth quarter

p0023 N75-13483 An evaluation of stereoscopic viewing of ERTS and Skylab p0060 N75-14220 nages p0060 N/5-14220
The potential of small-scale mapping from ERTS-1
nagery p0060 N75-14221

SATELLITE-BORNE RADAR

A preliminary feasibility study on a synthetic aperture radar /SARSAT/ for earth resources surveys p0067 A75-17194

SCATTERING

Scattering and absorption from poly-dispe p0019 N75-11553 Multiple scattering in cloud layers; some results --- emphasizing aerosol parameters on global basis

p0019 N75-11554

SCATTEROMETERS

Simultaneous active and passive microwave response of the earth - The Skylab radscat experiment p0012 A75-17114

Radar cross sections of vegetation canopies determined by monostatic and bistatic scatterometry

p0005 A75-17190 Detection of moisture and moisture related phenomena from Skylab --- radiometer and scatterometer analysis of Texas site

10007 75-10007]
Design data collection with Skylab/EREP microwave p0007 N75-13333 instrument S-193 --- Utah and measurements

E75-10070] p0059 N75-14189
Design data collection with Skylab/EREP microwave nstrument S-193 --- Texas, Utah, Illinois, and Minnesota [E75-10070] p0072 N75-14199 [F75-10080]

Irrigation scheduling, freeze warning and soil salinity detecting --- Cameron and Starr Counties, Texas [E75-10109] p0010 N75-15125 SCIENTIFIC SATELLITES

p0053 A75-11822 The community satellite. III

SEA ICE A study of microwave emission proper perties of sea ice -p0041 A75-17120

AIDJEX 1972 Investigation of radar discrimination of sea ice p0042 A75-17121

Use of ERTS-1 in sea-ice studies p0045 N75-14234 AIDJEX bulletin number 25 --- study of pack ice, sea and Beaufort Sea ice [PB-235298/7] p0045 N75-14262

SEA ROUGHNESS Skylab radar altimeter - Short-wavelength perturbations detected in ocean surface profiles p0066 A75-16096

SEA STATES

Bistatic radar sea state monitoring [NASA-CR-137469] p0043 N75-10673

BEA WATER

Estimation of sea surface temperature from remote sensing in the 11to 13-micron window

p0065 A75-14803 urements of the dielectric properties of sea water at 1.43 GHz

p0045 N75-13496 [NASA-CR-2458]

SFAS

Analysis of methods for the geodetic support of complex sea geophysical surveys [JPRS-63621] p0031 N75-13498

SECULAR VARIATIONS

ite techniques for the study of s of secular variations p0064 A75-12980 in position The effect of the secular change of the terrestrial gravity p0028 A75-18122 field on elevation level differences On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125

SEDIMENT TRANSPORT

Dynamics of suspended sediment plumes in Lake p0049 N75-10541

SEDIMENTS

Recent and present sedimentation complexity of the Rhone delta and the French mediterranean coast (Goliproject) --- ERTS-1 imagery p0045 N75-1422 p0045 N75-14228 SEISMOLOGY

Catalog of earthquake photographs --isoseismal maps p0072 N75-13455

Cruise data report. CHAIN 115 leg 2, Southlant expedition --- to survey geophysical parameters p0029 N75-10641 [AD-784421] SHOALS

Rapid stock assessment of pilchard populations by reraft-borne remote sensors p0003 A75-17168 aircraft-borne remote sensors SIDE-LOOKING RADAR

Feasibility of using multiplexed SLAR imagery for water resources management and communities mapping vegetation p0054 A75-17115

Four channel simultaneous X-L band imaging SAR radar n0054 A75-17116 Extraction of urban land cover data from multiplexed

synthetic aperture radar imagery n0054 A75-17117 Development of research in the USSR on the use of

radar images for geological purposes p0033 A75-17126 A preliminary feasibility study on a synthetic aperture radar /SARSAT/ for earth resources su p0067 A75-17194

Application of radar imagery to environmental geologic appling of Texas p0034 A75-17222 mapping of Texas Small scale thematic mapping -A case for radar

p0028 A75-18062 SIERRA NEVADA MOUNTAINS (CA)

Study to develop improved spacecraft snow survey methods using Skylab/EREP data --- Sierra Nevada, Cascades, and Great Plains [E75-10081] p0073 N75-14200

SIGNAL PROCESSING

Developing processing techniques for Skylab data ---Michigan test site [E75-10040] p0058 N75-12409

SIGNATURE ANALYSIS

Crude and refined petroleum oil structured luminescence signatures induced by UV laser or lamp and their remote sensing applications p0033 A75-17207 SILICATES

Mapping exposed silicate rock types and exposed ferric nd ferrous compounds from a space platfo p0036 N75-12392

SKYLAB PROGRAM

Automatic land use classification using Skylab S-192 itispectral data [AIAA PAPER 74-1224] Technology today for tomorrow; Proceedings of the Eleventh Space Congress, Cocoa Beach, Fla., April 17-19, 1974. Volumes 1 & 2 p0011 A75-13576

Summary of flight performance of the Skylab Earth Resources Experiment Package / EREP/

p0078 A75-17211 The Skylab S-1908 Earth Terrain Camera

n0069 A75-18071 Geologic evaluation of central Italy from ERTS-1 and Skylab data n0038 N75-14238

SKYLARK ROCKET VEHICLE

A comparison of the capabilities of ERTS-1 and the Skylark earth-resources rocket for resources surveying in central Argentina p0023 N75-14250

SNOW

Areal extent of snow in forested regions - A practical estimation technique using ERTS-1 data

n0002 A75-17122 SNOW COVER

Microwave radiometric characteristics of snow-covered p0049 A75-17200 Microwave profiling of snowpack free-water content p0049 A75-17950

Evaluate ERTS imagery for mapping and detection of changes of snowcover on land and on glaciers --- Alaska and Washington

p0049 N75-10539 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques --- geology. hydrology, vegetation, and snow cover

Study to develop improved spacecraft snow survey methods using Skylab/EREP data ··· Sierra Nevada. Cascades, and Great Plains [E75-10081]

Methodological aspects and regional examples of snow-cover mapping from ERTS-1 and EREP imager of

the Swiss Alps p0051 N75-14232
Study of recreational land and open space using Skylab imagery ---Michigan [E75-10092] --- snow and ice hydrology of southeast

p0024 N75-15110

SOIL EROSION

Application of Skylab EREP photographs to study of the modern episode of accelerated erosion in southern p0036 N75-13354 [E75-10054]

SOIL MAPPING Mapping soil features from multispectral scanner

p0027 A75-15435 sensed units for soil p0002 A75-17106 A method of specifying remotely s sample points Use of ERTS-1 imagery for land evaluation in Pennin p0055 A75-17135 County, South Dakota

The economic impact of remote sensing data as the source of nonpoint pollution monitoring and control D0076 A75-17166

Unsupervised mapping of geologic features and soils in p0034 A75-17225 California Optical film density values from color IR photography in wetland soils mapping p0005 A75-18073

for wetland soils mapping Radar response to vegetation --- soil il moisture mapping p0005 A75-18556 microwave backscattering ERTS-1 imagery interpretation techniques in the Tennessee Valley --- land use and soil mapping [E74-10811] p0016 N75-10538

Satellite pictures can help guide Sandhills irrigation ---enter-pivot irrigation in Sand Hills Region, Nebraska [F75-10023] p0021 N75-13339

Preliminary results of the interpretation of ERTS-1 imagery for soil surveys in the Merida region, Spain p0009 N75-14247

ne remote identification of terrain features and materials at Pennsylvania test sites: An investigative study of

p0031 N75-14263 Use of forest vegetation as an indicator in the interpretation and large scale mapping of soils in a forest

[AD-785967] p0010 N75-15146

SOIL MOISTURE

Active microwave measurement of soil water content IASA-CR-141685] p0001 A75-16242 [NASA-CR-141685] Passive microwave sensing of moist soils

p0066 A75-17124 Application of remote sensing to the location of hydrologically active /source/ areas p0066 A75-17141
Use of visible, near-infrared, and thermal infrared remote sensing to study soil moisture p0002 A75-17144 Moisture detection from Skylab ---

and Kansas sites p0002 A75-17145 Remote detection of soil surface m p0003 A75-17146

On the feasibility of remote monitoring of soil moisture ith microwave sensors p0067 A75-17147 with microwave senso Radar response to vegetation soil moisture mapping

via microwave backscattering p0005 A75-18556
Detection of moisture and moisture related phenomena
from Skylab --- Kansas, Oklahoma, and Arkansas p0007 N75-12405 [E75-10035]

Detection of moisture and moisture related phenomena from Skylab --- radiometer and scatterometer analysis of

[E75-10007] D0007 N75-13333 Detection of moisture and moisture from Skylab --- Texas and Kansas [E75-10073] DOOOR N75-14192

vertess, conducted in the USSR, in 1974
NASA-TT-F-16151 [NASA-TT-F-16151]

SOIL SCIENCE Microwave emission characteristics of moist soils

p0001 A75-15180 Microbial abundance and thermoluminescence antarctic dry valley soils p0005 A75-17973
Developing processing techniques for Skylab data ---

egetation and soil identification p0057 N75-10527 [E74-10799]

Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil --- Southern Texas

[E75-10012] p0006 N75-12394 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [275-10033] p0007 N75-12403

Irrigation scheduling, freeze warning and soil salinity detecting --- Cameron County, Texas [E75-10046] p0007 N75-12414 p0007 N75-12414

trrigation scheduling, freeze warning and soil satinity detecting --- Cameron County, Texas [E75-10050] DO050 N75-13350

Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical nd thermal characteristics of plants and soil p0008 N75-14196

Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 SOILS

The PUCE programme for terrain evaluation for engineering purposes. classification 2: Procedures for terrain p0029 N75-10625 [PB-233847/3]

Irrigation scheduling, freeze warning and soil satinity detecting --- ([E75-10084] -- Cameron and Starr County, Texas |4] p0008 N75-14202

The remote identification of terrain features and materials at a California test site: An investigative study of

p0031 N75-14261

trrigation scheduling, freeze warning and soil satinity detecting --- Cameron and Starr Counties, Texas [E75-10109] p0010 N75-15125

SOLAR HEATING

Remote sensing geophysics from Skylab --- solar heating of ground surface and infrared reflectance of rock [E75-10057] p0037 N75-13357

[IAF PAPER 74-156]

SPACE EXPLORATION

SPACE COMMUNICATION

More information and less data from space --- data

duction in remote sensing systems p0063 A75-11730

The progery of Skylab - The pioneering phase of our expansion into space [AIAA PAPER 74-1216] p0063 A75-12031

p0065 A75-13765

STRATOSPHERE SOLAR INSTRUMENTS Technology today for tomorrow; Proceedings of the Computer derived coastal water classifications Eleventh Space Congress, Cocoa Beach, Fla., April 17-19, 1974, Volumes 1 & 2 p0011 A75-13576 NASA-KSC's earth resources benefits from space Apollo Telescope Mount experiments technology p0064 A75-13589 p0042 A75-17177 spectral signatures Investigation into the spectral signature of agricultural tops during their state of growth p0004 A75-17189 SOLAR OBSERVATORIES crops during their state of growth p0075 A75-13592 coloration Skylab contributions to the future -The potentialities for determining content of minor gas SPACE MANUFACTURING achievements p0075 A75-13618 The final Skylab mission: Man at home and at work in constituents in the atmosphere by satellites SOLAR POSITION p0015 A75-17932 Effect of atmospheric haze and sun angle on automatic classification of ERTS-1 data p0014 A75-17157 Automated selection and location of terrain features on [NASA-TM-X-70394] p0078 N75-12019 the basis of spectral characteristics SPACE MISSIONS SOLAR RADIATION [AD-785698] p0030 N75-13375 Investigations of the ionosphere by sp Measurements of the extraterrestrial solar radiant flux Recognition and interpretation of spectral signatures of operation from aircraft and satellite imagery in Western Queensland, Australia p0008 N75-14241 p0065 A75-13868 from 2981 to 4000 A and its transmission through the earth's atmosphere as it is affected by dust and ozone The final Skylab mission: Man at home and at work in Queensland, Australia p0070 A75-19547 [NASA-TM-X-70394] SPECTROMETERS p0078 N75-12019 Ground-based measurements of solar radiation by SPACE SHUTTLES A microwave correlation spectros Technology today for tomorrow: Proceedings of the Eleventh Space Congress, Cocoa Beach, Fla., April 17-19, 1974, Volumes 1 & 2 p0011 A75-13576 geophysical monitoring for climatic change (USA) --- using pyroheliometers, pyranometers, and radiometers monitoring p0063 A75-11732 Investigation into the spectral signification of the spectral significance of the second section of the second section of the second section of the section of the second section of the s ature of agricultural n0018 N75-11527 SOLAR WIND A Shuttle/SpaceLab RF environment survey facility The remote Raman spectrometer is a viable instrument p0016 A75-19457 Solar wind and geomagnetic activity for remote sensing of the environment DO027 A75-11975 SPACEBORNE PHOTOGRAPHY p0068 A75-17214 Skylab, II - The three month vigil o0065 A75-13963 SOMALIA Feasibility study of a Gunn-diode microwave cavity Plutonic structure of the earth's crust on space images p0027 A75-17125 Mapping of the major structures of the African rift system as a formaldehyde monitor - Ethiopia, Somalia, and Yemen [UCID-16488] p0071 N75-11311 Correlation of coastal water turbidity and current p0030 N75-13340 Correlation of couster water considered circulation with ERTS-1 and Skylab imagery p0048 A75-17181 Microwave rotational spectroscopy: A physical technique SOUNDING ROCKETS for specific pollutant monitoring p0018 N75-11529 Remote sounding of stratospheric temperatures using high resolution radiance measurements from the IRIS-D Life cycle cost study of Army Spectrometric Oil Analysis Summary of flight performance of the Skylab Earth Program (ASOAP) [AD-786501] Resources Experiment Package / EREP/ [NASA-CR-141130] p0022 N75-13452 p0073 N75-14870 p0076 A75-17211 Elements of the deep structure of the earth's crust on SOUTH AFRICA SPECTROPHOTOMETRY Mineral resources, geological structures, and landfo observations of earth from the space images of the East Caucasus Spectrophotometric p0035 N75-10561 SULVEYS p0028 A75-17228 Sovuz-13 SOUTH DAKOTA [IAF PAPER 74-156] The earth in the lens of Selyut-3 p0065 A75-13765 Use of ERTS-1 imagery for land evaluation in Pennington [NASA-TT-F-16028] p0070 N75-10964 Investigation of the utility of kinetically controlled and County, South Dakota p0055 A75-17135 Dust storms according to data of space research --catalytic reactions to perform environmental determinations Experiment to evaluate feasibility of utilizing Skylab-EREP spaceborne photography of arid zones [NASA-TT-F-16021] by parallel spectrophotometric analysis [TID-26594] remote sensing data of tectonic analysis of the Bighorn p0021 N75-12463 p0021 N75-13063 Mountains region, Wyoming-Montana --- Black Hills, South Dakota/Wyoming Computer and photogrammetric general land use study SPECTRORADIOMETERS of central north Alabama In situ spectroradiometric calibration of EREP imagery [E75-10014] n0030 N75-12395 p0022 N75-13372 and oceanography of Block Island Sound --- Arizona Inventory of forest and rangeland resources, including SPACECRAFT COMPONENTS geology [E75-10042] ERTS 1 flight evaluation report, 23 October 1973 - 23 January 1974 --- orbits 6371 to 7651 forest stress --- Colorado, Georgia, and Black Hills, South nO072 N75-12411 Spectroradiometric and multispectral scanner over Lake [NASA-CR-140712] p0077 N75-10546 [E75-10032] DO007 N75-12402 Monroe ERTS 1 flight evaluation report, 23 January - 23 April 1974 --- orbits 7652 to 8907 Applications of Skylab EREP photographs to mapping E75-10066] DO072 N75-13366 of landforms and environmental geology in the Great Plains and Midwest --- Illinois, Iowa, Kansas, Missouri, Nebraska, SPECTROSCOPIC ANALYSIS [NASA-CR-140810] p0077 N75-10547 ERTS 1 flight evaluation report, January 23, 1973 to Laser Raman spectroscopy of solutes dissolved in water from a remote platform and South Dakota p0037 N75-13355 [E75-10055] April 23, 1973 --- analysis of ERTS-1 systems during orbits [PB-233432/4] n0056 N75-10202 600-3810 SOUTHEAST ASIA Correlation spectroscopy techniques for th global measurements of air pollution p0018 N75-11505 [NASA-CR-140809] p0077 N75-10548 Comparison of the Southern and Northern Hemisphere ERTS 1 flight evaluation report, 23 April 1974-23 July 1974 --- orbits 8908 to 10182 Investigation of the utility of kinetically controlled and catalytic reactions to perform environmental determinations general circulation characteristics as determined by satellite [NASA-CR-140711] [UCRL-75556] p0017 N75-10638 p0077 N75-10568 by parallel spectrophotometric analysis [TID-26594] SPACECRAFT CONFIGURATIONS p0021 N75-13063 SOUTHERN CALIFORNIA A preliminary feasibility study on a synthetic aperture radar /SARSAT/ for earth resources surveys Oil slick detection by X-band synthetic aperture radar SPECTRUM ANALYSIS Developing processing techniques for Skylab data --egetation and soil identification p0068 A75-17209 p0067 A75-17194 Unsupervised mapping of geologic features and soils SPACECRAFT ENVIRONMENTS p0034 A75-17225 p0057 N75-10527 California [E74-10799] Skylab induced environment --- spacecraft-originated Lineaments in basement terrane of the Peninsular Ranges, ST LOUIS-KANSAS CITY CORRIDOR (MO)
Mapping of the 1973 Mississippi River floods by the contaminant effects Southern California [AIAA PAPER 74-1225] n0011 A75-12037 [E75-10003] o0035 N75-11407 NOAA-2 satellite p0028 A75-17139 SPACECREWS alt tectonics and earthquake hazards in the Peninsular Skylab environmental and crew microbiology studies On evaluating compliance with air pollution levels 'not ges. Southern California p0024 N75-14366 [E75-10005] 75-10005] p0035 N75-11408 Remote sensing geophysics from Skylab --- Southern to be exceeded more than once a yea SPAIN p0011 A75-12803 Preliminary results of the interpretation of ERTS-1 imagery for soil surveys in the Merida region, Spain p0009 N75-14247 California and Nevada minerals and rocks STATISTICAL ANALYSIS p0030 N75-12413 [E75-10045] On evaluating compliance with air pollution levels 'not Fault tectonics and earthquake hazards in the Peninsular to be exceeded more than once a year Southern California SPANISH SAHARA nO011 A75-12803 A satellite-aircraft thermal study of the upwelled waters off Spanish Sahara p0041 A75-10645 STEREOPHOTOGRAPHY [E75-10022] p0030 N75-13338 p0041 A75-10645 Analysis of pseudocolor transformations of ERTS-1 An electronic drawing aid for map compiling p0027 A75-10076 mages of Southern California area --- geological faults and stereo-orthophotos SPECTRAL BANDS Hologrammetry - Concepts and applications A cloud physics investigation utilizing Skylab data p0036 N75-13346 p0025 N75-15117 p0065 A75-15434 [E75-10037] [E75-10099] Terrain measurement from holographic stereograms p0001 A75-16682 Remote sensing geophysics from Skylab --- near infrared SPECTRAL REFLECTANCE Mapping soil features from multispectral scann reflectance anomalies of andesites and basalts in Southern Automatic contouring on the Gestalt Photomapper 2-1 p0028 A75-18055 California and Nevada p0027 A75-15435 [E75-10051] p0036 N75-13351 Vegetation canopy reflectance p0001 A75-16241 Fault tectonics and earthquake hazards in the Peninsular Ranges, Southern California --- Gulf of California An evaluation of stereoscopic viewing of ERTS and Skylab On the natural limitations of target differentiation by p0060 N75-14220 means of spectral discrimination techniques --- ERTS remote [E75-10074] p0031 N75-14193 STEREOSCOPY p0054 A75-17132 Fault tectonics and earthquake hazards in the Peninsular Single flight stereo radar capabilities ilities --- for terrain p0028 A75-17201 Use of visible, near-infrared, and thermal infrared remote Ranges Southern California p0002 A75-17144 sensing to study soil moisture [E75-10108] p0039 N75-15124 An evaluation of stereoscopic viewing of ERTS and Skylab Grass canopy bidirectional spectral reflectance The availability of local serial photography in southern images p0003 A75-17170 STORMS (METEOROLOGY) California --- for solution of urban planning problems [NASA-CR-141314] p0025 N75-1 Dust storms according to data of space research --spaceborne photography of and zones
[NASA-TT-F-16021] p0021 N75-12463 p0025 N75-15128 Spectral reflectance studies on mineral deficiency in corn p0003 A75-17172 SOUTHERN HEMISPHERE niants Comparison of the Southern and Northern Hamisphere Investigation into the spectral signature of agricultural STRATIGRAPHY general circulation characteristics as determined by satellite crops during their state of growth Recordings of the spectral-radio-meter SIGNE during the summer of 1972

[FOA-2-C-2577-E1] p0072 N75-12283 Remote sensing geophysics from Skylab --- strationaphic analysis of weathered rocks [E75-10058] [UCRL-75556] p0017 N75-10638 p0037 N75-13358 SOYUZ SPACECRAFT STRATOSPHERE Spectrophotometric observations of earth from SPECTRAL SIGNATURES The 1973 program of measurement of the min Soyuz-13

Transference of ERTS-1 spectral signatures in tin

bus-5 microwave spectrometer

The effect of atmospheric water classification of ERTS data

Effect of atmospheric haze and s classification of ERTS-1 date

Signatures of various earth surfaces measured by the

p0066 A75-17111

p0066 A75-17119

vanor on automatic

p0013 A75-17131

n angle on automatic p0014 A75-17157

[E74-10808]

p0016 N75-10535

p0018 N75-11523

constituents of the stratosphere using the Concorde 001 p0063 A75-11720

Quantitative determination of stratospheric aerosol

Measurement of trace stratospheric gases by inverting

thermal infrared limb radiance profiles

Quantitative determination of stratospheric aerosol SURFACE TEMPERATURE ERTS-B (Earth Resources Technology Satellite) ---Some features of the urban environment of Tokyo by remote sensing --- infrared surface temperature acecraft design remote sensor description, and technology p0020 N75-12393 urface temperature p0012 A75-17102 utilization Remote sounding of stratospheric temperatures using high resolution radiance measurements from the IRIS-D [NASA-CR-141130] p0022 N75-13452 Environmental definition program — satellite monitoring of the stratosphere and troposphere p0079 N75-13370 measurements [NASA-NEWS-RELEASE-74-329] The Coho Project: Living resources prediction feasibility study. Volume 3: System evaluation report [PB-234059/4] p0044 N75-10683 New methods for studying the dynamics of the entire arth p0016 A75-18108
Problems and consequences of modern hypotheses on SURFACE WATER p0022 N75-13458 [UCID-16565] the global tectonics for geodesy and gravimetry Surface water modeling Everglades Water Basin, p0028 A75-18109 Quantitative determination of stratospheric aerosol Applications of satellite geodesy in the exploration of characteristics [E74-10814] p0049 N75-10540 the internal structure of the earth p0029 A75-181: Lineaments in basement terrane of the Peninsular Range [F75-10079] p0023 N75-14198 p0029 A75-18123 Mapping of water bodies in northern Germany from ERTS pes p0051 N75-14227 STRATOSPHERE RADIATION tapes Nimbus 4 observations of changes in total ozone and Southern California stratospheric temperatures during a sudde udden warming p0012 A75-14841 p0035 N75-11407 Measurements of meteorological parameters and ai Fault tectonics and earthquake hazards in the Peninsular pollution spread over Stockholm, Sweden, by means of remote sensing and airborne instrumentation Remote sounding of stratospheric temperatures using high resolution radiance measurements from the IRIS-D [NASA-CR-141130] p0022 N75-13452 Ranges, Southern California [E75-10005] p0035 N75-11408 p0018 N75-11531 Experiment to evaluate feasibility of utilizing Skylab-EREP remote sensing data of tectonic analysis of the Bighorn Mountains region, Wyoming-Montana --- Black Hills, South Processing of ERTS CCT imagery of Sweden. A pilot udy p0060 N75-14219 STRIP MINING Automatic mapping of strip mine operations from Dakota/Wyoming pacecraft data p0034 A75-18065 Geographic applications of ERTS-1 imagery to landscape spacecraft data SWITZERLAND Methodological aspects and regional examples of snow-cover mapping from ERTS-1 and EREP imagery of [E75-10014] p0030 N75-12395 --- Mississippi River and Great Smoky Mountains Fault tectonics and earthquake hazards in the Peninsular nessee and North Carolina Ranges, Southern California p0051 N75-14232 [E74-10810] p0034 N75-10537 SYNCHRONOUS SATELLITES [E75-10022] p0030 N75-13338 Fault tectonics and earthquake hazards in the Peninsular Ranges, Southern California --- Gulf of California [E75-10074] Investigation of transient earth resources phenomena:
Continuation study
[NASA-CR-139163] p0073 N75-15131 Automated strip mine and reclamation mapping from ERTS --- Ohio coal p0035 N75-10556 Applicability of Skylab remote sensing for detection and monitoring of surface mining activities --- strip mining and reclamation in Kentucky and Tennessee SYNOPTIC METEOROLOGY

A satellite view of weather systems over the North ERTS-1 data on Afar tectonics p0031 N75-14239
Fault tectonics and earthquake hazards in the Peninsular [E75-10038] p0044 N75-12407 American Arctic p0063 A75-10790 Ranges, Southern California [E75-10108] Applicability of Skylab remote sensing for detection and Rainfall estimations from geosynchronous satellite pagery p0047 A75-13596 n0039 N75-15124 monitoring of surface mining activities — evironmental effects and reclamation in Tennessee and Kentucky [E75-10085] p0038 N75-15103 TELEVISION EQUIPMENT SYNTHETIC ARRAYS Rapid stock assessment of pilchard populations by p0003 A75-17168 Performance of a hologram matrix radar rcraft-borne remote sensors A demonstration of ERTS-1 analog and digital techniques applied to strip mining in Maryland and West Virginia [NASA-TM-X-70798] p0039 N75-15133 p0065 A75-14792 TELEVISION SYSTEMS P0065 A75-14792

SYSTEMS ANALYSIS

ERTS 1 flight evaluation report, 23 October 1973 - 23

January 1974 --- orbits 6371 to 7651

[NASA-CR-140712]

ERTS 1 flight evaluation report, 23 January - 23 April 1974 --- orbits 7652 to 8907

[NASA-CR-140810]

ERTS 1 flight evaluation report, January 23, 1973 to April 23, 1973 --- analysis of ERTS-1 systems during orbits 7800-1810 Space iconics [NASA-TT-F-798] n0057 N75-11419 Geometric distortions of opticommercial panoramic devision systems p0057 N75-11425 STRUCTURAL BASINS A comparison of ERTS imagery with conventional aerial photography for land-resource surveys in less developed countries. Examples from the Rift Valley Lakes Basin, Ethiopia p0061 N75-14249 television systems Geographic correlation of television pictures obtained om weather satellites p0071 N75-11432 Geographical correlation of TV and IR images obtained from weather satellites STRUCTURAL PROPERTIES (GEOLOGY)

Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State --- Lake Ontario from weather satellites --- cloud cover photography
p0071 N75-11433 System for electronic transformation and geographic correlation of satellite television information --- cloud cover photography p0071 N75-11434 2600-3810 p0034 N75-10536 [NASA-CR-140809] p0077 N75-10548 ERTS 1 flight evaluation report, 23 April 1974-23 July 1974 --- orbits 8908 to 10182 [NASA-CR-140711] p0077 N75-10568 Mineral resources, geological structures, and landf p0035 N75-10561 Experience in integrated geological and geographical interpretation of space photographs obtained in the USSR Lineaments in basement terrane of the Peninsular Ranges. SYSTEMS ENGINEERING Southern California p0036 N75-11437 [E75-10003] Earth resources sensor data handling system: NASA JSC version p0035 N75-11407 Use of TV pictures obtained from weather satellites to Geological and geophysical studies from space and respects for their development p0035 N75-11436 study the geological structure of the earth INASA-CR-1403341 prospects for their development p0035 N75-11436
Use of TV pictures obtained from weather satellites to p0072 N75-12418 D0036 N75-11438 TEMPERATURE DISTRIBUTION study the geological structure of the earth Theoretical investigations on the determination of the vertical temperature distribution in the atmosphere assuming multispectral thermal radiation measurements at T p0036 N75-11438 Mapping of the major structures of the African rift system --- Ethiopia, Somalia, and Yemen [E75-10024] TARGET RECOGNITION 60 GHz p0030 N75-13340 Mathematical modeling for the detection of fish by an rborne laser p0065 A75-15953 [BMFT-F8-W-74-04] p0024 N75-14331 Analysis of pseudocolor transformations of ERTS-1 airborne laser TEMPERATURE EFFECTS On the natural limitations of target differentiation by images of Southern California area --- geological faults and Use of visible, near-infrared, and thermal infrared rem means of spectral discrimination technique ques --- ERTS remote p0054 A75-17132 sensing to study soil moisture p0002 A75-17144 [E75-10037] DO036 N75-13346 sensors TEMPERATURE MEASUREMENT TASMANIA Preliminary results in the comparison of Skylab, ERTS Remote sounding of stratospheric temperatures using high resolution radiance measurements from the IRIS-D [NASA-CR-141130] p0022 N75-13452 Australian and Soviet gravity surveys along the Australian and RB-57 images for the detection of lineaments and fractures in Precambrian, Paleozoic, and late Tertiary rocks on and near the Colorado Plateau, North-Central Arizona calibration line p0031 N75-13454 Measurements of the dielectric properties of sea water p0037 N75-13360 TECHNOLOGICAL FORECASTING [E75-10060] at 1.43 GHz U.S. energy resources - Outlook for the future The Great Basin investigation [NASA-CR-2458] p0075 A75-12999 [E75-10076] p0037 N75-14195 Design data collection with Skylab/EREP microwave instrument S-193 --- Texas, Utah, Illinois, and Minnesota [E75-10080] p0072 N75-14199 The potential benefits for development derived from space Geological features on ERTS-1 images of a test area in the west-central Alps --- Italy p0037 N75-14237 applications [IAF PAPER 74-147] [E75-10080] p0075 A75-13761 Geologic evaluation of central Italy from ERTS-1 and cylab data p0038 N75-14238 TEMPERATURE PROFILES TECHNOLOGY ASSESSMENT
The progeny of Skylab - The pioneering phase of our Skylab data Measurement of trace stratospheric gases by inverting thermal infrared limb radiance profiles Preliminary evaluation of ERTS-1 imagery of Central rica p0060 N75-14242 expansion into space [AIAA PAPER 74-1216] p0018 N75-11523 p0063 A75-12031 The Great Basin investigation --- photomapping lithology Energy development: Proceedings of the Energy Sources Conference, Anaheim, Calif., July 14-19, 1974 Remote sounding of stratospheric temperatures using and geological structures [E75-10102] high resolution radiance measurements from the IRIS-D [NASA-CR-141130] p0022 N75-13452 p0038 N75-15120 p0075 A75-12986 Geologic structure in California: Three studies with Comparison of existing earth resources data processing TENNESSEE Geographic applications of ERTS-1 imagery to landscape change ... Mississippi River and Great Smoky Mountains of Tennessee and North Carolina ERTS-1 imagery [NASA-TM-X-70799] [IAF PAPER 74-144] p0053 A75-13760 p0039 N75-15132 Electronics in meteorological instrum SULFATES p0065 A75-14726
First results from the crop identification technology [E74-10810] n0034 N75-10537 A technique for measurement of sulphur dioxide and sulphate concentrations from an aircraft techniques in the ERTS-1 imagery interpretation assessment for remote sensing / CITARS/ Tennessee Valley --- land use and soil mapping [E74-10811] p0016 N75-10538 p0018 N75-11498 p0004 A75-17175 Aircraft measurements of the concentration and size The application of remote sensing techniques: Technical distribution of atmospheric sulfate particles Hydrologic significance of lineaments in central and methodological issues
[NASA-CR-140674] p0057 N75-11504 p0070 N75-10564 [E75-10030] SULFUR OXIDES p0050 N75-12401 TECHNOLOGY UTILIZATION Can spinoffs from space provide water for a thirsty p0047 A.75-13578 Applicability of Skylab remote sensing for detection and monitoring of surface mining activities --- strip mining and reclamation in Kentucky and Tennessee A technique for measurement of sulphur dioxide and sulphate concentrations from an aircraft world p0018 N75-11498 The potential benefits for development derived from space p0044 N75-12407 SURFACE PROPERTIES [E75-10038] applications Remote sensing geophysics from Skylab --- solar heating ground surface and infrared reflectance of rock [IAF PAPER 74-147] p0075 A75-13761 Applicability of Skylab remote sensing for detection and AF PAPER 74-147]
Evaluation of ERTS data utilization in developing p0055 A75-17193 monitoring of surface mining activities --- evironmental effects and reclamation in Tennessee and Kentucky [E75-10057] DO038 N75-15103 p0037 N75-13357 An orbiting visible/infrared spectrometer for terrestrial. [E75-10085] TENNESSEE VALLEY (AL-KY-TN)
ERTS-1 imagery interpretation techniques in the Tennessee Valley --- land use and soil mapping [E74-10811] p0016 N75-10538 SURFACE ROUGHNESS EFFECTS atmospheric and oceanographic applications --- on Skylab Multi-frequency radiometric measurements of foam and a mono-molecular slick --- microwave sensing of ocean surface p0068 A75-17208 p0068 A75-17212

An overview of Texas activities in remote sensing p0076 A75-17235

SUBJECT INDEX The ERTS-1 investigation. Volume 7: Erts-1 land-use THERMOLUMINESCENCE TERRADYNAMICS nalysis of the Houston area test site Microbial abundance and thermoluminescence antarctic dry valley soils p0005 A75-17 New methods for studying the dynamics of the entire (E75-10009) p0016 A75-18108 n0021 N75-13334 Irrigation se cheduling, freeze warning and soil salinity THERMOSPHERE Problems and consequences of modern hypotheses on Morning-evening asymmetry of the thermosphere during geomagnetic disturbance p0011 A75-1110 the global tectonics for geodesy and graving detecting --- Cameron County, Texas pravimetry p0028 A75-18109 75-10050] p0050 N75-13350 The ERTS-1 investigation (ER-600). Volume 2: ERTS-1 [E75-10050] p0011 A75-11103 TIMBER IDENTIFICATION TERRAIN ANALYSIS coastal/estuarine analysis --- Galveston Bay, Texas Performance comparison of techniques for obtaining stereo radar images p0063 A75-10075 [NASA-TM-X-58118] p0051 N75-13371
Design data collection with Skylab/EREP microwave instrument S-193 --- Utah and Texas Rad/Scat cover types, and forest types by digital processing ERTS-1 data p0054 A75-17110 Terrain measurement from holographic stereograms p0001 A75-16682 TIMBER VIGOR asurements Inventory of forest and rangeland resources, including forest stress --- Colorado, Georgia, and Black Hills, South Four channel simultaneous X-L band imaging SAR dar p0054 A75-17118 p0059 N75-14189 [E75-10070] Detection of moisture and moisture from Skylab --- Texas and Kansas On the natural limitations of target differentiation by [E75-10032] DOOO7 N75-12402 n0008 N75-14192 means of spectral discrimination techniques --- ERTS remote [F75-10073] p0054 A75-17132 TOGO collection with Skylab/EREP microwave Design data ERTS image contribution to geological knowledge prering different types of climates (France, Togo, Dahomey, ladagascar) p0038 N75-14240 Use of ERTS-1 imagery for land evaluation in Pennington instrument S-193 --- Texas, Utah, Illinois, and Minnesota p0072 N75-14199 [E75-10080] County, South Dakota p0055 A75-17135 775-10080]
Irrigation scheduling, freeze warning and soil salinity attecting — Cameron and Starr County, Texas
775-10084] Madagascar) The use of remote sensing and natural indicators to delineate floodplains - Preliminary findings TOPOGRAPHY [E75-10084] Hologrammetry - Concepts and applications p0067 A75-17142 p0065 A75-15434 Irrigation scheduling, freeze warning and soil salinity steeting --- Cameron and Starr Counties, Texas Grass canopy bidirectional spectral reflectance p0003 A75-17170 The use of remote sensing and natural indicators to delineate floodplains - Preliminary findings [E75-10109] p0010 N75-15125 Coastal wetlands analysis from ERTS MSS digital data DO067 A75-17142 TEXTURES p0048 A75-17178 and field spectral measurements Texture analysis with Fourier series --- for crop Color terrain map of Yellowstone National Park, computer-derived from ERTS MSS data identification from color aerial photographs surface topography p0004 A75-17187 Terrain properties and topography from Skylab p0004 A75-17185 Use of feature extraction techniques for the texture and Inventories of Delaware's coastal vegeteurs utilizing digital processing of ERTS-1 imagery p0004 A75-17186 context information in ERTS imagery: Spectral and textural processing of ERTS imagery --- classification of Kansas [E75-10059] p0030 N75-13359 Investigation of accuracy of determination of plane land use [E75-10011] DO059 N75-13335 Radar cross sections of vegetation canopies determined by monostatic and bistatic scatterometry THAILAND p0005 A75-17190 airborne rangefinder ERTS applications in Thailand - A progress report [AD-785886] p0076 A75-17123 The usefulness of imaging passive microwave for rural The next problems in the area of the implementation of urban terrain analysis p0055 A75-17199 Thailand national programme of the Earth Resources planning for cartographical-geodetic work
[AD-785882] nO Technology Satellite [E74-10806] Microwave radiometric characteristics of snow-covered p0077 N75-10533 p0049 A75-17200 terrain The Thailand programme of the Technology Satellite [E74-10807] p TRACE CONTAMINANTS Earth Resources Single flight stereo radar capabilities --- for terrain p0028 A75-17201 p0077 N75-10534 The use of space photos for search of oil and gas fie THEMATIC MAPPING p0034 A75-17229 Automatic contouring on the Gestalt Photomapper 2-1 Measurement of trace stratospheric gases by inverting Small scale thematic mapping - A case for radar n0028 A75-18062 thermal infrared limb radiance profiles n0028 A75-18055 THERMAL EMISSION p0018 N75-11523 The Skylab S-1908 Earth Terrain Camer. p0069 A75-18071 Thermal emission of agglomerations as recorded on TRANSFER FUNCTIONS influence of the atmosphere on spectral radiance and contrasts of natural formations measured from space --nvironmental satellite imagery --- of European urban Early results from the experimental Automated Image Data Extraction System --- for aerial terrain sensing p0069 A75-18072 [IAF PAPER 74-139] n0011 A75-13757 using data from Soyuz and Salyut spacecraft
p0019 N75-11557 THERMAL MAPPING An extension of the combinatorial geometry technique for modeling vegetation and terrain features
[AD-782883] Scanning thermal plumes --- from power plant TRANSIENT RESPONSE DO014 A75-17162 p0006 N75-10577 A generator of high-power current pulses of semisinusoidal shape for an aerial variant of the method of transient processes --- for aerial prospecting Thermal behaviour of some rice fields affected by a The PUCE programme for terrain evaluation for yellows-type disease p0004 A75-17174
Emissivities and forward scattering of natural and engineering purposes. 2: Procedures for terrain DO053 A75-13310 [PB-233847/3] p0029 N75-10625 man-made material at three millimeter wavelength p0067 A75-17197 TRANSMISSION FEEICIENCY Use of a digital computer to construct isoline maps based on a stereomodel of terrain [AD-784170] p0057 N75-11453 More information and less data from space --- data reduction in remote sensing systems p0063 A75-11730 Deteriorating effects on 3 mm wave passive imagery TRANSMITTANCE Terrain properties and topography from Geologic interpretation of infrared thermal images in East Qatrani area, Western Desert, Egypt p0033 A75-17215 constituents in the atmosphere by satellite [E75-10059] p0030 N75-13359 Automated selection and location of terrain features on Statistical separability of agricultural cover types in subsets of one to twelve spectral channels TRIANGULATION p0005 A75-17216 the basis of spectral characteristics Earth parameters from global satellite triangulation and trilateration p0064 A75-12985 p0030 N75-13375 Remote sensing techniques applied to the study of Italian oldanic areas - The results of the repetition of the airborne The remote identification of terrain features and materials at Pennsylvania test sites: An investigative study of photography --- North Carolina and Virginia [E74-10804] I.R. survey compared to the previous data techniques [PB-235992/5] p0069 A75-17223 p0031 N75-14263 Skylab: A proposal aerotriangulation with very small scale photography --- North Carolina and Virginia Applications of remote sensing by the state of p0069 A75-17232 TERRESTRIAL RADIATION The earth as a radio source - Terrestrial kilometric [E75-10101] Automatic pattern recognition and mapping utilizing ermal infrared imagery p0070 A75-18074 p0011 A75-11211 radiation TROPICAL METEOROLOGY thermal infrared imagery The radio emission of the earth as a planet A remote sensing study of Pacific hurricane Ava An application of infrared remote sensing techniques to book n0076 A75-15319 p0012 A75-17112 ecological problems [AD-786028] TROPICAL REGIONS A hydrogeomorphic approach to evaluating flood potential in central Texas from orbital and suborbital remote sensing p0022 N75-13419 ERTS image contribution to geological knowledge covering different types of climates (France, Togo, Dahomey, Madagascar) p0038 N75-14240 rbital remote sensing p0047 A75-17140 THERMAL POLLUTION Application of remote sensing for prediction and detection imagery of thermal pollution Moisture detection from Skylab --- soil studies for Texas TROPOSPHERE nd Kansas sites p0002 A7
An overview of Texas activities in remote sens [NASA-CR-139182] p0002 A75-17145 p0017 N75-10572 Feasibility of remote sensing for detecting thermal pollution. Part 1: Feasibility study. Part 2: Implementation of the stratosphere and troposphere [UCID-16565] p0076 A75-17235
Land use classification with ERTS-1 data in the Houston dan --- coastal ecology THERIDITY p0015 A75-18059 [NASA-CR-134453] p0025 N75-15199 Remote sensing of western Lake Superior An evaluation of the suitability of ERTS data for the THERMAL RADIATION p0048 A75-17181 purposes of petroleum exploration --- Anadarko Basin in Oklahoma and Texas p0035 N75-10553 Geothermal reconnaissance from quantitative and thermal infrared images p0056 A75-17218 Land use and mapping p0017 N75-10560 Correlation of missions 191, 51M and helicopter

Contribution of space platforms to a ground and airborne sensing programme over active Italian volcanoes D0691 p0037 N75-14188

Theoretical investigations on the determination of the vertical temperature distribution in the atmosphere assuming multispectral thermal radiation measurements at

[BMFT-FB-W-74-04] n0024 N75-14331 Analysis of Skylab MSS data, and calibration of thermal over Lake Monroe, Indiana

[E75-10095] p0073 N75-15113

THERMODYNAMIC PROPERTIES

photography --- aerial photography and mapping of three areas in Test Site 175 in Texas

Develop techniques and procedures, using multispectral

systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil --- Southern

Irrigation scheduling, freeze warning and soil salinity detecting --- Cameron County, Texas

Detection of moisture and moisture related phenomena

from Skylab --- radiometer and scatterometer analysis of

p0058 N75-12279

p0006 N75-12394

p0007 N75-12414

n0007 N75-13333

[NASA-CR-134289]

Texas

[E75-10012]

Texas site

[E75-10007]

5-10046]

Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil p0007 N75-12403 [E75-10033]

p0005 A75-17973

Classification and mapping of coal refuse, vegetative

The gravity field and the definition of stationary sea p0042 A75-18124

coordinates by means of the Central Scientific Research Institute of Geodesy, aerial photography and cartography

p0061 N75-14255

p0032 N75-15143

The 1973 program of measurement of the minor constituents of the stratosphere using the Concorde 001 DO063 A75-11720

The potentialities for determining content of minor gas

DO015 A75-17932

Skylab a proposal aerotriangulation with very small scale

p0070 N75-10532

p0073 N75-15119

Environmental definition program --- satellite monitoring

p0022 N75-13458

Dynamics of suspended sediment plumes in Lake n0049 N75-10541

[E74-10815] Radiative transfer of visible radiation in turbid p0019 N75-11552 atmosphara

U

U.S.S.R.

Development of research in the USSR on the use of redar images for geological purposes p0033 A75-17126 Elements of the deep structure of the earth's crust on the space images of the East Caucasus

p0028 A75-17228 Soviet satellite geodesy: Basis and status of Soviet laser geodesy and long-base line interferometry [AD-781138/3] p0029 N75-10524

U.S.S.R. SPACE PROGRAM Overall evaluation of ERTS-1 imagery for cartographic U.S., and U.S.S.R. p0029 N75-10528 application --- Australia, Antarctica, U.S., [E74-10800] Orbital geology --- an aid to mineral exploration and mapping the earth's surface [NASA-TT-F-16000] p0035 N75-10887 Experience in integrated geological and geographical interpretation of space photographs obtained in the USSR p0036 N75-11437 Australian and Soviet gravity surveys along the Australian [BULL-161] o0031 N75-13454 Preliminary data on the UHF radiometric research of soil retness, conducted in the USSR, in 1974 [NASA-TT-F-16151] p0009 N75-15097 Use of forest vegetation as an indicator in the interpretation and large scale mapping of soils in a forest [AD-785967] p0010 N75-15146 U.S.S.R. SPACE PROGRAM The earth in the lens of Salyut-3
[NASA-TT-F-16028] p0070 N75-10964 ULTRAHIGH FREQUENCIES Preliminary data on the UHF radiometric research of soil vetness, conducted in the USSR, in 1974 [NASA-TT-F-18151] p0009 N75-15097 ULTRAVIOLET RADIATION Crude and refined petroleum oil structured luminescence signatures induced by UV laser or lamp and their remote sensing applications
UNDERWATER OPTICS p0033 A75-17207 Mathematical modeling for the detection of fish by irborne laser p0065 A75-15953 UNITED STATES OF AMERICA Skytab and ERTS-1 investigations of coastal land-use and water properties in Delaware Bay [AIAA PAPER 74-1220] p0041 A75-12034 U.S. energy resources - Outlook for the future p0075 A75-12999 .Coastal and estuarine studies with ERTS-1 and Skylab p0047 A75-16240 Flood inundation in the southeastern United States from aircraft and satellite imagery p0047 A75-17138
Mapping of the 1973 Mississippi River floods by the p0028 A75-17139 p0047 A75-17138 Correlation of coastal water turbidity and current circulation with ERTS-1 and Skylab imagery Civil Works Mission --- for ecological monitoring application --- Australia, Antarctica, U.S., [E74-10800] p002 Geographic applications of ERTS-1 imagery to landscape change ... Mississippi River and Great Smoky Mountains of Tennessee and North Carolina

p0048 A75-17181 he application of ERTS-1 data to the Corps of Engineers p0015 A75-18064 Overall evaluation of ERTS-1 imagery for cartographic and U.S.S.R. p0029 N75-10528

[E74-10810] p0034 N75-10537 Skylab-EREP investigations of wetlands ecology --- New and Chesapeake Bay [E74-10816] p0049 N75-10542 Utility of ERTS for monitoring the breeding habit of migratory waterfowl --- United States and Canada p0006 N75-10557

Land use and mapping p0017 N75-10560 A survey for the use of remote sensing in the Chesapeake Bay region [NASA-CR-137476] p0043 N75-10571

The value of improved (ERS) information based on domestic distribution effects of U.S. agriculture crops [NASA-CR-141048] p0007 N75-12423

Improved (ERTS) information and its impact on U.S. markets for agricultural commodities: A quantitiative economic investigation of production, distribution and net xport effects [NASA-CR-141044] p0078 N75-12424

Preliminary results in the comparison of Skylab, ERTS and RB-57 images for the detection of lineaments and fractures in Precambrian, Paleozoic, and late Tertiary rocks on and near the Colorado Plateau, North-Central Arizona [E75-10060] New England reservoir management [E75-10065]

o0050 N75-13365 The Great Basin investigation

[E75-10076] p0037 N75-14195 Cartographic evaluation of Skylab-A S-192 scanner images --- Chesapeake Bay, Phoenix, and San Francisco

p0031 N75-14197 [E75-10078] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources, agriculture. Part 3: The integrated impact of improved IERS) information on US egricultural commodities

p0079 N75-14207 [NASA-CR-141262] Future US activities based on experience gained from ERTS-1 and EREP --- land use, forestry, egri-water-resources --- p0009 culture, and

p0009 N75-14253 Monitoring physical and chemical parameters of Delaware
Bay waters with an ERTS-1 data collection platform
[E75-10090] p0051 N75-15108 p0051 N75-15108

The Great Basin investigation --- photomapping lithology and geological structures [E75-10102] p0038 N75-15120 **UPPER ATMOSPHERE**

Remote sounding of stratospheric temperatures using high resolution radiance measurements from the IRIS-D p0022 N75-13452 [NASA-CR-141130] UPPER VOLTA

Earth Resources Technology Satellite-1 (ERTS-1) data and anthropology: Use of these data in carrying capacity estimates for sites in Upper Volta and Niger [E75-10001] p0078 N75-11406

Utilization of ERTS-1 imagery in cultivation and settlement site identification and carrying capacity estimates in Upper Volta and Niger [E75-10029] p0008 N75-13345

VELLING WATER
A satellite-eircraft thermal study of the upwelled waters
p0041 A75-10645 UPWELLING WATER off Spanish Sahara

Extensive summer upwelling on Lake Michigan during 1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160

URBAN DEVELOPMENT

Some findings on the applications of ERTS and Skylab imagery for metropolitan land use analysis p0013 A75-17136

URBAN PLANNING Recent progress in urban mapping p0015 A75-18054 Investigation of land use of northern megalopolis using

ERTS-1 imagery p0017 N75-10545 [NASA-CR-2459]

The availability of local aerial photography in southern California --- for solution of urban planning problems [NASA-CR-141314] p0025 N75-15126

URBAN RESEARCH Thermal emission of agglomerations as recorded on environmental satellite imagery --- of European urban

[IAF PAPER 74-139] p0011 A75-13757 Some features of the urban environment of Tokyo by mote sensing --- infrared surface temperature

urface temperature p0012 A75-17102 measurements Estimation of population density in Tokyo districts from ERTS-1 data p0076 A75-17103

Extraction of urban land cover data from multiplexed p0054 A75-17117 synthetic aperture radar imagery The usefulness of imaging passive microwave for rural

p0055 A75-17199 and urban terrain analysis UTAH

Experimental evaluation of atmospheric effects on radiometric measurements using the EREP of Skylab ---Bonneville Salt Flats, Utah [E75-10036] p0020 N75-12406

Geologic and mineral and water resources investigations in Western Colorado, using Skylab EREP data --- Utah [E75-10043] p0058 N75-12412 [E75-10043]

Design data collection with Skylab/EREP microwave instrument S-193 --- Utah and Texas Rad/Scat measurements p0059 N75-14189 [E75-10070]

Design data collection with Skylab/EREP microwave strument S-193 --- Texas, Utah, Illinois, and Minnesota p0072 N75-14199 instrument S-[E75-10080]

VALLEYS

Microbial abundance and thermolumine tarctic dry valley soils p0005 A75-17973 VAPOR PHASES

Airborne vapor surveillance system [AD-785585] p0021 N75-12459 VEGETATION

A comparison of Skylab and ERTS data for agricultural crop and natural vegetation interpretation

AIAA PAPER 74-1219] p0001 A75-12033 Coastal and estuarine studies with ERTS-1 and Skylab P0047 A75-16240 Vegetation canopy reflectance [AIAA PAPER 74-1219]

Classification and mapping of coal refuse, vegetative over types, and forest types by digital processing ERTS-1 lata p0054 A75-17110

Feasibility of using multiplexed SLAR imagery for water resources management and mapping vegetation communities p0054 A75-17115

ERTS-I data for classifying native plant com

ERTS-I data for Central Colorado p0004 A/5-1/1/c Prediction of the Fraunhofer Illuminescent materials --- crude and refined oils, effluents p0088 A/5-1/221 Remote sensing to detect the toxic effects of metals

on vegetation for mineral exploration p0034 A75-17228 Radar response to vegetation --- soil moisture mapping p0005 A75-18558 via microwave backscattering p0005 A75-18556
Developing processing techniques for Skylab data --vegetation and soil identification

p0057 N75-10527 An interdisciplinary analysis of multispectral satellite data: for selected cover types in the Colorado Mountains, using tomatic data processing techniques

[E74-10803] n0057 N75-10531 determine native anid plant distribution --- Arizona Regional Ecological Test Site [E74-10818] A study to explore the use of orbital remote se

An extension of the combinatorial geometry technique for modeling vegetation and terrain feature
[AD-782883] pOC

p0006 N75-10577 Study of recreational land and open space using Skylab nagery --- southeast Michigan

[E75-10034] p0020 N75-12404 Detection of moisture and moisture related phenomena from Skylab --- Kansas, Oklahoma, and Arkansas [E75-10035]

1E75-10U35] p0007 N75-12405
An interdisciplinary analysis of multispectral satellite data
for selected cover types in the Colorado Mountains, using
automatic data processing techniques ... geology,
hydrology, vegetation, and snow cover
[E75-10072] p0037 N75-14191 p0007 N75-12405

Detection of moisture and moisture related phenomena from Skylab --- Texas and Kansas

[E75-10073] p0008 N75-14192 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141287] p0080 N75-14211

Recognition and interpretation of spectral signatures of vegetation from aircraft and satellite imagery in Western Queensland, Australia p0008 N75-14241 Morphographic and morphometric analysis of vegetative

cover photographs in aerial photography for the mapping of vegetation [AD-785932]

Usefulness of Skylab color photography and ERTS-1 multispectral imagery for mapping range vegetation types in southwestern Wyoming p0010 N75-15115

[E75-10097] VEGETATION GROWTH

Inventories of Delaware's coastal vegetation and land-use Inventories of Detaware a cuester regulation utilizing digital processing of ERTS-1 imagery p0004 A75-17186

Radar cross sections of vegetation canopies determined by monostatic and bistatic scatterometry

p0005 A75-17190

VEHICULAR TRACKS

The use of color infrared imagery for the study of marsh buggy tracks o0061 N75-15127

VELOCITY FRRORS Analytical estimates of precision in determining parameters of motion of artificial earth satellite when

nalyzing imprecise measurements with correlated erro p0053 A75-11419 VENETUELA

ERTS-1 contributions to the ecogenetical knowledge of the Lianos of the Orinoco (Colombia and Venezuela)

p0023 N75-14245 VERTICAL MOTION

The effect of the secular change of the terrestrial grav field on elevation level differences p0028 A75-18122

VIRGINIA Coastal wetlands analysis from ERTS MSS digital data and field spectral measurements p0048 A75-17178

Skylab a proposal aerotriangulation with very small scale photography --- North Carolina and Virginia [E74-10804] p0070 N75-10532

Remote sensing applications in marine science progra VIMS [NASA-CR-137475] p0043 N75-10566

Urban and regional land use analysis: CARETS and tensus cities experiment package [E75-10019] p0020 N75-12398

Remote sensing geophysics from Skylab --- rationing-type algorithms to enhance multispectral data

algorithms to emission pool N/b-1338 [E75-10052] p0030 N/b-1338 Urban and regional land use analysis: CARETS and Urban and regional land use analysis: New Haven, cersus cities experiment package --- New Haven, Connecticut: Pontiac, Michigan: Phoenix, Arizona; District of Columbia

p0024 N75-15109 [E75-10091] Urban and regional land use analysis: CARETS and

census cities experiment package [E75-10100] p0025 N75-15118

Skylab: A proposal aerotriangulation with very small scale photography --- North Carolina and Virginia [E75-10101] p0073 N75-15119

VIRIRILITY The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficien

p0070 A75-19266 VISUAL PHOTOMETRY

Automated remote measurement system in agricultural meteorology --- for crop survey by aerial photom

p0064 A75-12947 **VOLCANOFS**

Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne I.R. survey compared to the previous data p0069 A75-17223

Contribution of space platforms to a ground and airborne remote-sensing programme over active Italian volcances

VOLCANOLOGY

Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcances p0037 N75-14188 [E75-10069]

VORTICES

Satellite photography of eddies in the gulf loop current p0041 A75-11255

W

WASHINGTON

Evaluate ERTS imagery for mapping and detection of changes of snowcover on land and on glaciers --- Alaska

p0049 N75-10539 [E74-10812]

Study to develop improved spacecraft snow survey methods using Skylab/EREP data --- Sierra Nevada, Cascades, and Great Plains p0073 N75-14200

WATER COLOR

Lake Ontario water mass delineation from ERTS-1 p0042 A75-17184

Ocean color, a three component system? p0044 N75-11411 [NASA-CR-139145]

WATER CURRENTS

Correlation of coastal water turbidity and current circulation with ERTS-1 and Skylab imagery p0048 A75-17181

WATER DEPTH

Water depth penetration film test p0047 A75-12873 Skylah water denth determination --- Lake Erie and Lake

Michigan [E75-10048]

WATER EROSION

Dynamics of suspended sediment plumes in Lake

[E74-10815]

p0049 N75-10541

WATER FLOW From aerospace photographs to forecasting and

calculating flows [NASA-TT-F-16006] DO059 N75-13367

WATER MANAGEMENT

Water-management models in Florida from ERTS-1

p0047 A75-13579 Coastal wetlands - Prospects for satellite inventory

p0047 A75-13580 Operational use of satellite and high altitude remote sensing for the generation of input data f ta for water demand p0067 A75-17148

Water resources --- monitoring and management from PTS-1 data p0043 N75-10559 ERTS-1 data

Utilization of EREP data in geological evaluation regional planning, forest management, and water management in North Carolina

p0008 N75-13363

New England reservoir management [F75-10065] p0050 N75-13365

WATER POLLUTION

The economic impact of remote sensing data as the source of nonpoint pollution monitoring and control g and control p0076 A75-17166

pollution with a lidar p0014 A75-17205 Remote measurements of water pollution

Oil slick detection by X-band synthetic aperture radar

The application of spectrometric and polarization techniques for remote sensing of oil on sea water

Laser Raman spectroscopy of solutes dissolved in water

from a remote platform [PB-233432/4] p0056 N75-10202 A study of a dual polarization laser backscatter system for remote identification and measurement of water

[NASA-CR-140665] p0017 N75-10563 Oil pollution detection, monitoring and law enforcement

isiana Gulf Coast waters p0022 N75-13343

An application of infrared remote sensing techniques to ecological problems [AD-786028] DO022 N75-13419

Investigation of the environmental change pattern of Japan

[E75-10086] p0024 N75-15104

Remote sensing of western Lake Superior

p0048 A75-17161 Correlation of multispectral imagery with water analysis - The Ross Barnett Reservoir Remote Sensing Project p0014 A75-17165

Computer derived coastal water classifications via correlation of coastal water turbidity and current

circulation with ERTS-1 and Skylab imagery

p0048 A75-17181 Boundaries of ERTS and aircraft data within which useful water quality information can be obtained

p0014 A75-17182 Lake Ontario water mass delineation from ERTS-1

Monitoring trophic status of inland lakes - A quantitative application of ERTS imagery p0049 A75-18067
Skylab study of water quality --- Kansas reservoirs and

lakes [E75-10075] p0051 N75-14194

WATER RECLAMATION

Can spinoffs from space provide w p0047 A75-13578

world WATER RESOURCES

Feasibility of using multiplexed SLAR imagery for water resources management and mapping vegetation p0054 A75-17115

Rapid stock assessment of pilchard populations by aircraft-borne remote sensors p0003 A75-17168 Microwave profiling of snowpack free-water content

p0049 A75-17950 Genlooic and mineral and water reso Colorado, using Skylab EREP in Western Co [E74-10802]

p0034 N75-10530 Surface water modeling Everglades Water Basin, Florida

[E74-10814] n0049 N75, 10540

water resources p0042 N75-10554
Utility of ERTS for monitoring the breeding habit of migratory waterfowl --- United States and Canada

p0006 N75-10557 Water resources --- monitoring and management from ERTS-1 data

ERTS-1 data p0043 N75-10559 Water survey of Canada: Application for use of ERTS-A for retransmission of water resources data

[E74-10805] p0049 N75-11409 Geologic and mineral and water resources investigations Western Colorado, using Skylab EREP data --

75-10043] p0058 N75-12412
Water survey of Canada: Application for use of ERTS-A [E75-10043] for retransmission of water resources data

[E75-10026] p0050 N75-13342 Water survey of Canada: Application for use of ERTS-A

retransmission of water resources data p0050 N75-13353

New England reservoir management [E75-10065] p0050 N75-13365 Geologic and mineral and water resources investigations

western Colorado, using Skylab EREP data p0037 N75-14190 Skylab study of water quality --- Kansas reservoirs and

lakes [E75-10075] p0051 N75-14194

The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 5: Inland water resources [NASA-CR-141269] p0080 N75-14209

[NASA-CR-141253] Future US activities based on experience gained from ERTS-1 and EREP --- land use, forestry, agriculture, and water resources p0009 N75-14253

Water resource and hazard planning report for the Clark Fork River Valley above Missoula, Missoula County,

[PB-236041/0] p0051 N75-14260 Monitoring physical and chemical parameters of Delaware Bay waters with an ERTS-1 data collection platform

p0051 N75-15108 Geologic and mineral and water resources investigations in western Colorado, using Skylab EREP data p0025 N75-15123

WATER TABLES Evaluation of ERTS-1 and aircraft data for assessing internal drainage in irrigated agriculture

n0002 A75-17143 WATER TEMPERATURE

A satellite-aircraft thermal study of the upwelled waters off Spanish Sahara p0041 A75-10645 Monitoring and predicting El Nino invasions --- equatorial Pacific Ocean-atmosphere interactions

Estimation of sea surface temperature from remote sensing in the 11to 13-micron window region

p0065 A75-14803

WATER VAPOR

vapor on automatic p0013 A75-17131 effect of atmospheric water classification of ERTS data

Ocean internal waves off the North American and African coasts from ERTS-1

Bistatic radar sea state monitoring [NASA-CR-137469] p0043 N75-10673 A simulation of synthetic aperture radar imaging of ocean

'[NASA-TM-X-72629] p0044 N75-12172

WATERSHEDS

to the location of p0066 A75-17141 Application of remote sensing to hydrologically active /source/ areas Automatic classification of eutrophication of inland lakes from spacecraft data p0048 A75-17164 A Skylab program for the International Hydrological

Decade (IHD) --- Lake Ontario Basin [F75-10039] p0050 N75-12408

WEATHER FORECASTING

Satellites and computer in the service of meteorology: WWW - The global weather observation on system p0053 A75-11274

Rainfall estimations from geosynchronous satellite

imagery p0047 A75-13598 Conference on Weather Forecasting and Analysis, 5th. St. Louis, Mo., March 4-7, 1974, Preprints

p0016 A75-18151 Observing and forecasting local effects from satellite data p0016 A75-18153 --- mesoscale cloud patterns

The Coho Project: Living resources prediction feasibility study, volume 1 --- meteorological forecasting of fish

[PB-234057/8] p0044 N75-10681 The Coho Project: Living resources prediction feasibility Volume 2: Environmental report

[PB-234058/6] p0044 N75-10682 WEATHERING

Remote sensing geophysics from Skylab --- stratigraphic analysis of weathered rocks

[£75-10058] p0037 N75-13358

WEST VIRGINIA

A demonstration of ERTS-1 analog and digital techniques applied to strip mining in Maryland and West Vir [NASA-TM-X-70798] p0039 N75-15 p0039 N75-15133

WETLANDS Coastal wetlands - Prospects for satellite inventory p0047 A75-13580

Coastal wetlands analysis from ERTS MSS digital data and field spectral measurements p0048 A75-17178 Remote sensing for regional resource analysis p0015 A75-18061

Optical film density values from color IR photography r wetland soils mapping p0005 A75-18073 for wetland soils mapping

Skylab-EREP investigations of wetlands ecology --- New Jersey and Chesapeake Bay

[E74-10816] p0049 N75-10542 Evaluation of satellite remote sensing and automatic data techniques for characterization of wetlands and coastal marshlands -- detection and measurement of accretion in estuary system of Atchafalaya River in southern

Louisiana [E75-10025] p0044 N75-13341

Evaluation of satellite remote sensing and automatic data techniques for characterization of wetlands and coastal marshlands --- Atchafalava River Basin, Louisians

[E75-10028] p0050 N75-13344

WHALES

BESMEX: Bering Sea marine mammal experiment with the primary target species being the walrus and bowhead whale [NASA-TM-X-62399] p0008 N75-13503

WHITEOUT An examination of whiteouts and transmission through

ice fog in the visible and near infrared [DREO-TN-74-18] n0021 N75-12467

The use of high altitude photography and ERTS-1 imagery

for wildlife habitat inventory in Kansas p0005 A75-18066

Utility of ERTS for monitoring the breeding habit of migratory waterfowl --- United States and Canada

p0006 N75-10557 Utilization of Skylab (EREP) system for appraising

Utilization of Skylab (Enery system)
langes in continental migratory bird habitat
p0008 N75-13361 [E75-10061]

The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rang [NASA-CR-141263] p0080 N75-1 p0080 N75-14208

Utilization of Skylab (EREP) system for appraising changes in continental migratory bird habitat [E75-10087] p0009 p0009 N75-15105

The detection by ERTS-1 of wind-induced ocean surface features in the lee of the Antilles Islands

p0041 A75-10644 WIND EROSION

Use of ERTS-1 imagery to interpret wind-erosion hazard in the Sandhills of Nebraska

WIND EFFECTS

WIND RIVER RANGE (WY) Surface compositional mapping in the Wind River Range and Basin, Wyoming by multispectral techniques applied

p0059 N75-13347

WIND VELOCITY MEASUREMENT

The wind vector in a layer over water

n0041 A75-15179 Application of a single laser Doppler system to the

measurement of atmospheric winds [NASA-TM-X-64891] WISCONSIN

Monitoring trophic status of inland lakes - A quantitative indication of ERTS imagery p0049 A75-18067 application of ERTS imagery WORLD DATA CENTERS

Satellites and computer in the service of meteorology: WWW - The global weather observation system p0053 A75-11274

WYOMING Color terrain map of Yellowstone National Park, computer-derived from ERTS MSS data

p0004 A75-17185 Surface compositional mapping in the Wind River Range and Basin, Wyoming by multispectral techniques applied p0028 A75-17224

Mineral resources, geological structures, and landform surveys p0035 N75-1056* Experiment to evaluate feasibility of utilizing Skylab-EREF n0035 N75-10561

remote sensing data of tectonic analysis of the Bighorn Mountains region, Wyoming-Montana --- Black Hills. South

Dakota/Wyoming [E75-10014] o0030 N75-12395 Inventory of forest and rangeland resources, including forest stress --- Colorado, Georgia, and Black Hills, South

[E75-10032] p0007 N75-12402 Usefulness of Skylab color photography and ERTS-1 multispectral imagery for mapping range vegetation types in southwestern Wyoming

A-29

p0010 N75-15115



X RAY ASTRONOMY
Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667



YELLOWSTONE NATIONAL PARK (ID-MT-WY)
Color terrain map of Yellowstone National Park,
computer-derived from ERTS MSS data
p0004 A75-17185

- - -


Service Care State

PERSONAL AUTHOR INDEX

Earth Resources / A Continuing Bibliography (Issue 5)

OCTOBER 1975

Typical Personal Author **Index Listing**



Listings in this index are arranged alphabetically by personal author. The title of the document provides the user with a brief description of the subject matter. The report number helps to indicate the type of document listed (e.g., NASA report, translation, NASA contractor report). The page and accession numbers are located: beneath and to the right of the title, e.g., p0074 N75-15138. Under any one author's name the accession numbers are arranged in sequence with the AIAA accession numbers appearing first.

Α

ABDEL-AAL, R. M.

Spectral reflectance studies on mineral deficiency in corr p0003 A75-17172 nlants

ABDEL-HADY, M. A.

Geologic interpretation of ERTS-1 satellite images for st Aswan area, Egypt p0033 A75-17108 East Aswan area, Egypt Geologic interpretation of ERTS-1 satellite images for West Aswan area, Egypt p0033 A75-17109 Geologic interpretation of infrared thermal images in East Qatrani area, Western Desert, Egypt p0033 A75-17215 ABDEL-SAMIE, A. G.

Spectral reflectance studies on mineral deficiency in corn ants p0003 A75-17172 plants

ABRAM, P

The role of ERTS in the establishment and of a nationwide land cover information system [NASA-CR-141045]

p0020 N75-12425 ACKERMAN, M.

The 1973 program of measurement of the minor constituents of the stratosphere using the Concorde 001 p0063 A75-11720

ACTON, L L

Global measurements of air pollution from satellites p0018 N75-11521

ADAMS, M. J.

The role of lidar in environmental sen p0070 A75-19274

AEKERSTEN, S. I

Processing of ERTS CCT imagery of Sweden. A pilot p0060 N75-14219

AGREEN, R. W. A simulation of the San Andreas fault experiment p0033 A75-11068

AGREN. C. H. Recordings of the spectral-radio-meter SIGNE during the summer of 1972

p0072 N75-12283

[FOA-2-C-2577-E1]

ALDRICH, R. C. Inventory of forest and rangeland resources, including

n0007 N75-12402 [E75-10032]

ALEKSANDROV, N. N. Methods for the automatic recording of atmospheric dust

concentrations and some of the results obtained p0057 N75-11502

ALEXANDER, R.

Urban and regional land use analysis: CARETS and ensus cities experiment package [E75-10019] n0020 N75-12398 Urban and regional land use analysis: CARETS and

nsus cities experiment package [E75-10056] p0022 N75-13356

Urban and regional land use analysis: CARETS and census cities experiment package [E75-10091] p0024 N75-15109

Urban and regional land use analysis: CARETS and census cities experiment package

[E75-10100] p0025 N75-15118

ALISHOUSE, J. A cloud physics investigation utilizing Skylab data [E75-10021] p00020 N75-1

p0020 N75-12400 A cloud physics investigation utilizing Skylab data p0025 N75-15117

ALLEN, W., III The availability of local aerial photography in southern [NASA-CR-141314] p0025 N75-15126

ALSID, L J. Skylab-EREP investigations of wetlands ecology [F74-10816] p0049 N75-10542 ANDERLE, R. J.

Transformation of terrestrial survey data to Dopple p0054 A75-16569

ANDERSON, A. T.

A demonstration of ERTS-1 analog and digital techniques applied to strip mining in Maryland and West Virginia [NASA-TM-X-70798] p0039 N75-15133

ANDERSON, D.

New England reservoir management [E75-10065] p0050 N75-13365

ANDERSON, D. M.

Multidisciplinary applications of ERTS and Skylab data in Ohio DO015 A75-17234

ANDERSON, H. R.

Oil slick detection by X-band synthetic aperture radap0068 A75-17209

ANDERSON, J. R.

ERTS-1 and data for national land use planning p0075 A75-13581 Remote sensing and land use inventory and mapping p0015 A75-18057

ANDERSON, R.

Multispectral photography for earth resources [NASA-CR-141283] DO074 N75-15138

ANDERSON, R. R. Coastal wetlands - Prospects for satellite inventory p0047 A75-13580

Skylab-EREP investigations of wetlands ecology 74-10816] p0049 N75-10542 [E74-10816]

ANDREWS, J.

e economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Source document [NASA-CR-141193] p0079 N75-14204

The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity

Forestry, wildlife and rangela Volume 4: [NASA-CR-141263] DO080 N75-14208 ANDREYANOV, V. V

Fundamentals of the space indication of earth [NASA-TT-F-16108] p0074 N75-15136

ANGSUWATANA, P.

ERTS applications in Thailand - A progress report

p0076 A75-17123

APFL J R

Ocean internal waves off the North American and Africa coasts from ERTS-1 p0042 A75-1718 p0042 A75-17183 APPLEGATE, D. N.

A method of specifying remotely sensed units for soil p0002 A75-17106 sample points

Transference of ERTS-1 spectral signatures in time and p0066 A75-17111

ARDEN, W.

The remote Raman spectrometer is a viable instrument for remote sensing of the environme p0068 A75-17214

ARMSTRONG, A. C.

of ERTS data Generalisation for global-scale investigations DOOR1 N75-14224

ARNOLD, J. T.

Airborne vapor surveillance system
[AD-785585] p0021 N75-12459

ARNOLD, K.

Geodetic aspects of laser distance measurements to the moon and radio-interference measurements to quasars p0027 A75-14646

ARONSON, A. I.

Very high resolution radiometer p0064 A75-12109

ATKINS, D. H. F.

A technique for measurement of sulphur dioxide and sulphate concentrations from an aircraft p0018 N75-11498

ATTEMA F. P. W.

Radar cross sections of vegetation canopies determined by monostatic and bistatic scatterometry p0005 A75-17190

ATWELL B. H.

A review of initial investigations to utilize ERTS-1 data in determining the availability and distribution of living marine resources p0043 N75-10555

B. A remote sensing study of Pacific hurricane Ava p0012 A75-17112 AU. B.

Multi-frequency radiometric measurements of foam and p0068 A75-17208 a mono-molecular slick

AWALD, J. T.

A technology to renovate the search for new mineral p0063 A75-10434 deposits

В

BAILEY, P. L.

Measurement of trace stratospheric gases by inverting thermal infrared fimb radiance profiles o0018 N75-11523

Performance comparison of techniques for obtaining stereo radar images p0063 A75-10075

Single flight stereo radar capabilities p0028 A75-17201

BAKER, D.

Skylab. II - The three month vigil p0065 A75-13983 BAKER, V. R.

A hydrogeomorphic approach to evaluating flood potential in central Texas from orbital and suborbital r p0047 A75-17140 imagery

BALANIS, C. A.

Application and sensitivity investigation of Fourier transforms for microwave radiometric in p0070 N75-11310

[NASA-CR-140754] BALASUBRAMANIAN, N.

Coherent optics in mapping; Proceedings of the Tutorial Seminar and Technology Utilization Pro N.Y. March 27-29, 1974 ALDI € p0027 A75-16676

BALDI, G.

Thermal behaviour of some rice fields affected by p0004 A75-17174 yellows-type disease BALL M.

Applications of remote sensing by Nebraska pC by the state of p0069 A75-17232 BANNERT, D. ERTS-1 data on Afar tectonics p0031 N75-14239

BARLOW, B. C. Australian and Soviet gravity surveys along the Australian calibration line

[BULL-161] p0031 N75-13454

BARNES, J. C.

Study to develop improved spacecraft snow survey nethods using Skylab/EREP data p0073 N75-14200 [E75-10081]

BARNETT, T. L

Summary of flight performance of the Skylab Earth Resources Experiment Package / EREP/ p0076 A75-17211

BARR. J.

Applications of remote sensing Nebraska by the state of p0069 A75-17232 BARR, J. C.

Design data collection with Skylab/EREP microwave instrument S-193 [F75-10070] p0059 N75-14189

BARRICK, D. E.

Bistatic radar sea state monitoring

[NASA-CR-137469] p0043 N75-10673 BARTA, G. Applications of satellite geodesy in the exploration of

the internal structure of the earth p0029 A75-18123 BARTLE, E. R.

Global measurements of air pollution from satellites p0018 N75-11521

BARTLETT. D.

Skylab and ERTS-1 investigations of coastal land-use and water properties in Delaware Bay [AIAA PAPER 74-1220] p0041 A75-12034

Coastal and estuarine studies with ERTS-1 and Skylab

BASHARINOV, A. E. inventories of Delaware's coastal vegetation and land-use The use of remote sensing and natural indicators to BRUNELLE, D. N. utilizing digital processing of ERTS-1 imagery delineate floodolains - Preliminary findings The usefulness of imaging passive microwave for rural p0004 A75-17186 p0067 A75-17142 and urban terrain analysis p0055 A75-17199 BASHARINOV, A. E. BRUSCH, R. ROBIANI A The value of improved (ERS) information based on The radio emission of the earth as a planet Geological features on ERTS-1 images of a test are p0076 A75-15319 mestic distribution effects of U.S. agriculture crop the west-central Alps p0037 N75-14237 Passive microwave sensing of moist soils [NASA-CR-141046] p0007 N75-12423 BORODIN, L. F. p0066 A75-17124 BRYAN, B. A. Passive microwave sensing of moist soils BASHARINOV, A. Y. Correlation of missions 191, 51M and helicopter p0066 A75-17124 Preliminary data on the UHF radiometric research of soil retness, conducted in the USSR, in 1974 photography [NASA-CR-134289] Preliminary data on the UHF radiometric research of soil wetness, conducted in the USSR, in 1974 [NASA-TT-F-16151] p0009 N75-15097 p0058 N75-12279 p0009 N75-15097 [NASA-TT-F-16151] BRYAN, M. L. BASHILOVA, I. I. Extraction of urban land cover data from multiplexed nthetic aperture radar imagery p0054 A75-17117 RORODIN, N. L. Use of TV pictures obtained from weather satellites to synthetic aperture radar imagery Work dealing with densification of altimetric-planometric BUDENHOLZER, R. A. study the geological structure of the earth data points employing universal instruments and radio altimeter and statoscopic indications p0036 N75-11438 U.S. energy resources - Outlook for the future BATAMANOVSKII, B. A. p0075 A75-12999 p0061 N75-14257 Investigation of accuracy of determination of plane coordinates by means of the Central Scientific Research BULLRICH, K. BORRIELLO, L.
A clustering algorithm for classification Scattering and absorption from poly-dispersed aerosols unsupervised Institute of Geodesy, aerial photography and cartography p0019 N75-11553 unsupervised crop p0002 A75-17113 BUNNIK, N. J. J. p0061 N75-14255 [AD-785886] BOSLEY, R. J. crops during their state of growth BURNS, J. P. Investigation into the spectral signature of agricultural BATTRICK, B. T. Use of feature extraction techniques for the texture and context information in ERTS imagery: Spectral and textural p0004 A75-17189 European earth-resources satellite experiments [ESRO-SP-100] p0081 N7 p0081 N75-14216 processing of ERTS imagery [E75-10011] Recent progress in urban mapping p0015 A75-18054 o0059 N75-13335 Application of ERTS data to environmental inventory and First results from the crop identification technology BOUCHILLON, C. W.
Application of remote sensing to state and regional assessment for remote sensing /CITARS, p0004 A75-17175 BURNS, K. L. problems RAYMA R W [NASA-CR-140805] DO078 N75-11414 A study of the usefulness of Skylab EREP data for earth Investigation of microwave hologram techn resources studies in Australia [E75-10047] BOULANGER, Y. O.
Australian and Soviet gravity surveys along the Australian ram techniques for p0067 A75-17195 application to earth resources p0036 N75-12415 BURTON, T. BEAUMONT, T. E. calibration line Recognition and interpretation of spectral signatures of Design data collection with Skylab/EREP microwave p0031 N75-13454 [BULL-161] vegetation from aircraft and satellite imagery in Western Queensland, Australia p0008 N75-14241 instrument S-193 BOWHILL S. A [E75-10070] p0059 N75-14189 Investigations of the ionosphere by sp BUSCHMANN, E. BEHAR, J. V. p0065 A75-13868 Geodesy and physics of the earth: Proceedings of the Second International Symposium, Potsdam, East Germany, Airborne survey of major air basins in California BOWLING, J. L. p0011 A75-14360 Investigation of the utility of kinetically controlled and p0028 A75-18106 May 7-11, 1973. Parts 1 & 2 JSHUYEV, A. V. BELON, A. E. catalytic reactions to perform environmental determinations summary of ERTS data applications in Alask by parallel spectrophotometric analysis p0077 A75-17236 Geographic correlation of television pictures obtained [TID-26594] p0021 N75-13063 from weather satellites p0071 N75-11432 BEREZIN, A. M. BRADBURN. G. forest vegetation as an indicator in the BUZNIKOV, A. A. Airborne survey of major air basins in interpretation and large scale mapping of soils in a forest Spectrophotometric observations of earth from p0011 A75-14360 'Soyuz-13'
[IAF PAPER 74-156] [AD-785967] n0010 N75-15146 BRADFORD, D. F. p0065 A75-13765 The application of spectrometric BERTRAND, J. P. The value of improved (ERS) information based on and polarization Recent and present sedimentation complexity of the ution effects of U.S. agriculture techniques for remote sensing of oil on sea water Rhone delta and the French mediterranean coast (Golion p0007 N75-12423 [NASA-CR-141046] n0068 A75-17210 p0045 N75-14228 Influence of the atmosphere on spectral radiance and The economic value of remote sensing of earth resources BESSIS, J. L. from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources, agriculture. Part 2: Distribution effects contrasts of natural formations measured from space p0019 N75-11557 Orientation of methods of information processing for the control of terrestrial resources [IAF PAPER 74-140] BYRNE, R. J. D0053 A75-13758 p0079 N75-14206 [NASA-CR-141194] Remote sensing applications in marine science programs BIGELOW, R. C.
Prediction of the Fraunhofer line -+ VIMS BRAY, C. J. [NASA-CR-137475] DO043 N75-10566 detectivity The interpretation and use of false-colour infra-red and p0068 A75-17221 luminescent materials true colour photography of part of Argentina obtained by Skylark earth resources rockets p0067 A75-17192 C Investigation of radar discrimination of sea ice A comparison of the capabilities of ERTS-1 and the p0042 A75-17121 Skylark earth-resources rocket for resources sources surveying in p0023 N75-14250 BIRO, P. CABALLE, G. central Argentina e effect of the secular change of the Qualitative study of some ERTS-1 images of the Golion field on elevation level differences p0028 A75-18122 project. Ecological aspects of terrestrial The Coho Project: Living resources prediction feasibility p0023 N75-14248 study CAMERON, R. E. Ocean internal waves off the North American and African [PB-234057/8] p0044 N75-10681 Microbial abundance and thermole antarctic dry valley soils p0 coasts from ERTS-1 p0042 A75-17183 BROCKETT, R. M. p0005 A75-17973 RIAIRRAINS. A Skylab environmental and crew microbiology studie CAMPBELL C. W. A comparison of ERTS imagery with conventional aerial p0024 N75-14366 Computer and photogrammetric general land use study photography for land-resource surveys in less developed BROOKS J countries. Examples from the Rift Valley Lakes Basin, p0061 N75-14249 of central north Alabama An extension of the combinatorial geometry techn [NASA-TR-R-431] n0022 N75-13372 Ethiopia for modeling vegetation and terrain features CÂMPBELL, W. J. BLANCHARD, M. B. [AD-782883] p0006 N75-10577 A study of microwave emission properties of sea ice Use of visible, near-infrared, and thermal infrared remote BROOKS, R. L. AIDJEX 1972 p0041 A75-17120 sensing to study soil moisture p0002 A75-17144 Applicability of Skylab remote sensing for detection and monitoring of surface mining activities CANNON, P. J. BODECHTEL, J. Application of radar imagery to environ Geoscientific research with rockets and satellite [IAF PAPER 74-116] p0053 A7 mapping of Texas CANNON, T. W. [E75-10038] n0044 N75-12407 p0034 A75-17222 p0053 A75-13741 Applicability of Skylab remote sensing for detection and Evaluation of ERTS-1 data by analogue and digital chaigues p0059 N75-14218 monitoring of surface mining activities [E75-10085] A camera for photography of atmospheric particles from ircraft p0053 A75-13526 techniques 00038 N75-15103 aircraft Geologic evaluation of central Italy from ERTS-1 and BROSSIER R CAPOZZA, F. DO038 N75-14238 Skylab data Interpretation of ERTS-1 data for investigating the French clustering algorithm for classification p0002 A75-17113 BODENHEIMER, R. E. western coastal zones (Fralit program) ERTS-1 imagery interpretation techniques in the Tennessee Valley p0051 N75-14230 Digital land-use classification using an ERTS-1 image p0023 N75-14226 BROWN, C. W. [F74-10811] p0016 N75-10538 Laser Raman spectroscopy of solutes dissolved in water CARDONE, V. J. from a remote platform Preliminary analysis of Skylab RADSCAT results over BONRUD, L. O. p0056 N75-10202 the ocean [NASA-CR-141454] Correlation and registration of ERTS multispectral [PB-233432/4] BROWN, G. S. p0046 N75-14793 imagery [NASA-CR-134294] dications and scientific results Skylab altir n0058 N75-12420 Skylab altimeter applica [AIAA PAPER 74-1221] CARLSON, G. E. D0063 A75-12035 BOONYOBHAS, C. Performance comparison of techniques for obtaining stereo radar images ERTS applications in Thailand - A progress report p0063 A75-10075 An examination of whiteouts and transmission through p0076 A75-17123 Single flight stereo radar capabilities ice fog in the visible and near infrared [DREO-TN-74-18] p0028 A75-17201 BOOVARSSON, A p0021 N75-12467 Environmental studies of Iceland with ERTS-1 imagery CARLSON, M. BROWN, W. DO012 A75-17105 Applications of remote sensing by the state of Nebraska p0069 A75-17232 A remote sensing study of Pacific hurricane Ava BORDEN, F. Y. p0012 A75-17112 A method of specifying remotely sensed units for CARLSON, P. R. BRUKHANOV, V. sample points p0002 A75-17106 Surface currents along the California coast observed on Orbital geology [NASA-TT-F-16000]

ERTS imagery

CARNEGGIE, D. M.

p0035 N75-10887

p0038 N75-14244

numerical techniques for

Comparison of various

classification of multispectral data. Application to geology

p0048 A75-17180

p0003 A75-17169

Determining range condition and forage production potential in California from ERTS-1 imagery

Classification and mapping of coal refuse, vegetative

p0066 A75-17111

cover types, and forest types by digital processing ERTS-1 data p0054 A75-17110

Transference of ERTS-1 spectral sig

DOIRON, L. N. CLAYTON, K. M. DAVIDTS, D. Generalisation of ERTS data Application of remote sensing to state and regional for global-scale n0081 N75-14224 Reduction of information redundancy in ERTS-1 and investigations EREP data. A proposal for an experimental study [NASA-CR-140805] p0078 N75-11414 CLIFF, W. C. n0060 N75-14225 DAVIES, J. H. CARTER, V. Application of a single laser Doppler system to the measurement of atmospheric winds [NASA-TM-X-64891] Correlation spectroscopy techniques for the regional and global measurements of air pollution p0018 N75-11505 Coastal wetlands - Prospects for satellite inventory p0021 N75-12462 p0047 A75-13580 COFFMAN. J. L. Coastal wetlands analysis from ERTS MSS digital data DAVIS. D. D. Catalog of earthquake photographs p0072 N75-13455 p0048 A75-17178 Trace gas analysis of power plant plumes via aircraft spectral measurements COKER, A. E. measurement - 03, NOx, and SO2 che CARTER V P Water-management models in Florida from ERTS-1 Skylab-EREP investigations of wetlands ands ecology p0049 N75-10542 p0064 A75-13148 p0047 A75-13579 data DAVIS, M. R. Surface water modeling Everglades Water Basin, A study of the early detection of insect infestations and density/distribution of host plants CARTMILL, R. H. Florida Evaluation of satellite remote sensing and automatic data [E74-10814] n0049 N75-10540 p0008 N75-13362 techniques for characterization of wetlands and coastal [F75-10062] COLDRICK, J. R. A preliminary feasibility study on a synthetic aperture radar /SARSAT/ for earth resources surveys DAVISON, E. marshlands p0044 N75-13341 [E75-10025] Design data collection with Skylab/EREP microwave S-193 Evaluation of satellite remote sensing and automatic data p0067 A75-17194 instrument S-IE75-100701 p0059 N75-14189 techniques for characterization of wetlands and coastal DE BOER, T. A. marshlands Recognition and interpretation of spectral signatures of vegetation from aircraft and satellite imagery in Western Queensland. Australia p0008 N75-14241 Investigation into the spectral signature of agricultural roos during their state of growth p0004 A75-17189 [F75-10028] 50050 N75-13344 crops during their state of growth CASSINIS. R. Queensland, Australia Remote sensing techniques applied to the study of Italian pleanic areas - The results of the repetition of the airborne COLLIS R T H Radar cross sections of vegetation canopies determined Lidar observations of atmospheric partic by monostatic and bistatic scatterometry p0019 N75-11559 p0005 A75-17190 I.R. survey compared to the previous data p0069 A75-17223 COLWELL J. E. DE CAROLIS. C. Vegetation canopy reflectance o0001 A75-16241 Thermal behaviour of some rice fields affected by yellows-type disease p0004 A75-17174 Contribution of space platforms to a ground and airborne emote sensing programme over active Italian volcanoes Grass canopy bidirectional spectral reflectance p0004 A75-17174 n0003 A75-17170 [F75-10069] DE LOOR, G. P.
Investigation into the spectral signature of agricultural p0037 N75-14188 Contribution of space platforms to a ground and airborne A study to explore the use of orbital remote sensing to crops during their state of growth p0004 A75-17189
Radar cross sections of vegetation canopies determined remote-sensing programme over active Italian volcanoe determine native arid plant distribution [E74-10818] p0073 N75-14236 p0006 N75-10544 Agricultural resources investigations in northern Italy, puthern France and Madagascar p0009 N75-14252 by monostatic and bistatic scatterometry p0005 A75-17190 southern France and Madagascar Simultaneous active and passive microwave response CHAHINE, M. T. of the earth - The Skylab radscat experime Determining range condition and forage production potential in California from ERTS-1 imagery Remote atmospheric sensing with an airborne lase p0012 A75-17114 D0066 A75-16485 absorption spectrometer COOPER, S. n0003 A75-17169 CHANDLER, B. J. New England reservoir management [E75-10065] DEL TORO, G. A. The application of pattern classification techniques to DO050 N75-13365 An intensive natural resources inventory aids land use Skylark earth resources rocket imagery CORDES, E. H. planning and resources management in Puerto Ri p0056 A75-19268 Water-management models in Florida from ERTS-1 p0005 A75-18056 CHANG, D. T. data p0047 A75-13579 Experimental evaluation of atmospheric effects on Surface water modeling Everglades Water Basin, Measurements of the extraterrestrial solar radiant flux Florida iometric measurements using the EREP of Skylab from 2981 to 4000 A and its transmission through the earth's atmosphere as it is affected by dust and ozone [E74-10814] [F75-10036] n0020 N75-12406 n0049 N75-10540 COUTTS, D. A. CHANG, T. C. p0070 A75-19547 Australian and Soviet gravity surveys along the Australian A radio picture of the earth D0066 A75-16191 calibration line Interpretation of ERTS-1 data for investigating the French CHARNELL R L Ocean internal waves off the North American and African western coastal zones (Frafit program) COVOCORESSES, A. P. p0051 N75-14230 coasts from ERTS-1 D0042 A75-17183 Overall evaluation of ERTS-1 imagery for cartographic CHEOSAKUL P Thailand national programme of the Earth Resources Technology Satellite application Summary of flight performance of the Skylab Earth [F74-10800] a0029 N75-10528 Resources Experiment Package / EREP/ [E74-10806] p0077 N75-10533 p0076 A75-17211 Rapid stock assessment of pilchard oppulations by The Thailand programme of the Earth Resources DEN HOLLANDER, L. G. aircraft-borne remote sensors p0003 A75-17168 Radar cross sections of vegetation canopies determined by monostatic and bistatic scatterometry Technology Satellite [E74-10807] CRAM, R. p0077 N75-10534 The detection by ERTS-1 of wind-induced ocean surface CHESNOKOV, Y. M. 00005 A75-17190 features in the lee of the Antilles Islands Space iconics [NASA-TT-F-798] DEPLOEY, J. Preliminary evaluation of ERTS-1 imagery of Central p0060 N75-14242 p0041 A75-10644 p0057 N75-11419 CRANE, R. B. CHESWORTH F T Africa Adaptive processing of decision-directed Kalman filter MSS DESHLER, W. Ice particulates in the mesosphere p0012 A75-16752 data using a p0055 A75-17151 CHEUNG, M. An examination of the extent of fire in the grassland CRAWFORD, T. V. Automatic land use classification using Skylab S-192 and savanna of Africa along the southern sid Atmospheric release advisory capability: Research and multispectral data p0002 A75-17104 [AIAA PAPER 74-1224] p0064 A75-12036 DESPAIN, D. [UCRL-75644-REV-2] o0025 N75-15202 CHIGIREV, A. A. Color terrain map of Yellowstone National Park, computer-derived from ERTS MSS data Use of a digital computer to construct isoline maps based CRAWLEY, B. G. Automatic contouring on the Gestalt Photomapper 2-1 p0028 A75-18055 n a stereomodel of terrain p0004 A75-17185 AD-784170] DO057 N75-11453 DHOORE, J. CROUCH, L. W. Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 CHISMAN, H. An application of infrared remote sensing techniques to The potential of small-scale mapping from ERTS-1
nagery p0060 N75-14221 ecological problems DIETLEIN, L. F. imagery AD-7860281 p0022 N75-13419 CHRISTIE R J The Proceedings of the Skylab Life Sciences Symposium. CURLIN, J. W. e of remote sensing of earth resources volume 1 [NASA-TM-X-58154-VOL-1] National Environmental Policy Act of 1969. from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a nationwide p0081 N75-14358 Environmental indices, status of development pursuant to Sections 102(2)(B) and 204 of the act FRUSCIO, D. J.

An examination of whiteouts and transmission through [GPO-25-314] DO077 N75-10617 land cover information system ice fog in the visible and near infrared [DREO-TN-74-18] NASA-CR-141267] p0080 N75-14211 CURRAN, R. J. p0021 N75-12467 The detection of dust storms over land and water with CHUKHRAY, G. I. DILLINGER, A. E. Preliminary data on the UHF radiometric research of soil wetness, conducted in the USSR, in 1974 satellite visible and infrared measurements The effect of atmospheric water vapor on automaticlassification of ERTS data p0013 A75-1713 p0012 A75-16675 p0013 A75-17131 p0009 N75-15097 CURTIS, L. F. [NASA-TT-F-16151] . DILLMAN, R. D. Ground-truth measurements in relation to aircraft and satellite studies of agricultural land use and land classification in Britain p0009 N75-14251 CIHLAR, J.

Active microwave measurement of soil water content Unsupervised mapping of geologic features ar Ilifornia p0034 A75-17225
The remote identification of terrain features and materials California p0001 A75-16242 [NASA-CR-141685] CLAASSEN, J. D. at a California test site: An investigative study of CUSTANCE, N. D. E. Preliminary analysis of Skylab RADSCAT results over Recognition and interpretation of appearance in Western vegetation from aircraft and satellite imagery in Western p0008 N75-14241 techniques the ocean [NASA-CR-141454] B-235991/7] p0031 N75-14261 p0046 N75-14793 DIPAGLO, W. D. CLAASSEN, J. P.
Simultaneous active and passive microwave response Preliminary results in the comparison of Skylab, ERTS and RB-57 images for the detection of 'ineaments and fractures in Precambrian, Paleozoic, and lata Tertiary rocks CZARNECKI, W.

D

Fully automated compilation of gravity and geophysical

p0032 N75-15147

DALU. G.

p0012 A75-17114

p0016 N75-10469

p0042 A75-17177

maps [AD-785838]

of the earth - The Skylab radscat experiment

Modern trends in environmental testing

The application of ERTS-derived information to the regional land use planning process p0015 A75-18058

Computer derived coastal water classifications via

CLAPP, J. L.

CLARK A J. JR.

[SLA-73-5931]

spectral signatures

CLARK. D. K.

Estimation of sea surface temperature from remote sensing in the 11to 13-micron window region p0065 A75-14803

DOIRON, L N. The use of color infrared imagery for the study of marsh pool 1 N75-15127 buggy tracks

Skylab contributions to the future p0075 A75-13618

Evaluation of ERTS-1 data by analogue and digital

near the Colorado Plateau, North

on and near [E75-10060]

DISHER, J. H.

DITTEL R H.

techniques

p0037 N75-13360

p0059 N75-14218

DONZIGER, A.	EIDEN, R.	FAYMAN, D. L
The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service.	Scattering and absorption from poly-dispersed aerosols p0019 N75-11553	Simultaneous active and passive microwave response of the earth - The Skylab radscat experiment
Volume 2: Source document	EL GHAWABY, M. A.	p0012 A75-17114
[NASA-CR-141193] p0079 N75-14204 DORNBACH, J. E.	Geologic interpretation of ERTS-1 satellite images for	FEENBERG, D.
Land use classification with ERTS-1 data in the Houston	East Aswan area, Egypt p0033 A75-17108 Geologic interpretation of ERTS-1 satellite images for	The value of improved (ERS) information based on domestic distribution effects of U.S. agriculture crops
area test site p0015 A75-18059	West Aswan area, Egypt p0033 A75-17109	[NASA-CR-141046] p0007 N75-12423
DOSS, H. W. Investigation of microwave hologram techniques for	EL KASSAS, I. A.	FEGLEY, R.
application to earth resources p0067 A75-17195	Geologic interpretation of ERTS-1 satellite images for East Aswan area, Egypt p0033 A75-17108	Geophysical monitoring for climatic change. The NOAA program p0018 N75-11510
DOWDY, J. M. Multidisciplinary applications of ERTS and Skylab data	Geologic interpretation of ERTS-1 satellite images for	FEINBERG, E. B.
in Ohio p0015 A75-17234	West Aswan area, Egypt p0033 A75-17109	Application of ERTS-1 data to the protection and management of New Jersey's coastal environment
Remote sensing for regional resource analysis p0015 A75-18061	EL SHAZLY, E. M. Geologic interpretation of ERTS-1 satellite images for	p0042 A75-17233
DOYLE, F. L.	East Aswan area, Egypt p0033 A75-17108	FELLOWS, R. S.
Determination of points of entry for potential	Geologic interpretation of ERTS-1 satellite images for	Resource analysis applications in Michigan p0069 A75-17231
contaminants into limestone aquifers using thermal infrared imagery	West Aswan area, Egypt p0033 A75-17109 EL-SHAZLY, E. M.	FERGUSON, J. K.
[NASA-CR-120540] p0021 N75-12546	Geologic interpretation of infrared thermal images in East	Skylab environmental and crew microbiology studies
DRAEGER, W. C. Areal extent of snow in forested regions - A practical	Oatrani area, Western Desert, Egypt p0033 A75-17215	p0024 N75-14366 FERRIS. J. E.
estimation technique using ERTS-1 data	ELLIS, M. An overview of Texas activities in remote sensing	Investigation of microwave hologram techniques for
p0002 A75-17122 Usefulness of remote sensing techniques for the	p0076 A75-17235	application to earth resources p0067 A75-17195
environmental monitoring of the Sacramento-San Joaquin	ELSE, C. V. Electronics in meteorological instrumentation	FESENKO, A. I. Detection of synchronous changes in fluxes of energetic
Delta p0025 N75-15135 DRAHOVZAL, J. A.	p0065 A75-14726	electrons and gamma-quanta during a magnetic
The remote-sensing program of the Geological Survey	ELSTON, D. P.	disturbance p0064 A75-12317
of Alabama p0069 A75-17230	Preliminary results in the comparison of Skylab, ERTS and RB-57 images for the detection of lineaments and	FISCHER, K. Scattering and absorption from poly-dispersed aerosols
DRAKE, B. Feasibility of using multiplexed SLAR imagery for water	fractures in Precambrian, Paleozoic, and late Tertiary rocks	p0019 N75-11553
resources management and mapping vegetation	on and near the Colorado Plateau, North-Central Arizona [E75-10060] p0037 N75-13360	FISHER, L. T.
communities p0054 A75-17115 DRENNAN, D. S. H.	EMPLAINCOURT, J. L. G.	Interactive machine assessment of critical land resources using ERTS-1 data p0013 A75-17149
The interpretation and use of false-colour infra-red and	The remote-sensing program of the Geological Survey	FISHMAN, G. P.
true colour photography of part of Argentina obtained by Skylark earth resources rockets p0067 A75-17192	of Alabama p0069 A75-17230 ENSLIN, W. R.	Comparison of existing earth resources data processing
A comparison of the capabilities of ERTS-1 and the	Resource analysis applications in Michigan	facilities [IAF PAPER 74-144] p0053 A75-13760
Skylark earth-resources rocket for resources surveying in	p0069 A75-17231	FISHMAN, H.
central Argentina p0023 N75-14250 DREW, J.	ERADUS, W. J. Radar cross sections of vegetation canopies determined	The value of improved (ERS) information based on
Applications of remote sensing by the state of	by monostatic and bistatic scatterometry	domestic distribution effects of U.S. agriculture crops [NASA-CR-141046] p0007 N75-12423
Nebraska p0069 A75-17232 DREW, J. V.	p0005 A75-17190	FLORENSKII, P. V.
Satellite pictures can help guide Sandhills irrigation	ERB, R. B. The ERTS-1 investigation. Volume 7: Erts-1 land-use	Plutonic structure of the earth's crust on space images
[E75-10023] p0021 N75-13339 Use of ERTS-1 imagery to interpret wind-erosion hazard	analysis of the Houston area test site	p0027 A75-17125
in the Sandhills of Nebraska	[E75-10009] p0021 N75-13334 The ERTS-1 investigation (ER-600). Volume 2: ERTS-1	The use of space photos for search of oil and gas fields p0034 A75-17229
[E75-10044] p0059 N75-13347 DRISCOLL, R. S.	coastal/estuarine analysis	FONTANEL A.
ERTS-I data for classifying native plant communities -	[NASA-TM-X-58118] p0051 N75-13371	Comparison of various numerical techniques for
Central Colorado p0004 A75-17176	ESCHELBACH, G.	classification of multispectral data. Application to geology
Income of forces and consoland consumer including	Scattering and absorption from poly-dispersed aerosols	p0038 N75-14244
Inventory of forest and rangeland resources, including forest stress	Scattering and absorption from poly-dispersed aerosols p0019 N75-11553	p0038 N75-14244 FORMISANO, V.
forest stress [E75-10032] p0007 N75-12402	p0019 N75-11553 ESPARZA, F.	FORMISANO, V. Solar wind and geomagnetic activity
forest stress [E75-10032] p0007 N75-12402 DUBENSKIY, V. P.	p0019 N75-11553	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY. V. P. System for electronic transformation and geographic correlation of satellite television information	p0019 N75-11653 ESPARZA, F. Planning applications in east central Florida [£75-10015] p0021 N75-13337 ESTES, J. E.	FORMISANO, V. Solar wind and geomagnetic activity
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [875-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models p0067 A75-17148	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [£75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State
forest stress [E75-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [£75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models p0067 A75-17148 The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY. V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818]
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J.	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models p0067 A75-17148 The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [£75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S.	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J.	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E.
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [£75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S.	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17189 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-I data for classifying native plant communities - Central Colorado p0004 A75-17176 FRAZEE, C. J.
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAM, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques p0054 A75-17132	p0019 N75-11653 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-I data for classifying native plant communities - Central Colorado p0004 A75-17176 FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY. V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques p0054 A75-17132	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17189 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. 0II-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-I data for classifying native plant communities - Central Colorado p0004 A75-17176 FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAM, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques p0054 A75-17132	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [£75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models p0067 A75-17148 The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 EVANS, M. 8. Investigation of microwave hologram techniques for application to earth resources p0067 A75-17195	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-1 data for classifying native plant communities - Central Colorado p0004 A75-17176 FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135 FREDEN, S. C. Third Earth Resources Technology Satellite Symposium.
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, J. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques p0054 A75-17132 E EAGLEMAN, J. R. Moisture detection from Skylab p0002 A75-17145 Detection of moisture and moisture related phenomena from Skylab	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 EVANS, M. B. Investigation of microwave hologram techniques for application to earth resources p0067 A75-17195 EVERETT, J. R.	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-I data for classifying native plant communities - Central Colorado p0004 A75-17176 FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135 FREDEN, S. C. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY. V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESR0-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques E EAGLEMAN, J. R. Moisture detection from Skylab p0002 A75-17145 Detection of moisture and moisture related phenomens	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [£75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models p0067 A75-17148 The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 EVANS, M. 8. Investigation of microwave hologram techniques for application to earth resources p0067 A75-17195	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-1 data for classifying native plant communities - Central Colorado p0004 A75-17176 FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135 FREDEN, S. C. Third Earth Resources Technology Satellite Symposium.
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques p0054 A75-17132 E EAGLEMAN, J. R. Moisture detection from Skylab p0002 A75-17145 Detection of moisture and moisture related phenomena from Skylab [E75-10035] p0007 N75-12405 Detection of moisture and moisture related phenomena from Skylab [E75-10035] p0007 N75-12405 Detection of moisture and moisture related phenomena from Skylab [E75-10035] p0007 N75-12405	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] STES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural and urban terrain analysis P0055 A75-17198 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] P0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis EVANISKO, F. E. Investigation of microwave hologram techniques for application to earth resources EVERETT, J. R. An evaluation of the suitability of ERTS data for the purposes of petroleum exploration	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. 0il-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-I data for classifying native plant communities - Central Colorado p0004 A75-17176 FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135 FREDEN, S. C. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-SP-356] p0077 N75-10549 FRIEDMAN, D. B. Third Earth Resources Technology Satellite Symposium.
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY. V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESR0-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques E EAGLEMAN, J. R. Moisture detection from Skylab p0002 A75-17145 Detection of moisture and moisture related phenomena from Skylab [E75-10035] p0007 N75-12405 Detection of moisture and moisture related phenomena from Skylab [E75-10007] p0007 N75-13333	PO019 N75-11553 ESPARZA, F. Planning applications in east central Florida [875-10015] ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural and urban terrain analysis ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-755838] FETKIN. V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis EVANS, M. 8. Investigation of microwave hologram techniques for application to earth resources P0067 A75-17195 EVERETT, J. R. An evaluation of the suitability of ERTS data for the	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-I data for classifying native plant communities - Central Colorado p0004 A75-17176 FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135 FREDEN, S. C. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-SP-356] p0077 N75-10549 FRIEDMAN, D. B.
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAM, I. E. Feasibility study ASCS remote sensing/compliance determination system (NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques p0054 A75-17132 E EAGLEMAN, J. R. Moisture detection from Skylab p0002 A75-17145 Detection of moisture and moisture related phenomena from Skylab [E75-10035] p0007 N75-12405 Detection of moisture and moisture related phenomena from Skylab [E75-10007] p0007 N75-13333 Detection of moisture and moisture related phenomena from Skylab [E75-10007] p0007 N75-13333 Detection of moisture and moisture related phenomena from Skylab [E75-10007] p0007 N75-13333	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 EVANS, M. 8. Investigation of microwave hologram techniques for application to earth resources p0067 A75-17195 EVERETT, J. R. An evaluation of the suitability of ERTS data for the purposes of petroleum exploration	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] FOURNIER, G. Oil-on-Water sensor [AD-784104] FRANCIS, R. E. ERTS-I data for classifying native plant communities - Central Colorado D004 A75-17176 FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota P0055 A75-17135 FREDEN, S. C. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY. V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques p0054 A75-17132 E EAGLEMAN, J. R. Moisture detection from Skylab Detection of moisture and moisture related phenomena from Skylab [E75-10035] p0007 N75-12405 Detection of moisture and moisture related phenomena from Skylab [E75-10007] p0007 N75-13333 Detection of moisture and moisture related phenomena from Skylab [E75-10073] p0008 N75-14192	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models p0067 A75-17148 The usefulness of imaging passive microwave for rural p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 EVANS, M. B. Investigation of microwave hologram techniques for application to earth resources p0067 A75-17195 EVERETT, J. R. An evaluation of the suitability of ERTS data for the purposes of petroleum exploration F FAKUNDINY, R. H.	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-1 data for classifying native plant communities - Central Colorado p0004 A75-17176 FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135 FREDEN, S. C. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-5P-356] p0077 N75-10549 FRIEDMAN, D. B. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-5P-356] p0077 N75-10549 FRIEDMAN, D. B. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-5P-356] p0077 N75-10549 FRIEDMAN, D. B. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-5P-356] p0077 N75-10549 FRIORIKSSON, S. Environmental studies of Iceland with ERTS-1 imagery
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY. V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques p0054 A75-17132 E EAGLEMAN, J. R. Moisture detection from Skylab p0002 A75-17145 Detection of moisture and moisture related phenomena from Skylab [E75-10035] p0007 N75-12333 Detection of moisture and moisture related phenomena from Skylab [E75-10007] p0007 N75-13333 Detection of moisture and moisture related phenomena from Skylab [E75-10073] p0008 N75-14192 EBBERT, T. F. The application of Skylab EREP data for land use	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models p0067 A75-17148 The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 EVANS, M. 8. Investigation of microwave hologram techniques for application to earth resources p0067 A75-17195 EVERETT, J. R. An evaluation of the suitability of ERTS data for the purposes of petroleum exploration F FAKUNDINY, R. H. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-1 data for classifying native plant communities - Central Colorado p0004 A75-17176 FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135 FREDEN, S. C. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-5P-356] p0077 N75-10549 FRIEDMAN, D. B. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-5P-356] p0077 N75-10549 FRIEDMAN, D. B. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-5P-356] p0077 N75-10549 FRIORIKSSON, S. Environmental studies of Iceland with ERTS-1 imagery
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques p0054 A75-17132 E EAGLEMAN, J. R. Moisture detection from Skylab p0002 A75-17145 Detection of moisture and moisture related phenomena from Skylab [E75-10035] p0007 N75-12405 Detection of moisture and moisture related phenomena from Skylab [E75-10007] p0007 N75-13333 Detection of moisture and moisture related phenomena from Skylab [E75-10073] p0008 N75-14192 EBBERT, T. F. The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17198 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN. V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 EVANS, M. 8. Investigation of microwave hologram techniques for application to earth resources p0067 A75-17195 EVRETT, J. R. An evaluation of the suitability of p0035 N75-10553 F FAKUNDINY, R. H. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10558	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-I data for classifying native plant communities - Central Colorado p0004 A75-17176 FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135 FREDEN, S. C. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-SP-356] p0077 N75-10549 FRIEDMAN, D. B. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-SP-356] p0077 N75-10549 FRIEDMAN, D. B. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-SP-356] p0077 N75-10549 FRIEDRIKSSON, S. Environmental studies of Iceland with ERTS-1 imagery p0012 A75-17105
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques p0054 A75-17132 E EAGLEMAN, J. R. Moisture detection from Skylab p0002 A75-17145 Detection of moisture and moisture related phenomena from Skylab [E75-10035] p0007 N75-12405 Detection of moisture and moisture related phenomena from Skylab [E75-10007] p0007 N75-13333 Detection of moisture and moisture related phenomena from Skylab [E75-10073] p0008 N75-14192 EBBERT, T. F. The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus agreas	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models p0067 A75-17198 The usefulness of imaging passive microwave for rural p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 EVANS, M. 8. Investigation of microwave hologram techniques for application to earth resources p0067 A75-17195 EVERETT, J. R. An evaluation of the suitability of ERTS data for the purposes of petroleum exploration p0035 N75-10553 F FAKUNDINY, R. H. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FALLER, K. Preliminary results of fisheries investigation associated	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-1 data for classifying native plant communities contral Colorado p0004 A75-17176 FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135 FREDEN, S. C. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-5P-356] p0077 N75-10549 FRIEDMAN, D. B. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-5P-356] p0077 N75-10549 FRIEDMAN, D. B. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-5P-356] p0077 N75-10549 FRIEDRIKSSON, S. Environmental studies of Iceland with ERTS-1 imagery p0012 A75-17105 FROST, R. T. Remote detection of soil surface moisture p0003 A75-17146
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, J. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques p0054 A75-17132 E EAGLEMAN, J. R. Moisture detection from Skylab p0002 A75-17145 Detection of moisture and moisture related phenomena from Skylab [E75-10035] p0007 N75-12405 Detection of moisture and moisture related phenomena from Skylab [E75-10073] p0007 N75-13333 Detection of moisture and moisture related phenomena from Skylab [E75-10073] p0008 N75-14192 EBBERT, T. F. The application of Skylab EREP data for land use-classification and mapping in the Cleveland and Columbus areas [E75-10083] p0023 N75-14201	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 EVANS, M. B. Investigation of microwave hologram techniques for application to earth resources p0067 A75-17195 EVERETT, J. R. An evaluation of the suitability of ERTS data for the purposes of petroleum exploration p0035 N75-10553 FAKUNDINY, R. H. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] FALLER, K. Preliminary results of fisheries investigation associated with Skylab-3 p0003 A75-17167	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological enalysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-I data for classifying native plant communities - Central Colorado p0004 A75-17176 FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135 FREDEN, S. C. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-SP-356] p0077 N75-10549 FRIEDMAN, D. B. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-SP-356] p0077 N75-10549 FRIORIKSSON, S. Environmental studies of Iceland with ERTS-1 imagery p0012 A75-17105 FROST, R. T. Remote detection of soil surface moisture p0003 A75-17146
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY. V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques p0054 A75-17132 E EAGLEMAN, J. R. Moisture detection from Skylab p0002 A75-17145 Detection of moisture and moisture related phenomena from Skylab [E78-10035] p0007 N75-13333 Detection of moisture and moisture related phenomena from Skylab [E75-10007] p0007 N75-13333 Detection of moisture and moisture related phenomena from Skylab [E75-10007] p0008 N75-14192 EBBERT, T. F. The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus areas [E75-10083] p0023 N75-14201 EDWARDS, D. Applications of remote sensing by the state of	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17198 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 EVANS, M. 8. Investigation of microwave hologram techniques for application to earth resources p0067 A75-17195 EVENETT, J. R. An evaluation of the suitability of ERTS data for the purposes of petroleum exploration p0035 N75-10553 F FAKUNDINY, R. H. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FALLER, K. Preliminary results of fisheries investigation associated with Skylab-3 Po0003 A75-17167 FANTASIA, J. F. Development of an experimental airborne laser remote	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-1 data for classifying native plant communities central Colorado p0004 A75-17176 FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135 FREDEN, S. C. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-SP-356] p0077 N75-10549 FRIEDMAN, D. B. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-SP-356] p0077 N75-10549 FRIORIKSSON, S. Environmental studies of Iceland with ERTS-1 imagery p0012 A75-17105 FROST, R. T. Remote detection of soil surface moisture p0003 A75-17146 FUBARA, D. M. J. Interaction of marine geodesy, satellite technology and ocean physics
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, J. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques p0054 A75-17132 E EAGLEMAN, J. R. Moisture detection from Skylab p0002 A75-17145 Detection of moisture and moisture related phenomena from Skylab [E75-10035] p0007 N75-12405 Detection of moisture and moisture related phenomena from Skylab [E75-10073] p0007 N75-13333 Detection of moisture and moisture related phenomena from Skylab [E75-10073] p0008 N75-14192 EBBERT, T. F. The application of Skylab EREP data for land use-classification and mapping in the Cleveland and Columbus areas [E75-10083] p0023 N75-14201	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models p0067 A75-17148 The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 EVANS, M. 8. Investigation of microwave hologram techniques for application to earth resources p0067 A75-17195 EVERETT, J. R. An evaluation of the suitability of ERTS data for the purposes of petroleum exploration F FAKUNDINY, R. H. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FALLER, K. Preliminary results of fisheries investigation associated with Skylab-3 FANTASIA, J. F. Development of an experimental airborne faser remote sensing system for the detection and classification of oil	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological enalysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-I data for classifying native plant communities - Central Colorado FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135 FREDEN, S. C. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-SP-356] p0077 N75-10549 FRIEDMAN, D. B. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-SP-356] p0077 N75-10549 FRIORIKSSON, S. Environmental studies of Iceland with ERTS-1 imagery p0012 A75-17105 FROST, R. T. Remote detection of soil surface moisture p0003 A75-17146 FUBARA, D. M. J. Interaction of marine geodesy, satellite technology and ocean physics [NASA-CR-137470] p0043 N75-10671
[278-10032] p0007 N75-12402 DUBENSKIY. V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, I. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques p0054 A75-17132 E EAGLEMAN, J. R. Moisture detection from Skylab p0002 A75-17145 Detection of moisture and moisture related phenomena from Skylab [275-10035] p0007 N75-12405 Detection of moisture and moisture related phenomena from Skylab [275-10007] p0007 N75-13333 Detection of moisture and moisture related phenomena from Skylab [275-10073] p0008 N75-14192 EBBERT, T. F. The application of Skylab EREP data for land use-classification and mapping in the Cleveland and Columbus areas [275-10083] p0023 N75-14201 EDWARDS, D. Applications of remote sensing by the state of Nebraska EGAN, W. G. Boundaries of ERTS and aircraft data within which useful	ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 EVANS, M. B. Investigation of microwave hologram techniques for application to earth resources p0067 A75-17195 EVERETT, J. R. An evaluation of the suitability of ERTS data for the purposes of petroleum exploration p0035 N75-10553 F FAKUNDINY, R. H. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-1090] p0034 N75-10536 FALLER, K. Preliminary results of fisheries investigation associated with Skylab-3 p0003 A75-17167 FANTASIA, J. F. Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-I data for classifying native plant communities - Central Colorado p0004 A75-17176 FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135 FREDEN, S. C. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-SP-356] p0077 N75-10549 FRIEDMAN, D. B. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-SP-356] p0077 N75-10549 FRIORIKSSON, S. Environmental studies of Iceland with ERTS-1 imagery p0012 A75-17105 FROST, R. T. Remote detection of soil surface moisture p0003 A75-17146 FUBARA, D. M. J. Interaction of marine geodesy, satellite technology and ocean physics [NASA-CR-137470] p0043 N75-10671
forest stress [E78-10032] p0007 N75-12402 DUBENSKIY, V. P. System for electronic transformation and geographic correlation of satellite television information p0071 N75-11434 DUC, N. T. European earth-resources satellite experiments [ESRO-SP-100] p0081 N75-14216 DUGGAN, J. E. Feasibility study ASCS remote sensing/compliance determination system [NASA-CR-134288] p0007 N75-12419 DUGGIN, M. J. On the natural limitations of target differentiation by means of spectral discrimination techniques p0054 A75-17132 E EAGLEMAN, J. R. Moisture detection from Skylab p0002 A75-17145 Detection of moisture and moisture related phenomena from Skylab [E75-10035] p0007 N75-12405 Detection of moisture and moisture related phenomena from Skylab [E75-10007] p0007 N75-1333 Detection of moisture and moisture related phenomena from Skylab [E75-10073] p0008 N75-14192 EBBERT, T. F. The application of Skylab EREP data for land use-classification and mapping in the Cleveland and Columbus areas [E75-10083] p0023 N75-14201 EDWARDS, D. Applications of remote sensing by the state of Nebraska p0069 A75-17232	p0019 N75-11553 ESPARZA, F. Planning applications in east central Florida [E75-10015] p0021 N75-13337 ESTES, J. E. Operational use of satellite and high altitude remote sensing for the generation of input data for water demand models The usefulness of imaging passive microwave for rural p0055 A75-17199 ESTES, J. M. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 ETKIN, V. S. Microwave emission characteristics of moist soils p0001 A75-15180 EVANISKO, F. E. The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 EVANS, M. 8. Investigation of microwave hologram techniques for application to earth resources p0067 A75-17195 EVERETT, J. R. An evaluation of the suitability of p0067 A75-10553 F FAKUNDINY, R. H. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10558 FALLER, K. Preliminary results of fisheries investigation associated with Skylab-3 p0003 A75-17167 FANTASIA, J. F. Development of an experimental airborne laser remote sensing system for the detection and classification of oil p0068 A75-17167	FORMISANO, V. Solar wind and geomagnetic activity p0027 A75-11975 FORSTER, S. W. Assessment of ERTS-1 imagery as a tool for regional geological enalysis in New York State [E74-10809] p0034 N75-10536 FOSTER, K. E. A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] p0006 N75-10544 FOURNIER, G. Oil-on-Water sensor [AD-784104] p0072 N75-11454 FRANCIS, R. E. ERTS-I data for classifying native plant communities - Central Colorado FRAZEE, C. J. Use of ERTS-1 imagery for land evaluation in Pennington County, South Dakota p0055 A75-17135 FREDEN, S. C. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-SP-356] p0077 N75-10549 FRIEDMAN, D. B. Third Earth Resources Technology Satellite Symposium. Volume 2: Summary of results [NASA-SP-356] p0077 N75-10549 FRIORIKSSON, S. Environmental studies of Iceland with ERTS-1 imagery p0012 A75-17105 FROST, R. T. Remote detection of soil surface moisture p0003 A75-17146 FUBARA, D. M. J. Interaction of marine geodesy, satellite technology and ocean physics [NASA-CR-137470] p0043 N75-10671

purposes [E75-10103]

FAWKES, G.

The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 8: Atmosphere
[NASA-CR-141191] p0080 N75-14213

pi0025 N75-15121 FULDE. J.
On the detectability of atmospheric carbon monoxide by microwave remote sensing p0013 A75-17129

Investigation of radar discrimination of sea ice p0042 A75-17121

EGBERT, D.

Optical data processing analysis of stream patterns exhibited on ERTS-1 imagery p0056 A75-17219

G

GALLAGHER, D. B.

Urban and regional land use analysis: CARETS and census cities experiment package p0024 N75-15109 [F75-10091]

GALLERY, W. O.

Remote sounding of stratospheric temperatures using high resolution radiance measurements from the IRIS-D [NASA-CR-141130] p0022 N75-13452

GALLI DE PARATESI, S.
Thermal behaviour of some rice fields affected by yellows-type disease p0004 A75-17176 p0004 A75-17174

GALLIDEPARATESI, S.

Agricultural resources investigations in northern Ital southern France and Madagascar p0009 N75-14252 GALLOWAY, I. R.

The interpretation and use of false-colour infra-red and true colour photography of part of Argentina obtained by p0067 A75-17192 Skylark earth resources rockets

The application of pattern classification techniques to Skylark earth resources rocket imagery

p0056 A75-19268

GALPER, A. M.

Detection of synchronous changes in fluxes of energetic Detection of synchronous energieselectrons and gamma-quanta during a magnetic p0064 A75-12317 GARRETT, G. B.

Multidisciplinary applications of ERTS and Skylab data p0015 A75-17234 in Ohio

GAUTREAUX, M. R.

Land use classification and ground truth correlations from imultaneously acquired aircraft and ERTS-1 MSS data p0014 A75-17188

GAZLEY, C., JR.

Analysis of pseudocolor transformations of ERTS-1 images of Southern California area p0036 N75-13346

GEBHART, R. F.

Satellites and computer in the service of meteorology: WWW - The global weather observation syst

p0053 A75-11274 GEGG, S. R.

Cruise data report. CHAIN 115 leg 2, Southlant

GERTSENOVA K. N.

Investigation of accuracy of determination of plane coordinates by means of the Central Scientific Research Institute of Geodesy, aerial photography and cartography airborne rangefinder p0061 N75-14255

[AD-785886]

Nimbus 4 observations of changes in total ozone and stratospheric temperatures during a sudden war

CHOSH A K

A microwave correlation spectrometer for air pollutio p0063 A75-11732 monitorina GILL I L

p0012 A75-14841

p0061 N75-14255

Investigation of accuracy of determination of plane coordinates by means of the Central Scientific Research Institute of Geodesy, aerial photography and cartography

airborne rangefinder [AD-785886]

GILLE, J. C. Measurement of trace stratospheric gases by inverting

thermal infrared limb radiance profiles p0018 N75-11523

Methods of calculating infrared transfer:

p0019 N75-11560

GILMER, D. S. Utilization of Skylab (EREP) system for appraising changes in continental migratory bird habitat

[E74-10817] 74-10817] p0006 N75-10543
Utility of ERTS for monitoring the breeding habit of igratory waterfowl p0006 N75-10557 migratory waterfowl Utilization of Skylab (EREP) system for appraising hanges in continental migratory bird habitat

fE75-10061 75-10061] p0008 N75-13361 Utilization of Skylab (EREP) system for appraising

changes in continental migratory bird habitat [E75-10087] p0009 N75-15105

GIRARD-GANNEAU, C. M. Photographs from balloons - Their use in agronomy management of environment p0055 A75-17 p0055 A75-17191

GIRARD, M. C. Photographs from balloons - Their p0055 A75-17191 management of environment

GLIDDEN, W. S. Computer derived coastal water classifications

p0042 A75-17177 spectral signatures

p0066 A75-16191 A radio picture of the earth GLORIA, H. R.

Airborne survey of major air basins in

p0011 A75-14360

GLUMOV, I. F Analysis of methods for the geodetic support of complex

p0031 N75-13498 GLUMOV, V. P

Analysis of methods for the geodetic support of complex sea geophysical surveys [JPRS-63821] p0031 N75-13498

GOETTELMAN, R.

Use of visible, near-infrared, and thermal infrared remote sensing to study soil moisture GOETZ, A. F. H. p0002 A75-17144

Preliminary results in the comparison of Skylab ERTS and R8-57 images for the detection of lineaments ar fractures in Precambrian, Paleozoic, and late Tertiary rocks on and near the Colorado Plateau, North-Central Arizona [E75-10060] p0037 N75-13360

GOLDMAN, G. C.

Investigation of transient earth resources phenomena: Continuation study

[NASA-CR-139163] p0073 N75-15131 GOLDSTEIN, W.

Investigation of land use of northern megalopolis using ERTS-1 imagery

p0017 N75-10545

GOLENETSKII, S. V.
Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667 GOLUBEV, A. N.

Optical quantum generators in geodetic measure p0032 N75-15237 [AD-787281]

Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243

GOMENIUK, A. S. Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases

using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 GONIN, G. B.

Geological and geophysical studies from space and rospects for their development p0035 N75-11436 GOOSEN, D.

Preliminary results of the interpretation of ERTS-1 imagery for soil surveys in the Merida region, Spain p0009 N75-14247

GOPALAPILLAI, S.

Calibration and evaluation of Skylab altimetry for geodetic determination of the geoid p0030 N75-13364

GORDON, H. H. Remote sensing applications in marine science programs at VIMS

[NASA-CR-137475] p0043 N75-10566

GORDON, K. S.

A motion-compensated spatial scanner

p0068 A75-17213 GORDON, R. C.

Usefulness of Skylab color photography and ERTS-1 multispectral imagery for mapping range vegetation types

in southwestern Wyoming [E75-10097] p0010 N75-15115

GÔVOROV, V. M. Geometric distortions of opticommercial panoramic p0057 N75-11425 television systems

GOWER, J. F. R. Measurement of sea surface currents using airborne

Measurements of meteorological parameters and air pollution spread over Stockholm, Sweden, by means of remote sensing and airborne instrumentation

n0018 N75-11531 GRANDFIELD, A. L Summary of flight performance of Resources Experiment Package /EREP/ of the Skylab Earth

p0076 A75-17211 GRANT, K

The PUCE programme for terrain name ring purposes. 2: Procedu Procedures for terrain engineering purposes. [PB-233847/3] p0029 N75-10625

GRAVES, R. C.

Skylab environmental and crew microbiology studies p0024 N75-14366

GREELEY, R.

Use of visible, near-infrared, and thermal infrared ren p0002 A75-17144 sensing to study soil moisture GREEN, T., III

Scanning thermal plumes n0014 A75-17162

GREENBLAT, E. J.

Evaluation of ERTS data utilization in develo p0055 A75-17193 countries

GRIFFITH, C. G.

Rainfall estimations from geosynchronous satellite p0047 A75-13596

GRIGGS, M.

rmination of the aerosol content from ERTS-1 data p0013 A75-17130 Global measurements of air pollution from satellites p0018 N75-11521

GRIGORYEV, A. A.

Experience in integrated geological and geographical interpretation of space photographs obtained in the USSR p0036 N75-11437

Dust storms according to data of space research INASA-TT-F-16021 p0021 N75-12463

GROMAN, R. C. Cruise data report. CHAIN 115 leg 2, Southlant

[AD-784421] DO029 N75-10641 GROSS, H. G.

Crude and refined petroleum oil structured luminescence signatures induced by UV laser or lamp and their remote sensing applications p0033 A75-17207

The value of improved (ERS) information based on domestic distribution effects of U.S. agriculture crops
[NASA-CR-141046] p0007 N75-12423

GROTEN, W.

Use of terrestrial gravity data in combination with satellite results for determining geoid heights in Central Europ p0029 A75-18128

GROVES, T. O.

Skylab environmental and crew microbiology studies p0024 N75-14366 GUENTHER, R. ERTS-1 data on Afar tectonics p0031 N75-14239

GUIGNARD, J. P.
A preliminary feasibility study on a synthetic aperture radar / SARSAT/ for earth resources surveys p0067 A75-17194

GUNLIA G S

Methods for the automatic recording of atmospheric dust concentrations and some of the results obtain p0057 N75-11502

GURIAN, IU. A. a Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667 GURK, H. M.

RK, H. M.
More information and less data from space
p0063 A75-11730

GURNETT, D. A The earth as a radio source - Terrestrial kilometric radiation p0011 A75-11211

GURVICH, A. S. The radio emission of the earth as a planet

DO076 A75-15319

Recent and present sedimentation complexity of the Rhone delta and the French mediterranean coast (Golion project) p0045 N75-14228

GUYADER, M. Agricultural resources investigations in northern Italy, southern France and Madagascar p0009 N75-14252

Н

HAAGAARD, A.

HALLIDAY, R. A.

Measurements of meteorological parameters and air pollution spread over Stockholm, Sweden, by means of remote sensing and airborne instrumentation

p0018 N75-11531 HAASE, E. F.

A study to explore the use of orbital remote sensing to determine native arid plant distribution

p0006 N75-10544 [E74-10818]

HAEFNER, H Methodological aspects and regional examples of snow-cover mapping from ERTS-1 and EREP imagery p0051 N75-14232

HAENEL G. Scattering and absorption from poly-dispersed aerosols

HAHN, W. S.

Pacific range electromagnetic signature studies / Project RESS/ p0042 A75-17479 PRESS/ HALE, L C

Ice particulates in the mesosphere p0012 A75-16752 HALL F.G.

First results from the crop identification technology assessment for remote sensing /CITARS/ p0004 A75-17175

HALL, G. D.

Global measurements of air pollution from satellites p0018 N75-11521

Water survey of Canada: Application for use of ERTS-A for retransmission of water resources data [E74-10805] 74-10805] p0049 N75-11409 Water survey of Canada: Application for use of ERTS-A

for retransmission of water resources data p0050 N75-13342 [E75-10026]

Water survey of Canada: Application for use of ERTS-A for retransmission of water resources data p0050 N75-13353

HANNAH, J. W.

Planning applications in east central Florida [E75-10015] p0021 p0021 N75-13337 HANSON, K. The detection by ERTS-1 of wind-induced ocean surface

features in the lee of the Antilles Islands

p0041 A75-10644 HARALICK, R. H.

Use of feature extraction techniques for the texture and context information in ERTS imagery: Spectral and textural

processing of ERTS imagery [E75-10011] p0059 N75-13335

HARDY, E. Evaluation of Skylab imagery as an information service for investigating land use and natural resources [E74-10730] p0056 N75-10526

Design concepts for land use and natural resource inventories and information systems p0013 A75-17133

B-5

Evaluation of Skylab imagery as an information service	HENRIKSON, P. J.	HOOPER, A. J.
for investigating land use and natural resources	Correlation and registration of ERTS multispectral	Ground-truth measurements in relation to aircraft and
[E75-10018] p0020 N75-12397 Evaluation of Skylab imagery as an information service	imagery [NASA-CR-134294] p0058 N75-12420	satellite studies of agricultural land use and land classification in Britain p0009 N75-14251
for investigating land use and natural resources	HENSLEY, W. E.	HOOPER, J. O.
[E75-10013] p0059 N75-13336 Evaluation of Skylab imagery as an information service	Summary of flight performance of the Skylab Earth Resources Experiment Package /EREP/	The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199
for investigating land use and natural resources	p0076 A75-17211	Microwave radiometric characteristics of snow-covered
[E75-10068] p0059 N75-14187 Evaluation of Skylab imagery as an information service	HERBERT, G. A.	terrain p0049 A75-17200 HOPPIN, R. A.
for investigating land use and natural resources	Geophysical monitoring for climatic change. The NOAA	Experiment to evaluate feasibility of utilizing Skylab-EREP
[NASA-CR-141372] p0024 N75-15100 HARDY, J. R.	program p0018 N75-11510 Ground-based measurements of solar radiation by	remote sensing data of tectonic analysis of the Bighorn Mountains region, Wyoming-Montana
The interpretation and use of false-colour infra-red and	geophysical monitoring for climatic change (USA)	[E75-10014] p0030 N75-12395
true colour photography of part of Argentina obtained by Skylark earth resources rockets p0067 A75-17192	p0018 N75-11527	HORVATH, R. Oil pollution detection, monitoring and law
A comparison of the capabilities of ERTS-1 and the	HERD, L. O. Multidisciplinary applications of ERTS and Skylab data	enforcement
Skylark earth-resources rocket for resources surveying in central Argentina p0023 N75-14250	in Ohio p0015 A75-17234	[E75-10027] p0022 N75-13343 HOSFORD, K. R.
HARDY, N.	HERGENRADER, G. Applications of remote sensing by the state of	Resource analysis applications in Michigan
Detection of moisture and moisture related phenomena from Skylab	Nebraska p0069 A75-17232	p0069 A75-17231 HOUSE, F. B.
[E75-10035] p0007 N75-12405	HESKETH, W. D.	Measurement of trace stratospheric gases by inverting
HARRIS, J. C. Application of remote sensing to state and regional	Global measurements of air pollution from satellites p0018 N75-11521	thermal infrared limb radiance profiles p0018 N75-11523
problems	HESSIN, T. D.	HOYT, D. V.
[NASA-CR-140805] p0078 N75-11414 HART, J. L.	Prediction of the Fraunhofer line detectivity of luminescent materials p0068 A75-17221	Geophysical monitoring for climatic change. The NQAA program p0018 N75-11510
Extensive summer upwelling on Lake Michigan during	HIBBEN, S. G.	program p0018 N75-11510 Ground-based measurements of solar radiation by
1973 observed by NOAA-2 and ERTS-1 satellites p0048 A75-17160	Soviet satellite geodesy: Basis and status of Soviet laser geodesy and long-base line interferometry	geophysical monitoring for climatic change (USA)
HART, W. G.	[AD-781138/3] p0029 N75-10524	p0018 N75-11527
A study of the early detection of insect infestations and density/distribution of host plants	HIGER, A. L.	HRUBESH, L. W. Feasibility study of a Gunn-diode microwave cavity
[E75-10062] p0008 N75-13362	Water-management models in Florida from ERTS-1 data p0047 A75-13579	spectrometer as a formaldehyde monitor
HASELL, P. G., JR. The remote identification of terrain features and materials	Surface water modeling Everglades Water Basin,	[UCID-16488] p0071 N75-11311 Microwave rotational spectroscopy: A physical technique
at a California test site: An investigative study of	Florida [E74-10814] p0049 N75-10540	for specific pollutant monitoring p0018 N75-11529
techniques [PB-235991/7] p0031 N75-14261	HILL, I. E.	HSU, SY. Automatic pattern recognition and mapping utilizing
The remote identification of terrain features and materials	Generalisation of ERTS data for global-scale investigations p0081 N75-14224	thermal infrared imagery p0070 A75-18074
at Pennsylvania test sites: An investigative study of	HILWIG, F. W.	HUBBLE, L Image selection system
techniques [PB-235992/5] p0031 N75-14263	Preliminary results of the interpretation of ERTS-1	[NASA-CR-2475] p0058 N75-12422
HATCHER, N. M.	imagery for soil surveys in the Merida region, Spain p0009 N75-14247	HUFF, D. D. Application of remote sensing to the location of
Simultaneous active and passive microwave response of the earth - The Skylab radscat experiment	HIRSCHFELD, T.	hydrologically active /source/ areas p0066 A75-17141
p0012 A75-17114	The remote Raman spectrometer is a viable instrument for remote sensing of the environment	HUFFAKER, R. M. Application of a single laser Doppler system to the
Summary of flight performance of the Skylab Earth Resources Experiment Package /EREP/	p0068 A75-17214	measurement of atmospheric winds
p0076 A75-17211	HO, W. W. Measurements of the dielectric properties of sea water	[NASA-TM-X-64891] p0021 N75-12462 HULKE, S. D.
HAWTHORNE, C. A. The usefulness of imaging passive microwave for rural	at 1.43 GHz	A hydrogeomorphic approach to evaluating flood potential
and urban terrain analysis poops A75-17199	[NASA-CR-2458] p0045 N75-13496	in central Texas from orbital and suborbital remote sensing imagery p0047 A75-17140
and urban terrain analysis p0055 A75-17199 HAY, C. M.	HODAPP, C. J. Skylab environmental and crew microbiology studies	imagery p0047 A75-17140 HUNT, G. R.
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366	imagery p0047 A75-17140
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R.	HODAPP, C. J. Skylab environmental and crew microbiology studies	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and near-infrared p0033 A?5-17220 HURD, D.
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN. R. Evaluation of ERTS-1 data by analogue and digital	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and near-infrared p0033 A75-17220
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN. R. Evaluation of ERTS-1 data by analogue and digital techniques p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and near-infrared HURD, D. Image selection system
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN. R. Evaluation of ERTS-1 data by analogue and digital techniques p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and near-infrared HURD, D. Image selection system
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN. R. Evaluation of ERTS-1 data by analogue and digital techniques p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422
and urban terrain analysis p0055 A75-17199 HAY, C. M. A gricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital techniques p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803]	imagery HUNT, G. R. Remote sensing of rock type in the visible and p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] Image selection system P0058 N75-12422
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data	imagery HUNT, G. R. Remote sensing of rock type in the visible and p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422 IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion
and urban terrain analysis p0055 A75-17199 HAY, C. M. A gricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital p0059 N75-14218 Geologic evaluation of central Harly from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] p0057 N75-10531 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques	imagery HUNT, G. R. Remote sensing of rock type in the visible and near-infrared HURD, D. Image selection system [NASA-CR-2475] P0058 N75-12422
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141288] p0079 N75-14203	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite date for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] p0078 N75-12390	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and p0033 A75-17220 HUND, D. Image selection system [NASA-CR-2475] p0058 N75-12422 IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital techniques p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14204 FRTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains. using automatic data processing techniques [E75-10002]	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J.	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] p0057 N75-10531 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] p0078 N75-12390 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and near-infrared hURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422 IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: AERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration p0075 A75-13592	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types in the Colorado Mountains and the selected cover types and the	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422 IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration p0075 A75-13592	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] p0057 N75-10531 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] p0078 N75-12390 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] p0037 N75-14191 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using sutomatic data processing techniques	imagery HUNT, G. R. Remote sensing of rock type in the visible and near-infrared HURD, D. Image selection system [NASA-CR-2475] IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667 INGRAM, J. D. Microbial abundance and thermoluminescence of
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: AERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration p0075 A75-13592	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite date for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104]	imagery HUNT, G. R. Remote sensing of rock type in the visible and p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration p0075 A75-13592 HEINTZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0019 N75-11553	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] p0057 N75-10531 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] p0078 N75-12390 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] p0037 N75-14191 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104] p0073 N75-15122	imagery HUNT, G. R. Remote sensing of rock type in the visible and near-infrared HURD, D. Image selection system [NASA-CR-2475] IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667 INGHAM, J. D. Microbial abundance and thermoluminescence of antarctic dry valley soils INGLE, S. J. A study of the early detection of insect infestations and
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration p0075 A75-13592 HEINTZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0019 N75-11553 HEISS, K. P. The economic value of remote sensing of earth resources	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite date for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104]	imagery HUNT, G. R. Remote sensing of rock type in the visible and near-infrared HURD, D. Image selection system [NASA-CR-2475] IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV INGHAM, J. D. Microbiał abundance antarctic dry valley soils INGLE, S. J. A study of the early detection of insect infestations and density/distribution of host plants [E75-10062]
and urban terrain analysis p0055 A75-17199 HAY, C. M. A gricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration p0075 A75-13592 HEINTZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0019 N75-11553 HEISS, K. P. The economic value of remote sensing of earth resources from space. An ERTS overview and the value of continuity of service. Volume 1: Summary	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10104] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104] HOFFMAN, R. Applications of remote sensing by the state of Nobreska p0069 A75-17232	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422 IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII. V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667 INGHAM, J. D. Microbial abundance antercite dry valley soils p0005 A75-17973 INGLE, S. J. A study of the early detection of insect infestations and density/distribution of host plants [E75-10062] p0008 N75-13362
and urban terrain analysis p0055 A75-17199 HAY, C. M. A gricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital techniques p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14204 HECKER, E. J. NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration HEINZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0019 N75-11553 HEISS, K. P. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service.	Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104] P0073 N75-15122 HOFFMAN, R. Applications of remote sensing by the state of Nebraska	imagery HUNT, G. R. Remote sensing of rock type in the visible and near-infrared HURD, D. Image selection system [NASA-CR-2475] IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667 INGHAM, J. D. Microbial abundance anterctic dry valley soils INGLE, S. J. A study of the early detection of insect infestations and density/distribution of host plants [E75-10062] INGRAO, H. C. Development of an experimental airborne laser remote sensing system for the detection and classification of oil
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration p0075 A75-13592 HEINTZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0019 N75-111553 HEISB, K. P. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] p0057 N75-10531 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] p0078 N75-12390 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] p0037 N75-14191 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104] p0073 N75-15122 HOFFMAN, R. Applications of remote sensing by the state of Nebraska p0069 A75-17232 HOLLYDAY, E. F. Hydrologic significance of lineaments in central Tennessee [E75-10030]	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422 IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667 INGHAM, J. D. Microbial abundance and thermoluminescence of antarctic dry valley soils INGLE, S. J. A study of the early detection of insect infestations and density/distribution of host plants [E75-10062] p0008 N75-13362 INGRAO, H. C. Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills
and urban terrain analysis p0055 A75-17199 HAY, C. M. A gricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration p0075 A75-13592 HEINTZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0019 N75-11553 HEISS, K. P. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite date for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10014] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104] HOFFMAN, R. Applications of remote sensing by the state of Nebraska policy pol	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422 IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667 INGHAM, J. D. Microbial abundance antercite dry valley soils p0005 A75-17973 INGLE, S. J. A study of the early detection of insect infestations and density/distribution of host plants [£75-10062] p0008 N75-13362 INGRAO, H. C. Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 ISACMSEN, Y. W. Assessment of ERTS-1 imagery as a tool for regional
and urban terrain analysis p0055 A75-17199 HAY, C. M. A gricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital pcosing poof poof poof poof poof poof poof poo	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-100104] P0037 N75-14191 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-100104] P0037 N75-14191 Applications of remote sensing by the state of Nebraska HOLLYOAY, E. F. Hydrologic significance of lineaments in central Tennessee [E75-10030] HOLMES, J. J. Application and sensitivity investigation of Fourier transforms for microwave radiometric inversions	imagery HUNT, G. R. Remote sensing of rock type in the visible and near-infrared HURD, D. Image selection system [NASA-CR-2475] IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV INGHAM, J. D. Microbial abundance and thermoluminescence of p0005 A75-17973 INGLE, S. J. A study of the early detection of insect infestations and density/distribution of host plants [E75-10062] INGRAO, H. C. Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills ISACHSEN, Y. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State
and urban terrain analysis p0055 A75-17199 HAY, C. M. A gricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital techniques p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration p0075 A75-13592 HEINTZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0019 N75-11553 HEISS, K. P. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141193] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HEMPENIUS, S. A. The ITC 8-image comparator and its possible use in ERTS	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-1002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-1002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-1004] P0073 N75-14191 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104] P0073 N75-15122 HOLFMAN, R. Applications of remote sensing by the state of Nebraska p0098 A75-17232 HOLLYDAY, E. F. Hydrologic significance of lineaments in central Tennessee [E75-10030] P0050 N75-12401 HOLMES, J. J. Application and sensitivity investigation of Fourier transforms for microwave radiometric inversions [NASA-CR-140754]	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422 IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667 INGHAM, J. D. Microbial abundance antercite dry valley soils p0005 A75-17973 INGLE, S. J. A study of the early detection of insect infestations and density/distribution of host plants [£75-10062] p0008 N75-13362 INGRAO, H. C. Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0068 A75-17206 ISACMSEN, Y. W. Assessment of ERTS-1 imagery as a tool for regional
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital pcomposed poots p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration p0075 A75-13592 HEINTZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0019 N75-11553 HEISB, K. P. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HEMPENIUS, S. A. The ITC B-image comparator and its possible use in ERTS and EREP investigations p0060 N75-14222	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite date for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104] P0073 N75-14191 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104] P0073 N75-1224 HOFFMAN, R. Applications of remote sensing by the state of Nebraska HOLLYDAY, E. F. Hydrologic significance of lineaments in central Tennessee [E75-10030] HOLMES, J. J. Application and sensitivity investigation of Fourier transforms for microwave radiometric inversions [NASA-CR-140754] P0070 N75-11310 HOLTZMAN, J. C. Simultaneous active and passive microwave response	imagery HUNT, G. R. Remote sensing of rock type in the visible and p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422 IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV INGHAM, J. D. Microbial abundance and thermoluminescence of antarctic dry valley soils INGLE, S. J. A study of the early detection of insect infestations and density/distribution of host plants [E75-10062] p0008 N75-13362 INGRAO, H. C. Development of an experimental airbome laser remote sensing system for the detection and classification of oil spills p008 A75-17206 ISACHSEN, Y. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 ISNAQ, A. M. Application of remote sensing to the location of
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital techniques p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration p0075 A75-13592 HEINTZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0019 N75-11553 HEISS, K. P. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141193] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141193] p0079 N75-14204 HEMPENIUS, S. A. The ICS -image comparator and its possible use in ERTS end EREP investigations p0060 N75-14222 HEMPHILL, W. R. Prediction of the Fraunhofer line detectivity of	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic date processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104] HOFFMAN, R. Applications of remote sensing by the state of Nebraska p0069 A75-17232 HOLLYDAY, E. F. Hydrologic significance of lineaments in central Tennessee [E75-10030] HOLMES, J. J. Application and sensitivity investigation of Fourier transforms for microwave radiometric inversions [MOLTZMAN, J. C.	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and near-infrared p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422 IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667 INGHAM, J. D. Microbial abundance and thermoluminescence of antarctic dry valley soils p0005 A75-17737 INGLE, S. J. A study of the early detection of insect infestations and density/distribution of host plants [E75-10062] INGRAO, H. C. Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0088 A75-17208 ISACHSEN, Y. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] ISHAO, A. M. Application of remote sensing to the location of hydrologically active /source/ areas
and urban terrain analysis p0055 A75-17199 HAY, C. M. A gricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141258] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration p0075 A75-13592 HEINTZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0019 N75-11553 HEISS, K. P. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141288] p0079 N75-14203 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141288] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HEMPENIUS, S. A. The ITC 8-image comparator and its possible use in ERTS and EREP investigations p0060 N75-14222 HEMPHILL, W. R. Prediction of the Fraunhofer line detectivity of public pounds.	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10072] Applications of remote sensing by the state of Nebraska p0069 A75-17232 HOLLYDAY, E. F. Hydrologic significance of lineaments in central Tennessee [E75-10030] PO050 N75-12401 HOLLER, J. J. Application and sensitivity investigation of Fourier transforms for microwave radiometric inversions [NASA-CR-140754] p0070 N75-11310 HOLLER, J. C. Simultaneous active and passive microwave response of the earth - The Skylab radscat experiment	imagery HUNT, G. R. Remote sensing of rock type in the visible and near-infrared HURD, D. Image selection system [NASA-CR-2475] PO058 N75-12422
and urban terrain analysis p0055 A75-17199 HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital techniques p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration p0075 A75-13592 HEINTZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0019 N75-11553 HEISS, K. P. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141193] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141193] p0079 N75-14204 HEMPENIUS, S. A. The ICS -image comparator and its possible use in ERTS end EREP investigations p0060 N75-14222 HEMPHILL, W. R. Prediction of the Fraunhofer line detectivity of	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10072] HOFFMAN, R. Applications of remote sensing by the state of Nebraska p0093 A75-17232 HOLLYDAY, E. F. Hydrologic significance of lineaments in central Tennessee [E75-10030] HOLMES, J. J. Application and sensitivity investigation of Fourier transforms for microwave radiometric inversions [NASA-CR-140754] p0070 N75-11310 HOLTZMAN, J. C. Simultaneous active and passive microwave response of the earth - The Skylab radscat experiment p0012 A75-17114	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and near-infrared p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422 IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667 INGHAM, J. D. Microbial abundance and thermoluminescence of antarctic dry valley soils p0005 A75-17737 INGLE, S. J. A study of the early detection of insect infestations and density/distribution of host plants [E75-10062] INGRAO, H. C. Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0088 A75-17208 ISACHSEN, Y. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] ISHAO, A. M. Application of remote sensing to the location of hydrologically active /source/ areas
and urban terrain analysis HAY, C. M. A gricultural inventory techniques with orbital and high-altitude imagery HAYDN, R. Evaluation of ERTS-1 data by analogue and digital techniques Geologic evaluation of central Italy from ERTS-1 and Skylab data HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] PO079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration HEINTZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0019 N75-11553 HEISS, K. P. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141193] The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] PO079 N75-14203 HEMPENIUS, S. A. The ITC 8-image comparator and its possible use in ERTs and EREP investigations PO068 A75-17221 HENDERSON, F. M. Small scale thematic mapping - A case for radar imagery PO068 A75-18062	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104] HOFFMAN R. Applications of remote sensing by the state of Nebraska p0069 A75-17232 HOLLYDAY, E. F. Hydrologic significance of lineaments in central Tennessee [E75-10030] HOLMES, J. J. Application and sensitivity investigation of Fourier transforms for microwave radiometric inversions [NASA-CR:140754] HOLTZMAN, J. C. Simultaneous active and passive microwave response of the earth - The Skylab radscat experiment p0012 A75-17114	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and near-infrared p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422 IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667 INGHAM, J. D. Microbial abundance and thermoluminescence of antarctic dry valley soils p0005 A75-17973 INGLE, S. J. A study of the early detection of insect infestations and density/distribution of host plants [E75-10062] p0008 N75-13362 INGRAO, H. C. Development of an experimental airbome laser remote sensing system for the detection and classification of oil spills p0068 A75-17208 ISACHSEN, Y. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] p0034 N75-10536 ISHAQ, A. M. Application of remote sensing to the location of p0066 A75-17141 IURKIN, IU. T. Detection of synchronous changes in fluxes of energetic electrons and gamma-quanta during a magnetic dURKINA, M. I.
and urban terrain analysis p0055 A75-17199 HAY, C. M. A gricultural inventory techniques with orbital and high-altitude imagery p0001 A75-12872 HAYDN, R. Evaluation of ERTS-1 data by analogue and digital p0059 N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration p0075 A75-13592 HEINTZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0019 N75-11553 HEISB, K. P. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-1411268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HEMPENIUS, S. A. The ITC 8-image comparator and its possible use in ERTS and EREP investigations p0068 A75-17221 HENDERSON, F. M. Small scale thematic mapping - A case for radar p0028 A75-18062	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10072] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10074] HOFFMAN, R. Applications of remote sensing by the state of Nebraska p0073 N75-15122 HOLLYDAY, E. F. Hydrologic significance of lineaments in central Tennessee [E75-10030] P0050 N75-12401 HOLMES, J. J. Application and sensitivity investigation of Fourier transforms for microwave radiometric inversions [NASA-CR-140754] p0070 N75-11310 HOLTZMAN, J. C. Simultaneous active and passive microwave response of the earh - The Skylab radscat experiment p0012 A75-17114 HOLZ, R. K. A hydrogeomorphic approach to evaluating flood potential in central Texas from orbital and suborbital remote sensing imagery p0047 A75-17140	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and near-infrared p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422
and urban terrain analysis HAY, C. M. A gricultural inventory techniques with orbital and high-altitude imagery HAYDN, R. Evaluation of ERTS-1 data by analogue and digital techniques Geologic evaluation of central Italy from ERTS-1 and Skylab data HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] HECKER, E. J. NASA-KSC's earth resources benefits from space exploration HEINZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0079 N75-11553 HEISS, K. P. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141193] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141193] The economic value of remote sensing by satellitie: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] HEMPENUS, S. A. The ITC 8-image comparator and its possible use in ERTS and EREP investigations HEMPENUS, S. A. The ITC 8-image comparator and its possible use in ERTS and EREP investigations HENDERSON, F. M. Small scale thematic mapping - A case for radar imagery HENNEY, M. R Skylab-environmental-and-crew-microbiology-studies—p0024 N75-14366	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] p0057 N75-10531 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] p0078 N75-12390 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] p0037 N75-14191 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104] p0037 N75-14191 HOFFMAN, R. Applications of remote sensing by the state of Nebraska p0069 A75-17232 HOLLYDAY, E. F. Hydrologic significance of lineaments in central Tennessee [E75-10030] p0050 N75-12401 HOLTZMAN, J. C. Simultaneous active and passive microwave response of the earth - The Skylab radscat experiment p0012 A75-171140 HOLZ, R. K. A hydrogeomorphic approach to evaluating flood potential in central Texas from orbital and suborbital remote sensing imagery p0047 A75-17140 HONEY, F. R. Feasibility of using-S-191 infrared spectra-for geological—studies from space	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422 IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667 INGHAM, J. D. Microbial abundance and thermoluminescence of antarctic dry valley soils p0005 A75-17973 INGLE, S. J. A study of the early detection of insect infestations and density/distribution of host plants [E75-10062] p0008 N75-13382 INGRAO, H. C. Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0088 A75-17208 ISACMSEN, Y. W. Assessment of ERTS-1 imagery as a tool for regional geological enalysis in New York State [E74-10809] ISHAQ, A. M. Application of remote sensing to the location of hydrologically active /source/ areas p0084 A75-12317 IURKIN, IU. T. Detection of synchronous changes in fluxes of energetic electrons and gamma-quanta during a magnetic disturbance p0084 A75-12317 IURKINA, M. I. On the determination of secular variation in the earth's gravity field and of secular variation in the earth's gravity field and of secular variation in the earth's p0084 A75-18125
and urban terrain analysis HAY, C. M. Agricultural inventory techniques with orbital and high-altitude imagery HAYDN, R. Evaluation of ERTS-1 data by analogue and digital poops N75-14218 Geologic evaluation of central Italy from ERTS-1 and Skylab data p0038 N75-14238 HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HECKER, E. J. NASA-KSC's earth resources benefits from space exploration p0075 A75-13592 HEINTZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0019 N75-11553 HEISS, K. P. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141193] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141193] p0079 N75-14203 The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204 HEMPENIUS, S. A. The ICS -image comparator and its possible use in ERTS and EREP investigations p0060 N75-14222 HEMPERSON, F. M. Small scale thematic mapping - A case for radar imagery HENNEY, M. RSkylab-environmental-and-crew-microbiology-studies—p0024 N75-14366	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10014] P0037 N75-14191 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104] P0073 N75-14191 HOFFMAN, R. Applications of remote sensing by the state of Nebraska HOLLYDAY, E. F. Hydrologic significance of lineaments in central Tennessee [E75-10030] HOLMES, J. J. Application and sensitivity investigation of Fourier transforms for microwave radiometric inversions [NASA-CR-140754] HOLZMAN, J. C. Simultaneous active and passive microwave response of the earth - The Skylab radscat experiment p0012 A75-17114 HOLZ, R. K. A hydrogeomorphic approach to evaluating flood potential in central Texas from orbital and suborbital remote sensing imagery p0047 A75-17140 HONEY, F. R. Feasibility of using-S-191 infrared spectra-for geological—studies from space [E75-10088]	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422 IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667 INGHAN, J. D. Microbial abundance and thermoluminescence of antractic dry valley soils p0005 A75-1773 INGLE, S. J. A study distribution of host plants [E75-10062] p0008 N75-13362 INGRAO, H. C. Development of an experimental airbome laser remote sensing system for the detection and classification of oil spills ISACHSEN, Y. W. Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State [E74-10809] ISHAQ, A. M. Application of remote sensing to the location of hydrologically active / source/ areas p0066 A75-17141 IURKIN, IU. T. Detection of synchronous changes in fluxes of energetic electrons and gamma-quanta during a magnetic p0064 A75-12317 IURKIN, M. I. T. On the determination of secular variation in the earth's gravity-field and of-secular-polar-motion p0029 A75-18125
and urban terrain analysis HAY, C. M. A gricultural inventory techniques with orbital and high-altitude imagery HAYDN, R. Evaluation of ERTS-1 data by analogue and digital techniques Geologic evaluation of central Italy from ERTS-1 and Skylab data HAZELRIGG, G. A., JR. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] HECKER, E. J. NASA-KSC's earth resources benefits from space exploration HEINZENBERG, J. Scattering and absorption from poly-dispersed aerosols p0079 N75-11553 HEISS, K. P. The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141193] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141193] The economic value of remote sensing by satellitie: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] HEMPENUS, S. A. The ITC 8-image comparator and its possible use in ERTS and EREP investigations HEMPENUS, S. A. The ITC 8-image comparator and its possible use in ERTS and EREP investigations HENDERSON, F. M. Small scale thematic mapping - A case for radar imagery HENNEY, M. R Skylab-environmental-and-crew-microbiology-studies—p0024 N75-14366	HODAPP, C. J. Skylab environmental and crew microbiology studies p0024 N75-14366 HOFER, R. Emissivities and forward scattering of natural and man-made material at three millimeter wavelength p0067 A75-17197 HOFFER, R. M. An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E74-10803] p0057 N75-10531 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10002] p0078 N75-12390 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using automatic data processing techniques [E75-10072] p0037 N75-14191 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques [E75-10104] p0037 N75-14191 HOFFMAN, R. Applications of remote sensing by the state of Nebraska p0069 A75-17232 HOLLYDAY, E. F. Hydrologic significance of lineaments in central Tennessee [E75-10030] p0050 N75-12401 HOLTZMAN, J. C. Simultaneous active and passive microwave response of the earth - The Skylab radscat experiment p0012 A75-171140 HOLZ, R. K. A hydrogeomorphic approach to evaluating flood potential in central Texas from orbital and suborbital remote sensing imagery p0047 A75-17140 HONEY, F. R. Feasibility of using-S-191 infrared spectra-for geological—studies from space	imagery p0047 A75-17140 HUNT, G. R. Remote sensing of rock type in the visible and p0033 A75-17220 HURD, D. Image selection system [NASA-CR-2475] p0058 N75-12422 IEREMEEV, V. F. On the determination of secular variation in the earth's gravity field and of secular polar motion p0029 A75-18125 IIZUKA, K. Performance of a hologram matrix radar p0065 A75-14792 ILINSKII, V. N. Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-17667 INGHAM, J. D. Microbial abundance and thermoluminescence of antarctic dry valley soils p0005 A75-17973 INGLE, S. J. A study of the early detection of insect infestations and density/distribution of host plants [E75-10062] p0008 N75-13382 INGRAO, H. C. Development of an experimental airborne laser remote sensing system for the detection and classification of oil spills p0088 A75-17208 ISACMSEN, Y. W. Assessment of ERTS-1 imagery as a tool for regional geological enalysis in New York State [E74-10809] ISHAQ, A. M. Application of remote sensing to the location of hydrologically active /source/ areas p0084 A75-12317 IURKIN, IU. T. Detection of synchronous changes in fluxes of energetic electrons and gamma-quanta during a magnetic disturbance p0084 A75-12317 IURKINA, M. I. On the determination of secular variation in the earth's gravity field and of secular variation in the earth's gravity field and of secular variation in the earth's p0084 A75-18125

KELEJIAN H H KOMHYR, W. IVANOV, V. The earth in the lens of Salyut-3 The value of improved (ERS) information based on Geophysical monitoring for climatic change. The NOAA p0070 N75-10964 p0018 N75-11510 [NASA-TT-F-16028] domestic distribution effects of U.S. agriculture crops program [NASA-CR-141046] p0007 N75-12423 KONDRATEV, K. IA. IZOTOV, A. A. [NASA-CR-141040]
The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources, agriculture. Parl 2: Distribution effects Spectrophotometric observations of earth Soyuz-13 Status and prospects of development of space geodesy [JPRS-63231] p0029 N75-10869 p0029 N75-10869 [IAF PAPER 74-156] p0065 A75-13765 Space iconics The application of spectrometric and polarization techniques for remote sensing of oil on sea water p0068 A75-17210 p0057 N75-11419 [NASA-CR-141194] p0079 N75-14206 KELLER, M. Skylab a proposal aerotriangulation with very small scale J KONDRATYEV, K. Y. photography [E74-10804] Influence of the atmosphere on spectral radiance and p0070 N75-10532 contrasts of natural formations measured from red from space p0019 N75-11557 Skylab: A proposal aerotriangulation with very small JACOBOWITZ, H. scale photography [E75-10101] A cloud physics investigation utilizing Skylab data [E75-10021] p0020 N75-1 KONKOV, S. A. p0020 N75-12400 p0073 N75-15119 Methods for the automatic recording of atmospheric dust KEMMERER, A. J.

A review of initial investigations to utilize ERTS-1 data A cloud physics investigation utilizing Skylab data [£75-10099] p0025 N75-1 concentrations and some of the results obtain p0025 N75-15117 p0057 N75-11502 in determining the availability and distribution of JARMAN, J. W. p0043 N75-10555 KONSHIN, M. D. marine resources ERTS program of the US Army Corps of Engineers a panoramic aerial camera KENNEY, J. Use of a [AD-784162] p0042 N75-10554 Multi-frequency radiometric measurements of foan p0071 N75-11450 JAYROE, R. R. p0068 A75-17208 mono-molecular slick KORCHAGIN, E. K. Computer and photogrammetric general land use study KHARITONOVA, T. V. The wind vector in a layer over water Investigations of diffuse cosmic gamma radiation in the range 28 keV-4.1 MeV p0015 A75-1766 of central north Alabama p0041 A75-15179 [NASA-TR-R-431] p0022 N75-13372 p0015 A75-17667 KORNHAUSER, A. L JOATTON, R. KHODAIR M. M. The economic value of remote sensing of earth resources The 1973 program of measurement of the minor Spectral reflectance studies on mineral deficiency in corn from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources: Agriculture. Part 1: Overview constituents of the stratosphere using the Concorde 001 nlants n0003 A75-17172 p0063 A75-11720 KIEFER, R. W. The application of ERTS-derived information to the JOHNSON, D. A. [NASA-CR-141265] p0079 N75-14205 A technique for measurement of sulphur dioxide and regional land use planning process p0015 A75-18058 KORUNOV S. S. sulphate concentrations from an aircraft Economic effectiveness of using the space systems of D0018 N75-11498 Optical film density values from color IR photograph distant indication of natural resources [IAF PAPER 74-165] for wetland soils mapping KING, R. B. p0005 A75-18073 JOHNSON, W. R. p0075 A75-13770 Satellite photography of eddies in the gulf loop current p0041 A75-11255 KOTLARSKI, J. R. A comparison of ERTS imagery with conventional serial Oil slick detection by X-band synthetic aperture rada photography for land-resource surveys in less developed n0068 A75-17209 countries. Examples from the Rift Valley Lakes Basin. The Proceedings of the Skylab Life Sciences Symposium, KOVALENKO, V. A. p0081 N75-14249 Automated remote measurement system in agricultural KIRILLOV-UGRIUMOV, V. G. [NASA-TM-X-58154-VOL-1] p0081 N75-14358 meteorology p0064 A75-12947 Detection of synchronous changes in fluxes of energetic electrons and gamma-quanta during a magnetic gamma-quanta during a magnetic p0064 A75-12317 remote measuring system in p0007 N75-12518 Automated Implementation of an advanced table look-up classis agrometeorology disturbance for large area land-use classification p0014 A75-17153 KIRVIDA, L. KRISTOF, S. J. JONES, R. C. Automatic land use classification using Skylab S-192 Mapping soil features from multispectral scanner data logic evaluation and applications of ERTS-1 imagery p0027 A75-15435 multispectral data p0034 N75-10552 over Georgia [AIAA PAPER 74-1224] p0064 A75-12036 KRITIKOS, G. JOYCE, A. T. Mapping of water bodies in northern Germany from ERTS KLAINER, S. M. p0017 N75-10560 p0051 N75-14227 Land use and mapping The remote Raman spectrometer is a viable instrument KROECK, R. M. for remote sensing of the environment JUDAY, R. D. Summary of flight performance of the Skylab Earth Resources Experiment Package / EREP/ D0068 A75-17214 Image selection system [NASA-CR-2475] KLANKAMSORN, B. p0058 N75-12422 p0076 A75-17211 KRONBERG, P. ERTS applications in Thailand - A progress report p0076 A75-17123 p0031 N75-14239 JUSTICE, C. O. ERTS-1 data on Afar tectonics The interpretation and use of false-colour infra-red and KLAUBER, G. Trace gas analysis of power plant plumes via aircraft measurement - O3, NOx, and SO2 chemistry p0064 A75-13148 Skylab environmental and crew microbiology studies p0024 N75-14366 true colour photography of part of Argentina obtained by Skylark earth resources rockets p0067 A75-17192 A comparison of the capabilities of ERTS-1 and the KUHLOW, W. W. KLAUSNER. S. earth-resources rocket for resources surveying in Interactive machine assessment of critical land resources using ERTS-1 data p0013 A75-17149

The application of ERTS-derived information to the p0023 N75-14250 Modifications in a computer-implemented method for the central Argentina detection and extraction of objects in aerial photographs p0055 A75-17154 regional land use planning process p0015 A75-18058 Κ KUHNER, M. Skylab and ERTS-1 investigations of coastal land-use and water properties in Delaware Bay [AIAA PAPER 74-1220] p0041 A75-12034 Calibration and evaluation of Skylab altimetry for geodetic determination of the geoid JA, R.
Multispectral photography for earth resources
p0074 N75-15138 [E75-10064] p0030 N75-13364 Coastal and estuarine studies with ERTS-1 and Skylab p0047 A75-16240 KUMAR, R [NASA-CR-141283] Statistical separability of agricultural cover types in KALININ, G. P. Correlation of coastal water turbidity and current From aerospace photographs to forecasting and calculating flows
[NASA-TT-F-16006] p0059 N75-13367 subsets of one to twelve spectral channels Correlation of coastal water to constitute circulation with ERTS-1 and Skylab imagery p0048 A75-17181 p0005 A75-17216 KUNDE, V. G. Estimation of sea surface temperature from remote sensing in the 11to 13-micron window region p0065 A75-14803 Inventories of Delaware's coastal vegetation and land-use KALISZEWSKI, T. Bistatic radar sea state monitoring [NASA-CR-137469] utilizing digital processing of ERTS-1 imagery p0004 A75-17186
Monitoring physical and chemical parameters of Delaware p0043 N75-10673 Bay waters with an ERTS-1 data collection platform [E75-10090] p0051 N75-15108 Signatures of various earth surfaces measured by the Thermal emission of agglomerations as recorded on Nimbus-5 microwave spectrometer p0066 A75-17119 KURDINOVSKII, O. U. environmental satellites and ERTS-1 imagery p0059 N75-14217 mental satellite imagery KLESHCHENKO, A. D. p0011 A75-13757 Automated remote measurement system in agricultural eteorology p0064 A75-12947 Economic effectiveness of using the space systems of distant indication of natural resources
[IAF PAPER 74-165] meteorology p0075 A75-13770 Automated ring system in p0007 N75-12518 KARMELI. D. KURIYAN, J. G. agrometeorology Proceedings of the UCLA International Conference on Radiation and Remote Probing of the Atmosphere KLETT, A. T. TT, A. T.
Utility of ERTS for monitoring the breeding habit of igratory waterfowl p0006 N75-10557

Modifications in a computer-implemented method for the detection and extraction of objects in aerial photographs p0055 A75-17154

į.

KASHIN, L.A. The next problems in the area of the implementation of planning for cartographical-geodetic work [AD-785882] pO

p0032 N75-15143 KATIUSHINA, V. V. Morning-evening asymmetry of the thermosphere during p0011 A75-11103 neomagnetic disturbance

KATSIERIS, D. Preliminary results of the interpretation of ERTS-1

imagery for soil surveys in the Merida region, Spain p0009 N75-14247

KAUTZLEBEN, H.

Geodesy and physics of the earth; Proceedings of the Second International Symposium, Potsdam, East Gerr May 7-11, 1973. Parts 1 & 2 p0028 A75-18106 Problems and consequences of modern hypotheses on the global tectonics for geodesy and gravimetry p0028 A75-18109

KEATHLEY, W. C. Apollo Telescope Mount experiments technology DO064 A75-13589

Environmental definition program [UCID-16565] p0022 N75-13458 Atmospheric release advisory capability: Research and

progress p0025 N75-15202 [UCRL-75644-REV-2]

p0072 N75-11454

p0058 N75-12422

KNUTSON, M. A. on system

Image selection [NASA-CR-2475]

migratory waterfowl

Oil-on-Water sensor

KNIGHT. W

KNOX, J. B.

[AD-784104]

KÔCH, M.

ERTS image contribution to geological knowledge overing different types of climates (France, Togo, Dahomey, fadagascar) p0038 N75-14240 Madagascar) KOMAROV, V. B.

Development of research in the USSR on the use of radar images for geological purposes p0033 A75-17126 KUSCER. I. Some analytical results for radiative transfer in thick

[NASA-CR-140816]

KUROCHKIN, A. V.

atmospheres

off Spanish Sahara

L LA VIOLETTE, P. E. A satellite-aircraft thermal study of the upwelled waters p0041 A75-10645

Particulate sizes from polarization measurements

Detection of synchronous changes in fluxes of energetic electrons and gamma-quanta during a magnetic disturbance p0064 A75-12317

Qualitative study of some ERTS-1 images of the Golion project. Ecological aspects of terrestrial environments p0023 N75-14248

B-7

p0078 N75-11549

p0019 N75-11558

p0019 N75-11555

LAKHTANOV, G. A.

The application of spectrometric on particular techniques for remote sensing of oil on sea water p0068 A75-17210

LALIEMAND C.

Comparison of various numerical techniques for classification of multispectral data. Application to geology p0038 N75-14244

LAMAR, D. L.

Lineaments in basement terrane of the Peninsular Ranges, Southern California p0035 N75-11407 [E75-10003] Fault tectonics and earthquake hazards in the Peninsular

Ranges, Southern California [E75-10022] n0030 N75-13338

Analysis of pseudocolor transformations of ERTS-1 images of Southern California area DO036 N75-13346

[E75-10037] LAMAR, J. V. Analysis of pseudocolor transformations of ERTS-1

images of Southern California area [E75-10037] p0036 N75-13346

Water depth penetration film test p0047 A75-12873 LANDINI A. J.

The availability of local aerial photography in southern California p0025 N75-15128

NASA-CR-141314) LARSEN, P. A.

Computer and photogrammetric general land use study of central north Alabama p0022 N75-13372 [NASA-TR-R-431]

LARSON, R. W. Investigation of microwave hologram techniques for

p0067 A75-17195 application to earth resources

Areal extent of snow in forested regions - A practical estimation technique using ERTS-1 data p0002 A75-17122

LAVROVA, V. S.

Use of a digital computer to construct isoline maps based on a stereomodel of terrain [AD-784170] p0057 N75-11453

Microbial abundance and thermoluminescence

antarctic dry valley soils LEBEDEV, V. V. p0005 A75-17973

Spectrophotometric observations of earth from [IAF PAPER 74-156] n0065 A75-13765

Thermal behaviour of some rice fields affected by a vellows-type disease D0004 A75-17174

Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcanoes [E75-10069] p0037 N75-14188 Contribution of space platforms to a ground and airborne remote-sensing programme over active Italian volcanoes p0073 N75-14236

LEDLEY, B. G.

Magnetopause rotational forms p0011 A75-11221

Geologic and mineral and water resources investigations in Western Colorado, using Skylab EREP data p0034 N75-10530

Index of Skylab data available at Colorado School of

p0072 N75-12410 [E75-10041] Geologic and mineral and water resources investigations Western Colorado, using Skylab EREP data

[F75-10043]

[E79-10043] Goologic and mineral and water resources investigations in western Colorado, using Skylab EREP data [E75-10071] p0037 N75-14190 [E75-10071] p0037 N75-14190 Geologic and mineral and water resources investigations

in western Colorado, using Skylab EREP data p0025 N75-15123 [E75-10105]

LEE. S. L. On the feasibility of remote monitoring of soil mois with microwave sensors p0067 A75-17147

LEE, S. S. Application of remote sensing for prediction and detection

of thermal pollution [NASA-CR-139182] DO017 N75-10572

Feasibility of remote sensing for detecting thermal pollution. Part 1: Feasibility study. Part 2: Implementation plan [NASA-CR-134453]

n0025 N75-15199

LEGENDRE, G.

Comparison of various numerical techniques for classification of multispectral data. Application to geology p0038 N75-14244

Coherent optics potential applications to mapping p0027 A75-14683 Coherent optics in mapping: Proceedings of the Tutorial Seminar and Technology Utilization Program, Rochester, N.Y., March 27-29, 1974 p0027 A75-16676

LEIKOV, N. G.

Detection of synchronous changes in fluxes of energetic electrons and gamma-quanta during a magnetic disturbance p0064 A75-12317 disturbance LEITAO, C. D.

Skylab altimeter applications and scientific results [AIAA PAPER 74-1221] p0063 A75-12035

Skylab radar altimeter - Short-wavelength perturbations atected in ocean surface profiles p0066 A75-16096 detected in ocean surface profiles LEPLEY, L. K.

A study to explore the use of orbital remote sensing to determine native arid plant distribution [E74-10818] n0006 N75-10544

LEWIS, A. J. Optical film density values from color IR photography wetland soils mapping p0005 A75-18073 for wetland soils mapping

LEWIS, D. T. Satellite pictures can help guide Sandhills irrigation [E75-10023] p0021 N75-13339 Use of ERTS-1 imagery to interpret wind-erosion hazard

Sandhills of Nebraska [E75-10044] n0059 N75-13347

LHOMMER, A.

Recent and present sedimentation complexity of the Rhone delta and the French mediterrar p0045 N75-14228 project)

LIDSTER, W. A.

Evaluation of ERTS-1 and aircraft data for assessing internal drainage in irrigated agriculture p0002 A75-17143

UEGEL L H. An intensive natural resources inventory aids land use planning and resources management in Pu p0005 A75-18056

The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction nd overview

p0080 N75-14210 [NASA-CR-141266] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 7: Nonreplenishable natural resources: Minerals, fossil fuels and geothermal energy sources

p0080 N75-14212 [NASA-CR-141264] The economic value of remote sensing of earth resources m space: An ERTS overview and the value of continuity service. Volume 9: Oceans

of service. Volume 9
[NASA-CR-141270] p0080 N75-14214 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity Volume 10: Industry [NASA-CR-141192] p0080 N75-14215

LIN. W. C. Detection of moisture and moisture related phenomena from Skylab

[E75-10073] p0008 N75-14192 LINDGREN, D. T.

Investigation of land use of northern megalopolis using ERTS-1 imagery
[NASA-CR-2459] p0017 N75-10545

LINLOR, W. I. Microwave profiling of snowpack free p0049 A75-17950

LINS. H. F., JR Urban and regional land use analysis: CARETS and census cities experiment package

p0020 N75-12398 5-10019] Urban and regional land use analysis: CARETS and nsus cities experiment package

p0022 N75-13356 Urban and regional land use analysis: CARETS and nsus cities experiment package

[F75-10091] p0024 N75-15109 Urban and regional land use analysis: CARETS and ensus cities experiment package

n0025 N75-15118 [E75-10100] LINTON, L. W The GLOBAL computer program and GLOBEMAP: A

special purpose three-dimensional plotter subroutine [AD-785869] p0032 N75-1 p0032 N75-15141 LIPATOV, V. B.

Dust storms according to data of space research [NASA-TT-F-16021] p0021 N75p0021 N75-12463 LOCKWOOD H. E.

Water depth penetration film test p0047 A75-12873 LOEFFLER, A. Reduction of information redundancy in ERTS-1 and

EREP data. A proposal for an experiment p0060 N75-14225

Qualitative study of some ERTS-1 images of the Golion

project. Ecological aspects of terrestrial environments p0023 N75-14248

LOVE. A. W.

Measurements of the dielectric properties of sea water at 1.43 GHz D0045 N75-13496 [NASA-CR-2458]

Comparison of the Southern and Northern Hemisphere general circulation characteristics as determined by satellite [UCRL-75556] D0017 N75-10638

Environmental definition program
[UCID-16565] D0022 N75-13458 LOWE, D. S.

Evaluation of ERTS data utilization in developing D0055-A75-17193 countries

LOWMAN, P. D., JR. Geologic structure in California: Three studies with ERTS-1 imag [NASA-TM-X-70799] DO039 N75-15132 LUCHKOV, B. I.

Detection of synchronous changes in fluxes of energetic on of synchronous changes in magnetic and gamma-quanta during a magnetic p0064 A75-12317 electrons disturbance

LUDWIG, C. B. Global measurements of air pollution from satellites

p0018 N75-11521 Cloud screening from aerial photography applying

coherent optical pattern recognition techniques p0054 A75-18690

LUTSYCHIN, A. S.

A generator of high-power current pulses of semisinusoidal shape for an aerial variant of the method p0053 A75-13310 of transient processes LUTZEN, E. E.

Land-use planning aided by computer modelling/mapping system to combine remote sensing.
natural resources, social and economic data p0076 A75-17118

An analysis of applications development systems for remotely sensed, multispectral data for the Earth Observations Division of the NASA Lyndon B. Johnson Space Center

[NASA-CR-140345] LYON, J. P.

Feasibility of using S-191 infrared spectra for geological studies from space [E75-10088] n0038 N75-15106

Feasibility of using S-191 infrared spectra for geological studies from space [F75.10089]

p0038 N75-15107 LYZENGA, D. R.

Skylab water depth determination [E75-10048] p0050 N75-13348

М

MACARTHUR, J. L.

A radar altimeter for satellite application [AD-784373] p0056 N75-10425 MACDONALD, R. B.

Agriculture, forestry, range resources

p0006 N75-10558 MACOMBER, R. T.

Application of ERTS-1 data to the protection and management of New Jersey's coastal environment p0042 A75-17233

MADDING, R. P. Scanning thermal plumes DO014 A75-17162

MAIRS, R. L. Application of ERTS-1 data to the protection and management of New Jersey's coastal environment

p0042 A75-17233 MAKAROV, V. I. Plutonic structure of the earth's crust

p0027 A75-17125 MALILA, W. A.

First results from the crop identification technology assessment for remote sensing / CITARS/ p0004 A75-17175

Multi-aspect techniques in remote sensing p0068 A75-17217 Developing processing techniques for Skylab data

[F74-10799] n0057 N75-10527 Developing processing techniques for Skylab data

[E75-10040] p0058 N75-12409 Developing processing techniques for Skylab data [E75-10082] p0061 N75-1 p0061 N75-15102

MALKMUS, W. Global measurements of air pollution from satellite

p0018 N75-11521

Theoretical investigations on the determination of the vertical temperature distribution in the atmosphere assuming multispectral thermal radiation measurements at 60 GH2 [BMFT-FB-W-74-04] p0024 N75-14331

MAMAEV. V. A.

A generator of high-power current pulses semisinusoidal shape for an aerial variant of the meth p0053 A75-133 current pulses of p0053 A75-13310 MANDUROVSKII, I. A.

The wind vector in a layer over water

p0041 A75-15179

1.1.

MARCHUK, G. I. Solution of the radiative transfer theory problems b Monte Carlo method p0058 N75-11551 MARINO, C. M.

Remote sensing techniques applied to the study of Italian volcanic areas - The results of the repetition of the airborne

I.R. survey compared to the previous data p0069 A75-17223

remote sensing programme over active Italian volcanoes [E75-10069] p0037 N75-14188

Geological features on ERTS-1-images of a test a the west-central Alps p0037 N75-14237 MARTIN, A. M.

Environmental satellites and El\tal{1}\tal{1}\tal{1}\tal{1}\tal{1}\tal{1}\tal{1}\tal{1}\tal{2}\tal{1

MCQUEEN, J. L. MILLER, W. F. MARTIN, H. L. Life cycle cost study of Army Spectrometric Oil Analysis Program (ASOAP) Skylab environmental and crew microbiology studies p0024 N75-14366 Application of remote sensing to state and regional problems p0073 N75-14870 MEAD, R. A [AD-786501] NASA-CR-1408051 n0078 N75-11414 ERTS-I data for classifying native plant com-MARTIN, L U. MILLER, W. L. Central Colorado The economic impact of remote sensing data as the source of nonpoint pollution monitoring and control p0004 A75-17176 Multi-frequency radiometric measuren p0068 A75-17208 MEALOR, W. T. mono-molecular slick Correlation of multispectral imagery with water analysis p0076 A75-17166 MARUYASU, T. - The Ross Barnett Reservoir Remote Investigation of the environmental change pattern of MINAEV. E. S. p0014 A75-17165 Economic effectiveness of using the space systems of Japan MEEKS, D. C. [F75-10086] n0024 N75-15104 distant indication of natural resources A study of a AIDJEX 1972 of microwave emission prop perties of sea ice -p0041 A75-17120 (IAF PAPER 74-165) p0075 A75-13770 MATHER, R. S. The gravity field and the definition of stationary se-MINTER, T. C., JR. MEIER C. J. p0042 A75-18124 Feasibility study ASCS remote sensing/compliance surface topography Multidisciplinary applications of ERTS and Skylab data determination system MAUGHAN, P. M. p0015 A75-17234 in Ohio A review of initial investigations to utilize ERTS-1 data [NASA-CR-134288] p0007 N75-12419 MEIER, M. F. in determining the availability and distribution of living Microwave profiling of snowpack free-water content MINTZER O.W. p0043 N75-10555 Remote sensing for regional resource analysis marine resources p0049 A75-17950 p0015 A75-18061 Evaluate ERTS imagery for mapping and detection of MAUL, G. A Satellite photography of eddies in the gulf loop current p0041 A75-11255 changes of snowcover on land and on glaciers [E74-10812] p0049 p0049 N75-10539 Electrically scanning microwave radiometers p0056 A75-17202 MEL, M. R. Remote sensing of ocean current boundary layer n0042 N75-10529 [E74-10801] The usefulness of imaging passive microwave for rural MOELLER, H. urban terrain analysis p0055 A75-17199 An electronic drawing aid for map compiling An assessment of the potential contributions to oceanography from Skylab visual observations and n0027 A75-10076 MENZIES, A. T. Remote atmospheric sensing with an airborne laser psorption spectrometer p0066 A75-16485 MOHR, P. A. hand-held photography p0044 N75-12396 [E75-10017] absorption spectrometer Mapping of the major structures of the African rift MERCANTI, E. P. Remote sensing of ocean current boundary layer 75-10031] p0045 N75-14185 system [E75-10024] Third Earth Resources Technology Satellite Symposium. [E75-10031] p0030 N75-13340 Volume 2: Summary of results MONGET, J. M. Remote sensing of ocean current boundary layer 75-10067] p0045 N75-14186 [NASA-SP.356] p0077 N75-10549 Colour composites of ERTS multispectral imagery for Colour composites or End oceanography applications in hydrology and oceanography p0045 N75-14231 [E75-10067] MERCHANT, J. W., JR. Remote se oundary layer p0046 N75-15116 The use of high altitude photography and ERTS-1 imagery [E75-10098] for wildlife habitat inventory in Kansas MOODY, H. J. MAURER, H. D0005 A75-18066 A microwave correlation spectrometer for air pollution MFREMBECK, B. F. Texture analysis with Fourier series n0004 A75-17187 p0063 A75-11732 Classification and mapping of coal refuse, vegetative MAY. G. A. MOONEY, H method of specifying remotely sense cover types, and forest types by digital processing ERTS-1 ed units for soi A preliminary feasibility study on a synthetic aperture p0054 A75-17110 p0002 A75-17106 sample points Transference of ERTS-1 spectral signatures in time and radar /SARSAT/ for earth resources su MAYO, W. T., JR. p0067 A75-17194 p0066 A75-17111 Remote measurements of water pollution with a I MOORE, B. H.
Feasibility study ASCS remote sensing/compliance MERIFIELD, P. M. p0014 A75-17205 polarimete Lineaments in basement terrane of the Peninsular Ranges. MAZETS, E. P. Southern California determination system of diffuse cosmic gamma radiation in the [NASA-CR-134288] p0007 N75-12419 [E75-10003] p0035 N75-11407 range 28 keV-4.1 MeV p0015 A75-17667 Fault tectonics and earthquake hazards in the Peninsu Ranges, Southern California [E75-10005] p0035 N75-114 MOORE, D. G.
Evaluation of ERTS-1 and aircraft data for assessing MCALLUM, W. E.

The effect of atmospheric water vapor on autom on automatic pO013 A75-17131 Summary of flight performance of the Skylab Earth Resources Experiment Package /EREP/ n0035 N75-11408 internal drainage in irrigated agriculture Fault tectonics and earthquake hazards in the Peninsular p0002 A75-17143 Ranges, Southern California MOORE, G. K. p0030 N75-13338 [E75-10022] Flood inundation in the southeastern United States from Analysis of pseudocolor transformations of ERTS-1 nages of Southern California area aircraft and satellite imagery p0047 A75-17138 MCCASLIN, M. Hydrologic significance of lineaments in central An assessment of the potential contributions to ceanography from Skylab visual observations and p0036 N75-13346 Fault tectonics and earthquake hazards in the Peninsular [E75-10030] p0050 N75-12401 hand-held photography es, Southern California [F75-10017] n0044 N75-12396 MOORE H.I [E75-10074] p0031 N75-14193 MCCAULEY, J Forest defoliation assessment with satellite image Fault tectonics as d earthquake hazards in the Peninsular p0003 A75-17171 Ontical data processing analysis of stream patterns exhibited on ERTS-1 imagery MCCAULEY, J. R. p0056 A75-17219 Ranges, Southern California MOORE, R. K. p0039 N75-15124 [F75-10108] wave measurement of soil water content 6851 p0001 A75-16242 Active mic INASA-CR-141685] Skylab study of water quality [E75-10075] MESZAROS, E. p0051 N75-14194 Aircraft measurements of the concentration and size Simultaneous active and passive microw of the earth - The Skylab radscat experiment e microwave response MCCLAIN, E. P. distribution of atmospheric sulfate particles Images from the NOAA-3 very high resolution radiometer p0057 N75-11504 p0012 A75-17114 Investigation of radar discrimination of sea ice p0042 A75-17121 over the North Sea and adjoining countries MIESZKUC B. J. p0066 A75-16188 Skylab environmental and crew micro MCCLENDON, R. W. Design data collection with Skylab/EREP microwave p0024 N75-14366 Mathematical modeling for the detection of fish by an instrument S-193 MIKHAIL, E. M. p0065 A75-15953 [E75-10070] DO059 N75-14189 Hologrammetry - Concepts and applications MCCORMACK, K. Preliminary analysis of Skylab RADSCAT results over p0065 A75-15434 Oil-on-Water senso [AD-784104] the ocean MIKHAILOV, G. A NASA-CR-141454] p0072 N75-11454 p0046 N75-14793 Solution of the radiative transfer theory problems by the MCCORMICK, N. J. MOORE, R. P. p0058 N75-11551 Monte Carlo method Some analytical results for radiative transfer in thick The usefulness of imaging passive microwave for MIKHAILOVA, S. M. p0019 N75-11555 p0055 A75-17199 and urban terrain analysis The application of spectrometric and polarization MCCRICKERD, J. T. Microwave radiometric characteristics of snow-covere Terrain measurement from holograph techniques for remote sensing of oil on sea water p0049 A75-17200 p0068 A75-17210 p0001 A75-16682 MORENO, G. MCFADDEN, J. MILAZZO V. A. Solar wind and geomagnetic activity Some findings on the applications of ERTS and Skytab , p0027 A75-11975 A remote sensing study of Pacific hurricane Ava p0012 A75-17112 imagery for metropolitan land use analysis MORGAN, J RGAN, J. ERTS applications in Thailand - A progress report p0076 A75-17123 p0013 A75-17136 MCGINNIES, W. G. A study to explore the use of orbital remote sensing to determine native arid plant distribution MILES. R. The economic value of remote ser p0006 N75-10544 from space: An ERTS overview and the value of continuity Some first accuracy estimates for applications of aerial [E74-10818] Volume 8: Atmosphere MCGINNIS, D. F. [NASA-CR-141191] [AD-786418] Mapping of the 1973 Mississippi River floods by the p0080 N75-14213 p0032 N75-14335 MORLEY, L. W. The Canadian ERTS program NOAA-2 satellite p0028 A75-17139 IILLARD, J. J. MCGOOGAN, J. T. p0077 N75-10550 Planning applications in east central Florida Skylab altimeter applications and scientific results MORRISON, R. B. [E75-10015] p0021 N75-13337 [AIAA PAPER 74-1221] Skylab radar altimeter - Short-wavele p0063 A75-12035 Application of Skylab EREP photographs to study of the MILLER, J. modern episode of accelerated erosion in southern length perturbations p0066 A75-16096 Geophysical monitoring for climatic change. The NOA detected in ocean surface profiles p0018 N75-11510 program [E75-10054] MCGREGOR, D. N. p0036 N75-13354 MILLER, J. M. Applications of Skylab EREP photographs to mapping A Shuttle/SpaceLab RF environment survey facility A summary of ERTS data applications in Alaska p0016 A75-19457 of la ndforms and environmental geology in the Great Plains p0077 A75-17238 and Midwe MCLAURIN, J. D. Environmental surveys in Alaska based upon ERTS sta p0017 N75-10551 e Skylab S-190B Earth Terrain Camera [F75-10055] o0037 N75-13355 p0069 A75-18071 Cartographic evaluation of Skylab-A S-192 scanner data MILLER, J. R. Geologic interpretation of infrared thermal images in East A motion-compensated spatial scanner area, Western Desert, Egypt p0033 A75-17215 images [E75-100]. p0068 A75-17213 p0031 N75-14197 MOURAD, A. G. MCNAUGHTON, J. MILLER, L. S. Interaction of marine geodesy, satellite technology and Optical data processing analysis of stream patterns exhibited on ERTS-1 imagery p0056 A75-17219 neter applications and scientific results ocean physics [NASA-CR-137470]

[AIAA PAPER 74-1221]

p0063 A75-12035

o0043 N75-10671

MOWER, R. D. PERSONAL AUTHOR INDEX Calibration and evaluation of Skylab altimetry for geodetic NITHACK, J. PARASHAR, S. K. Investigation of radar discrimination of sea ice Geologic evaluation of central Italy from ERTS-1 and determination of the geoid p0038 N75-14238 [F75-10064] n0030 N75-13364 Skylab data p0042 A75-17121 NOEL. J. An application of infrared remote sensing techniques to A preliminary feasibility study on a synthetic aperture radar /SARSAT/ for earth resources surveys On the feasibility of remote monitoring of soil moisture ecological problems [AD-786028] microwave sensors p0022 N75-13419 D0067 A75-17194 PARKER, R. A. R. MUELLER, I. I. Planning and management of science programs on NORRIS. D. R. Earth parameters from global satellite triangulation and Satellite photography of eddies in the gulf loop [AIAA PAPER 74-1215] trilateration p0064 A75-12985 DO075 A75-12030 p0041 A75-11255 PARMENTER, F. C.

Observing and forecasting local effects from satellite MUNDAY, J. C., JR. Lake Ontario water mass delineation from ERTS-1 Flood inundation in the southeastern United States from p0016 A75-18153 p0042 A75-17184 aircraft and satellite imagery p0047 A75-17138 NOSWORTHY, C. T. Estimation of population density in Tokyo districts from Preliminary results of fisheries investigation associated pi0003 A75-17167 Feasibility study ASCS remote sensing/compliance p0076 A75-17103 FRTS-1 data p0003 A75-17167 determination system PATERSON, M. P. NASA-CR-1342881 p0007 N75-12419 Quality control in global and regional air chemistry monitoring networks: Lessons of the European Air Chemistry Network p0018 N75-11540 Crude and refined netroleum oil structured luminescence signatures induced by UV laser or lamp and their re p0033 A75-17207 ensing applications 0 MURARKA, R. S. PATTMAN, J. D. R. An extension of the combinatorial geometry technique An examination of whiteouts and transmission through ice fog in the visible and near infrared [DREO-TN-74-18] for modeling vegetation and terrain features OBRIEN, J. J. [AD-782883] MURPHREE, D. L. p0006 N75-10577 p0021 N75-12467 The Coho Project: Living resources prediction feasibility PAUL, C. K. The availability of local aerial photography in southern California Volume 2: Environmental report Mathematical modeling for the detection of fish by [PB-234058/6] p0044 N75-10682 p0065 A75-15953 The Coho Project: Living resources prediction feasibility udy. Volume 3: System evaluation report p0044 N75-10683 [NASA-CR-141314] p0025 N75-15126 MURPHY, L. P. Early results from the experimental Automated Ima Data Extraction System p0069 A75-180 [PB-234059/4] An electronic drawing aid for map compiling p0027 A75-10076 p0069 A75-18072 MUSICK, H. B. OGURA, H. A study to explore the use of orbital remote sensing to Performance of a hologram matrix radar PAVLOV, V. I. p0065 A75-14792 Space iconics [NASA-TT-F-798] determine native arid plant distribution [E74-10818] p0006 N75-10544 OGUROK, D. D. DO057 N75-11419 Optical-acoustic detection of low concentrations of MYERS, V. I. PEACOCK, K. Use of ERTS-1 imagery for land evaluation in Pennin hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation Ground-based determination of atmos correction of ERTS-1 data p County, South Dakota p0055 A75-17135

Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical p0054 A75-18476 p0063 A75-11443 PENNEWELL, J. D.

Applicability of Skylab remote sensing for detection and and thermal characteristics of plants and soil Measurements of meteorological parameters and air pollution spread over Stockholm, Sweden, by means of monitoring of surface mining activities p0006 N75-12394 [E75-10012] [E75-10038] 275-10038] p0044 N75-12407 Applicability of Skylab remote sensing for detection and Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil remote sensing and airborne instrumentation monitoring of surface mining activities [E75-10085] p0018 N75-11531 p0038 N75-15103 OMAKUPT, M. and thermal c [E75-10033] p0007 N75-12403 AKUPT, M.
ERTS applications in Thailand - A progress report
p0078 A75-17123 PENNEY, M. E. Develop techniques and procedures, using multispectral Remote sensing applications in marine science programs systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil VIMS [NASA-CR-137475] p0043 N75-10566 Application of ERTS data to environmental inventory and p0008 N75-14196 [E75-10077] PERRY, L p0016 A75-18069 Water depth penetration film test p0047 A75-12873 ONUOHA, D. PERSSON C N An extension of the combinatorial geometry technique Measurements of meteorological parameters and air for modeling vegetation and terrain features pollution spread over Stockholm, Sweden, by means of remote sensing and airborne instrumentation [AD-782883] p0006 N75-10577 NALEPKA, R. F. p0018 N75-11531 ORHAUG, T. Developing processing techniques for Skylab data [E74-10799] p0057 N75-Processing of ERTS CCT imagery of Swe p0057 N75-10527 PETERSEN, G. W. of Sweden. A pilot p0060 N75-14219 A method of specifying remotely sensed units for soil mple points p0002 A75-17106
The use of remote sensing and natural indicators to study Developing processing techniques [E75-10040] for Skylab data p0058 N75-12409 ORLOV, V. M. sample points -Spectrophotometric observations of earth from Developing processing techniques for Skylab data 'Soyuz-13'
[IAF PAPER 74-156]

condication of spectrometric delineate floodplains - Preliminary findings [E75-10082] p0061 N75-15102 p0085 A75-13765 p0087 A75-17142 NAUMANN, R. J. The application of spectrometric and potechniques for remote sensing of oil on sea water PETERSEN, R. and polarization Skylab induced environment Applications of remote sensing by the Nebraska state [AIAA PAPER 74-1225] p0011 A75-12037 p0069 A75-17232 p0068 A75-17210 NEECE, R. N. An overview of Texas activities in remote sensing ERTS-1 images in glaciology and hydrology: Norweg The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 5: Inland water resources [NASA-CR-141269] p0080 N75-14209 - p0076 A75-17235 p0051 N75-14233 NELSON, E. E. Interactive computer processing for land use plan . p0080 N75-14209 Correlation of coastal water turbidity and current p0054 A75-17134 NEMKOVSKIY, B. L. circulation with ERTS-1 and Skylab imagery p0048 A75-17181 ETTINGER, L R. · A comparison of Skylab and ERTS data for agricultural System for electronic transformation and geographic Comparison of Skyled end Error School Corp and natural vegetation interpretation [AIAA PAPER 74-1219] p0001 A75-12033
Application of ERTS-1 data to the protection and correlation of satellite television information Measurements of meteorological parameters and air p0071 N75-11434 NEPOKLONOV, B. N. pollution spread over Stockholm, Sweden by means of remote sensing and airborne instrumentation management of New Jersey's coastal en Geographical correlation of TV and IR images obtained from weather satellites p0071 N75-11433 p0018 N75-11531 p0042 A75-17233 p0071 N75-11433 OWEN-JONES, E. S.
The interpretation and use of false-colour infra-red and PETTYJOHN, R. L Signatures of various earth surfaces measurables 5 NEUSTADTER, H. E. On evaluating compliance with air pollution levels 'not to be exceeded more than once a year' true colour photography of part of Argentina obtained by Skylark earth resources rockets p0067 A75-17192 Nimbus-5 microwave spectrometer p0066.A75-17119 ETTYJOHN, W. A. p0011 A75-12803 The application of pattern crassings of the state of the The application of pattern classification techniques to Automatic mapping of strip mine operations from NEWTON, R. W. p0034 A75-18065 spacecraft data On the feasibility of remote monitoring of soil moisture Automated strip mine and reclamation mapping from p0067 A75-17147 Recognition and interpretation of spectral signatures of p0035 N75-10556 FOTS NEZ. G. regetation from aircraft and satellite imagery in Western Queensland, Australia p0008 N75-14241 Land-use planning aided by computer PETZEL, G. An evaluation of the suitability of ERTS data f modelling/mapping system to combine remote sensing, natural resources, social and economic data OWEN. W. P. p0035 N75-10553 numoses of petroleum exploration Ocean color, a thre [NASA-CR-139145] a three component system p0076 A75-17118 PHILLIPS, E. p0044 N75-11411 Evaluation of Skylab imagery as an information service

NGUYEN, V. K.

Performance of a hologram matrix radar p0065 A75-14792

NIAVRO, B. P.

Development of research in the USSR on the use of radar images for geological purposes p0033 A75-17126

NICHOL, J. E. Land type analysis for regional land use planning from photomorphic mapping - An example for B p0013 A75-17137 Colorado

NIEMANN, B. J., JR. The application of ERTS-derived information to the regional land use planning process p0015 A75-18058

NISHIOKA. K. Systems study of an automated fire weather data p0008 N75-12277 --[NASA-TM-X-62402]

PACK, D. H. Geophysical monitoring for climatic change. The NOAA rogram p0018 N75-11510 PAINTING, D. J.

p

Electronics in meteorological instrumentation p0065 A75-14726

PALGEN, J. O. Optical processing of ERTS-1 (MSS) images: Examples for the Arcachon basin and Rhone delta DO060 N75-14223

PALMASON, G. Environmental studies of Iceland with ERTS-1 imagery p0012 A75-17105

PICKERING, S. M. Geologic evaluation and applications of ERTS-1 im over Georgia PIECH, K. R.

PHILPOT, W.

p0034 N75-10552 Special color analysis techniques p0084 A75-12874

Coastal and estuarine studies with ERTS-1 and Skylab

Correlation of coastal water turbidity and current

Correlation of coastal water turbers, circulation with ERTS-1 and Skylab imagery p0048 A75-17181

Meteorological interpretation of high resolution imag

for remote sensing of earth resources p0060 N75-14229

resources p0056 N75-10526

p0047 A75-18240

for investigating land use and natural re [E74-10730] p

PERSONAL AUTHOR INDEX PIERSON, W. J., JR. PRILEPIN, M. T. Automated strip mine and reclamation mapping from Preliminary analysis of Skylab RADSCAT results over Optical quantum generators in geodetic measurem p0035 N75-10556 [AD-787281] p0032 N75-15237 REKDER J R PRINCE, R. E. [NASA-CR-141454] p0046 N75-14793 Geographic applications of ERTS-1 imagery to landscape PILLARS, W. W. Skylab S-192 ratio codes of soil, mineral, and rock spectra A Shuttle/SpaceLab RF environment survey facility p0016 A75-19457 [E74-10810] p0034 N75-10537 PRITCHARD, J. A. for ratio image selection and interpretation REICHLE, H. Mapping of the 1973 Mississippi River floods by th NOAA-2 satellite p0028 A75-1713 p0003 A75-17158 Global measurements of air pollution from satellites p0028 A75-17139 p0018 N75-11521 Extensive summe: upwelling on Lake Michigan during The application of Skylab EREP data for land use REID, I. A. 1973 observed by NOAA-2 and ERTS-1 satellites classification and mapping in the Cleveland and Columbus Water survey of Canada: Application for use of ERTS-A p0048 A75-17160 for retransmission of water resources date areas PROST G I [£75-10083] n0023 N75-14201 Index of Skylab data available at Colorado School of p0049 N75-11409 PINCURA, P. G. Water survey of Canada: Application for use of ERTS-A or retransmission of water resources data Mines Multidisciplinary applications of ERTS and Skylab data [E75-10041] p0072 N75-12410 p0015 A75-17234 in Ohio PRUITT, G. R. [E75-10026] p0050 N75-13342 PINSON, J. W. Water survey of Canada: Application for use of ERTS-A An orbiting visible/infrared spectrometer for terrestrial atmospheric and oceanographic applications Correlation of multispectral imagery with water analysis retransmission of water resources data Sensing Project p0014 A75-17165 - The Ross Barnett Reservoir Remote p0068 A75-17212 p0050 N75-13353 PUTNAM M REINING, P. Earth Resources Technology Satellite-1 (ERTS-1) data and anthropology: Use of these data in carrying capacity estimates for sites in Upper Volta and Niger [E75-10001] p0078 N75-11406 PIPES, F. J. The economic value of remote sensing of earth resources Skylab environmental and crew micro from space: An ERTS overview and the value of continuity p0024 N75-14366 of service. Volume 5: Inland water resource [NASA-CR-141269] p0080 p0080 N75-14209 PIRIE, D. M. California nearshore processes - ERTS 1 Utilization of ERTS-1 imagery in cultivation and p0048 A75-17179 settlement site identification and carrying capacity estimates PITTS, D. E.
The effect of atmospheric water vapor in Upper Volta and Niger [E75-10029] DOOO8 N75-13345 p0013 A75-17131 classification of ERTS data QUADE, J. G. REINISCH, R. F. The Great Basin investigation PITTS, J. N., JR. Airborne survey of major air basins in California Celifornia [E75-10076] DO037 N75-14195 p0011 A75-14360 Airborne survey of major air basins p0011 A75-14360 The Great Basin investigation REY, P. A [E75-10102] n0038 N75-15120 PIYAPONGSE, P. Optical processing of ERTS-1 (MSS) images: Examples ERTS applications in Thailand - A progress report for the Arcachon basin and Rhone delta p0076 A75-17123 Monitoring and predicting El Nino invasions p0060 N75-14223 p0041 A75-12719 ARNICA/ERTS-1 experiment. Results and p0081 N75-14246 PLUHOWSKI, E. J. Dynamics of suspended sediment plumes in Lake Ontario RIABOV, E. A R [E74-10815] p0049 N75-10541 Optical-acoustic detection of low concentrations of POHN. H. A. hydrogen fluoride, nitrous oxide, and carbon dioxide in gases hydrogen fluoride, nitrous oxide, and cardon dioxide in gen-using pulsed hydrogen-fluoride laser radiation p0063 A75-11443 RAAD, J. H. ND, J. H. Resource analysis applications in Michigan p0069 A75-17231 nsing geophysics from Skylal [E75-10045] p0030 N75-12413 Remote se [E75-10051] RICHARDS, D. A. ng geophysics from RADIONOV, B. N. p0036 N75-13351 Cruise data report. CHAIN 115 leg 2, Southlant Space iconics [NASA-TT-F-798] expedition [AD-784421] ng geophysics from Skylat DO057 N75-11419 p0030 N75-13352 p0029 N75-10641 [F75-10052] RAGUSA, J. M.
NASA-KSC's earth resources benefits from space RICHARDSON, A. J. ng geophysics from Skylab Remote sens p0037 N75-13357 [E75-10057] Land use classification and ground truth correlations from exploration p0075 A75-13592 imultaneously acquired aircraft and ERTS-1 MSS da Remote sensing geophysics from Skylab RAHN, F. H. p0037 N75-13358 p0014 A75-17188 [F75-10058] An extension of the combinatorial geometry technique for modeling vegetation and terrain features [AD-782883] p0006 N75-10577 The application of remote sensing techniques: Technical Environmental studies of Iceland with ERTS-1 imager nd methodological issues p0012 A75-17105 RAHN, P. H. [NASA-CR-140674] p0070 N75-10564 RIVEREAU, J. C. Use of ERTS-1 imagery for land evaluation in Penningto County, South Dakota p0055 A75-1713 Comparison of various numerical techniques for classification of multispectral data. Application to geology p0038 N75-14244 A Skylab program for the International Hydrological p0055 A75-17135 cade (IHD) p0050 N75-12408 RAMSEIER, R. O. [E75-10039] ROBBIANO, P. Skylab water depth determination A study of microwave emission properties of sea ice Airborne vapor surveillance system [AD-785585] 10048] p0041 A75-17120 [E75p0050 N75-13348 AIDJEX 1972 A Skylab program for the International Hydrological lecade (IHD) RAO. M. M. n0021 N75-12459 Geophysical variations at Sodankyla during a geomagnetic storm, 17-18 December 1971 ROBINETTE, R. p0052 N75-15111 [E75-10093] Application of remote sensing to state and regional p0030 N75-13432 A Skylab program for the International Hydrological [REPT-9] oroblems [NASA-CR-140805] p0078 N75-11414 RAO, P. K. n0052 N75-15112 [F75-10094] Images from the NOAA-3 very high resolution radiometer RORINSON J. G. Resource analysis applications in Michigan p0069 A75-17231 over the North Sea and adjoining countries p0066 A75-16188 Microwave emission characteristics of moist soils p0001 A75-15180 RODGERS, C. D. RATTI. B. POPOVA, T. P. p0072 N75-11562 Infrared remote sounding Digital land-use classification using an ERTS-1 image p0023 N75-14226 Space iconics [NASA-TT-F-798] RODIONOV. B. N. Geographical correlation of TV and IR images obtained om weather satellites p0071 N75-11433 p0057 N75-11419 RAWSON, R. PORFIREV. L. F. from weather satellites Four channel simultaneous X-L band imaging SAR pdar p0054 A75-17116 Analytical estimates of precision in determining parameters of motion of artificial earth satellite when System for electronic transformation and geographic correlation of satellite television informati RAY, G. C. p0071 N75-11434 analyzing imprecise measurements with correlated errors p0053 A75-11419 BESMEX: Bering Sea marine mammal experiment [NASA-TM-X-62399] p0008 N75-13503 ROGERS, R. Skylab and ERTS-1 investigations of coastal land-use POTTER, A. E. Summary of flight performance of the Skylab Earth Resources Experiment Package / EREP/ REAL R. R. and water properties in Delaware Bay [AIAA PAPER 74-1220] MAA PAPER 74-1220] p0041 A75-12034 Coastal and estuarina studies with ERTS-1 and Skylab An electronic drawing aid for map cor p0027 A75-10076 00076 A75-17211 p0047 A75-16240 Correlation of coastal water turbidity and current POTTER. J REBEL, D. L. Effect of atmospheric haze and sun angle on automatic A Skylab program for the International Hydrological p0014 A75-17157 classification of ERTS-1 data Decade (IHD) circulation with ERTS-1 and Skylab imager p0048 A75-17181 POULTON, C. E. [E75-10093] p0052 N75-15111 A comparison of Skylab and ERTS data for agricultural Inventories of Delaware's coastal vegetation and land-use A Skylab program for the International Hydrological utilizing digital processing of ERTS-1 imagery p0004 A75-17186 crop and natural vegetation interpretation Decade (IHD) p0001 A75-12033 [E75-10094] p0052 N75-15112 POUQUET, J. ROGERS, R. H. REED, J. C., JR. Earth sciences in the age of the satellis Automatic classification of eutrophication of inland lake computer cellular Land-use planning aided by p0070 A75-18693
Optical processing of ERTS-1 (MSS) images: Examples from spacecraft data p0048 A75-17164 modelling/mapping system to combine remote sensing, Automatic mapping of strip mine operations from acceraft data p0034 A75-18065 natural res sources, social and economic data for the Arcachon basin and Rhone delta p0076 A75-17118 spacecraft data p0060 N75-14223 Automated strip mine and reclam poods N75-10556 POZDIAKOV, D. V. Coastal and estuarine studies with ERTS-1 and Skylab p0047 A75-16240 The application of spectrometric and polarization techniques for remote sensing of oil on sea water ROHDE, W. G. Forest defoliation assessment with satellite imagery p0003 A75-17171 p0068 A75-17210 Inventories of Delaware's coastal vegetation and land-use utilizing digital processing of ERTS-1 imagery p0004 A75-17186 PRABHAKARA, C. ROLLER, N. E. G. Estimation of sea surface temperature from remote sensing in the 11to 13-micron window region Color terrain map of Yellowstone computer-derived from ERTS MSS data vstone National Park, p0065 A75-14803 p0004 A75-17185 Automatic classification of eutrophication of inland lakes

p0048 A75-17164

Automatic mapping of strip mine operations from pacecraft data p0034 A75-18065

Remote sensing to detect the toxic effects of metals

on vegetation for mineral exploration p0034 A75-17226

ROMANOVSKII, IU. A.

Morning-evening asymmetry of the thermosphere during a geomagnetic disturbance p0011 A75-1110:

p0011 A75-11103

p0030 N75-13359

ROOT, R. R.

Color terrain map of Yellowstone National Park, computer-derived from ERTS MSS data n0004 A75-17185

ROSENBERG, N.

Applications of remote sensing by the state of Nebraska p0069 A75-17232 ROSS, D.

A remote sensing study of Pacific h p0012 A75-17112 Multi-frequency radiometric measure

ents of foam and a mono-molecular slick ROUSE, J. W., JR. p0068 A75-17208

On the feasibility of remote monitoring of soil moisture vith microwave sensors p0067 A75-17147 Remote measurements of water pollution with a lidar p0014 A75-17205 oolarimete: ROWELL E. M.

Soviet satellite geodesy: Basis and status of Soviet laser geodesy and long-base line interferometry
[AD-781138/3] p0029 N75-10524

ROZANOV, V. V.

The potentialities for determining content of minor gas constituents in the atmosphere by satellites

p0015 A75-17932 RUCK, G. T.

Bistatic radar sea state monitoring

[NASA-CR-137469] p0043 N75-10673 RÚML, D. J.

Investigation of land use of northern megalopolis using ERTS-1 imagery [NASA-CR-2459]

n0017 N75-10545 RUMMEL R.

Use of terrestrial gravity data in combination with satellite results for determining geoid heights in Central Europe p0029 A75-18128

RUSSELL J. M., III

Measurement of trace stratospheric gases by inverting thermal infrared limb radiance profiles p0018 N75-11523

RUSSELL, P. B.

Lidar observations of atmospheric particulate content p0019 N75-11559

RYLAND, b. W. Evaluation of ERTS-1 and aircraft data for assessing internal drainage in irrigated agriculture

p0002 A75-17143

S

SABINS, F. F., JR.

Oil exploration needs for digital processing of imager p0033 A75-10437

SACCHI, R CHI, R.
Geological features on ERTS-1 images of a test area in
west-central Albs p0037 N75-14237 the west-central Alps

SADOWSKI, F. G. The remote identification of terrain features and materials

at Pennsylvania test sites: An investigative study of techniques

[PB-235992/5] p0031 N75-14263

SAEMUNDSSON, K

Environmental studies of Iceland with ERTS-1 image p0012 A75-17105

Mapping of water bodies in northern Germany from ERTS p0051 N75-14227

SALISBURY J W

Remote sensing of rock type in the visible and pools A75-17220 near-infrared

SALMON, B.

Surface compositional mapping in the Wind River Range and Basin, Wyoming by multispectral techniques applied to ERTS-1 data p0028 A75-17224

SALOMONSON, V. V. Water resources

n0043 N75-10559 SAND, F.

The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. Volume 2: Source document [NASA-CR-141193] p0079 N75-14204

SANDS, K. J. T.

Electronics in meteorological instrumentation DO065 A75-14726

SARKISIAN, S. A.

Economic effectiveness of using the space systems of distant indication of natural resources
[IAF PAPER 74-165] p0075 A75-13770

SATTINGER, I. J.

Resource analysis applications in Michigan

p0069 A75-17231 Study of recreational land and open space using Skylab

[E75-10020] p0020 N75-12399 Study of recreational land and open space using Skylab [E75-10034] p0020 N75-12404

Study of recreational land and open space using Skylab imagery [E75-10092]

p0024 N75-15110

SAUER, G. E.

Water depth penetration film test p0047 A75-12873

SAVASTANO, K.

Preliminary results of fisheries investigation associate with Skylab-3 p0003 A75-17167

SAVIGEAR, R. A. G.

The interpretation and use of false-colour infra-red and true colour photography of part of Argentina obtained by Skylark earth resources rockets p0067 A75-17192

A comparison of the capabilities of ERTS-1 and the Skylark earth-resources rocket for resources surveying in p0023 N75-14250 SAXON, D. S.

Lectures on the scattering of light p0019 N75-11556 SCANVIC, J. Y.

ERTS image contribution to geological knowledge covering different types of climates (France, Togo, Dahomey, p0038 N75-14240 Madagascar)

SCARPACE, F. L p0014 A75-17162 Scanning thermal plumes

Monitoring trophic status of inland lakes - A quantitative application of ERTS imagery p0049 A75-18067 SCHAFRER G.

Deteriorating effects on 3 mm wave passive imager p0055 A75-17198

SCHALL, M. A.
SICLOPS - A system of computer progra mapping of airborne scanner imagery p0055 A75-17156 SCHANDA, E.

On the detectability of atmospheric carbon monoxide by p0013 A75-17129 microwave remote sensing Emissivities and forward scattering of natural and man-made material at three millimeter wavelength

Deteriorating effects on 3 mm wave passive imagen p0055 A75-17198

DO067 A75-17197

SCHAR, S. W.

Resource analysis applications in Michigan p0069 A75-17231

SCHEGLOV, S. N.

Australian and Soviet gravity surveys along the Australian calibration line

p0031 N75-13454 [BULL-161]

SCHILLING, P. E.

Optical film density values from color IR photography for wetland soils mapping p0005 A75-18073 SCHMER, F. A.

ation of ERTS-1 and aircraft data for assessing internal drainage in irrigated agriculture

n0002 A75-17143

SCHMITT, H. H.

The progeny of Skylab - The pioneering phase of our expansion into space [AIAA PAPER 74-1216] p0063 A75-12031

SCHMUGGE, T. J.

A radio picture of the earth n0066 A75-16191 SCHNEIDER, W. C. p0075 A75-13618

Skylab contributions to the future SCHOENFELD. M. p0031 N75-14239

ERTS-1 data on Afar tectonics SCHUBERT, J. Coastal wetlands analysis from ERTS MSS digital

and field spectral measurements n0048 A75-17178 A demonstration of ERTS-1 analog and digital techniques applied to strip mining in Maryland and West Virginia [NASA-TM-X-70798] p0039 N75-15133

SCORER, R. S. Quality control in global and regional air chemistry monitoring networks: Lessons of the European Air p0018 N75-11540 Chemistry Network SEEVERS, P. M.

Satellite pictures can help guide Sandhills e Sandhills irrigation p0021 N75-13339 [E75-10023]

Use of ERTS-1 imagery to interpret wind-erosion hazard in the Sandhills of Nebraska [F75-10044] n0059 N75-13347

SEIDEL, A. D.

The economic value of remote sensing of earth resources The economic value of remote sensing or earth resources from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources, agriculture. Part 3: The integrated impact of improved (ERS) information on US agricultural commodities [NSAS-CR-141262] p0079 N75-14207

SEIDEL K. Methodological aspects and regional examples of snow-cover mapping from ERTS-1 and EREP imagery of

p0051 N75-14232 the Swiss Alps SEMENOV, R. N.

The wind vector in a layer over water

p0041 A75-15179

SERGEEV. V. A. On the longitudinal localization of the substorm active

region and its changes during the substp0027 A75-11861

Planning and management of science programs on Skylab [AIAA PAPER 74-1215] p0075 A75-12030

SHAH. N.

Correlation of coastal water turbidity and current circulation with ERTS-1 and Skylab imagery p0048:A75-17181

Automatic classification of eutrophication of inland lakes from spacecraft data p0048 A75-17164 SHAIDUROV, V. O.

Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases ing pulsed hydrogen-fluoride laser radiation

p0063 A75-11443

SHAKLEE R V.

Land use mapping using ERTS-1 imagery for The Four Rivers Resource Conservation and Developnent District in p0016 A75-18070

SHAPIRO, A. Terrain properties and topography from Skylab altimetry [£75-10059]

SHARKOV, E. A. Microwave emission characteristics of p0001 A75-15180

SHCHUKIN, IU. K. Plutonic structure of the earth's crust on space images p0027 A75-17125

SHEFFIELD, L

Applications of remote sensing by the state of Nebraska p0069 A75-17232

SHEFFIELD, L F. Satellite pictures can help guide Sandbills irrigation [E75-10023] p0021 N75-13339

SHEIVES, T. C. Remote measurements of water pollution with a lidar plarimeter p0014 A75-17205 A study of a dual polarization laser backscatter system for remote identification and measurement of water

pollution [NASA-CR-140665] p0017 N75-10563

HELTON, M.

Effect of atmospheric haze and sun angle on automatic p0014 A75-17157 classification of ERTS-1 data SHELTON, R. L.

Design concepts for land use and natural resource inventories and information systems p0013 A75-17133
SHENK, W. E.

The detection of dust storms over land and water with

satellite visible and infrared measurements p0012 A75-16675

SHEPHERD, J. A study of the usefulness of Skylab EREP data for earth

resources studies in Australia p0036 N75-12415 SHILIN, B. V.

Geological and geophysical studies from space and rospects for their development p0035 N75-11436 SHORT, N. M.

Mineral resources, geological structures, and landform surveys p0035 N75-10561 Exploration for fossil and nuclear fuels from orbital

[NASA-TM-X-70781] p0017 N75-11413

SHUCHMAN, R. A.

Feasibility of using multiplexed SLAR imagery for water resources management and mapping vegetation mapping vegetation p0054 A75-17115

communities SHUTKO, A. M.

> Passive microwave sensing of moist soils p0066 A75-17124

Preliminary data on the UHF radiometric research of soil wetness, conducted in the USSR, in 1974
[NASA-TT-F-16151] p00 p0009 N75-15097

SIDIK, S. M. On evaluating compliance with air pollution levels 'not to be exceeded more than once a year

SIGNORINI, C.

Solar wind and geomagnetic activity

p0027 A75-11975

p0073 N75-15113

p0011 A75-12803

SIGTRYGGSSON, H. Environmental studies of Iceland with ERTS-1 imagery p0012 A75-17105

Statistical separability of agricultural cover types in subsets of one to twelve spectral channels p0005 A75-17216

Spectroradiometric and multispectral scanner over Lake [E75-10066] p0072 N75-13366

Reformatting Skylab and MSS data over northern [E75-10016] p0061 N75-15101

Analysis of Skylab MSS data, and calibration of thermal data over Lake Monroe, Indiana [E75-10095]

SIMPSON, R. B.

Investigation of land use of northern megalopolis using p0017 N75-10545

ERTS-1 imagery [NASA-CR-2459]

SIRY, J. W. Satellite techniques for the study of secular variations in position p0064 A75-12980

SITALOV. A. E.

Scientific research and experimental design planning and coordination for 1973 [AD-785998] p0081 N75-14259

SKALEY, J.

Evaluation of Skylab imagery as an information service for investigating land use and natural resources
[E74-10730] p0056 N o0056 N75-10526

Magnetopause SPENCER, M. M.

SPENCER, W. E.

use rotational forms

of the earth - The Skylab radscat experiment

SICLOPS - A system of computer programs for rectified

us active and passive microwave response

mapping of airborne scanner imagery p0055 A75-17158

p0011 A75-11221

p0012 A75-17114

SUVARNASUDHI, K.

SWEET, D. C.

in Ohio

ERTS applications in Thailand - A progress report p0076 A75-17123

Multidisciplinary applications of ERTS and Skylab data

p0015 A75-17234

The application of Skylab EREP data for land use SKATCHKOV, V. The potential benefits for development derived from space applications STAELIN, D. H. Signatures of various earth surfaces measured by the Nimbus-5 microwave spectrometer p0066 A75-17119 classification and mapping in the Cleveland and Columbus [IAE BAPER 74.147] n0075 A75-13761 STANCZUK D T [F75-10083] n0023 N75-14201 of ERTS-1 data to the protection and SKENDERGEF. C. SWIFT, C. T. A preliminary feasibility study on a synthetic aperture radar /SARSAT/ for earth resources surveys management of New Jersey's coastal em A simulation of synthetic aperture radar imaging of ocean p0042 A75-17233 surveys p0067 A75-17194 waves [NASA-TM-X-72629] p0044 N75-12172 Development of research in the USSR on the use of SWINNERTON, J. R. SKORFLEV. S. F. radar images for geological purposes p0033 A75-17126 Geometronics-earth resource measurements and displays futonic structure of the earth's crust on space images STEELE, M. L p0027 A75-17125 for the Forest Service p0005 A75-18063 Correlation of multispectral imagery with water analysis Correlation of multispectral Imagery with water of the Ross Barnett Reservoir Remote Sensing Project p0014 A75-17165 The availability of local aerial photography in southern California SLEDGE, B. SYDOR, M. Remote sensing of western Lake Superior p0048 A75-17161 [NASA-CR-141314] p0025 N75-15126 Theoretical investigations on the determination of the SLOAN, R.
Detection of moisture and moisture related phenomena vertical temperature distribution in the atmosphere assuming multispectral thermal radiation measurements at T from Skylab [E75-10007] 60 GHz p0007 N75-13333 n0024 N75-14331 [BMFT-FB-W-74-04] TALMON, T. L. SMAIL H. E. STEINBERG, H. A. Land use mapping using ERTS-1 imagery for The Four Rivers Resource Conservation and Development District in Multidisciplinary applications of ERTS and Skylab data An extension of the combinatorial recometry technique p0015 A75-17234 in Ohio for modeling vegetation and terrain feature
[AD-782883] p00 p0016 A75-18070 The application of Skylab EREP data for land use p0006 N75-10577 TANAKA, M classification and mapping in the Cleveland and Columbus STELLER, D. D. Radiative transfer of visible radiation California nearshore processes - ERTS 1 areas p0019 N75-11552 atmosphere [E75-10083] p0048 A75-17179 n0023 N75-14201 TAYLOR, C. D. STEPHAN, J. G. SMEDES, H. W. Mathematical modeling for the detection of fish by an Land-use planning aided by computer cellular modelling/mapping system to combine remote sensing. computer cellular Multidisciplinary applications of ERTS and Skylab data p0015 A75-17234 airborne laser p0065 A75-15953 in Ohio TAYLOR, R. E.
A Shuttle/SpaceLab RF environment survey facility The application of Skylab EREP data for land use natural resources, social and economic data p0076 A75-17118 classification and mapping in the Cleveland and Columbus p0016 A75-19457 Color terrain map of Yellowstone National Park, computer-derived from ERTS MSS data areas [E75-10083] TENGSTROM, E. p0023 N75-14201 New methods for studying the dynamics of the entire 00004 A75-17185 STERCKX, J. D0016 A75-18108 earth SMIRNOV, A. V. Preliminary evaluation of ERTS-1 imagery of Central frica p0050 N75-14242 Africa THAMAN, K. H.

Operational use of satellite and high altitude remote Analysis of methods for the geodetic support of complex ea geophysical surveys STEVENSON, P. sensing for the generation of input data for water demand models p0067 A75-17148 p0031 N75-13498 The economic value of remote sensing by satellite: An SMIRNOV, V. V. ERTS overview and the value of continuity of service. determining Analytical estimates of precision in determining parameters of motion of artificial earth satellite when Volume 2: Source document [NASA-CR-141193] THAMAN, R. R. Operational use of satellite and high altitude remote n0079 N75-14204 nalyzing imprecise measurements with correlated errors sensing for the generation of input data for water dema-The economic value of remote sensing of earth resources p0067 A75-17148 from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland The usefulness of imaging passive microwave for rural and urban terrain analysis p0055 A75-17199 SMITH D. E. of service. Volume [NASA-CR-141263] mulation of the San Andreas fault p0080 N75-14208 and urban terrain analysis DO033 A75-11068 STEVENSON, P. A. THOMAS, G. SMITH, D. V. Comparison of various numerical techniques for The economic value of remote sensing of earth resources Comparison of various numerical techniques for classification of multispectral data. Application to geology p0038 N75-14244 Automated image processing in forestry from space: An ERTS overview and the value of continuity of service. Volume 6: Land-use. Part 1: Introduction DO053 A75-10435 SMITH, F. and overview
[NASA-CR-141266] THOMAS, G. L. Four channel simultaneous X-L band imaging SAR p0054 A75-17116 Planning applications in east central Florida [E75-10015] n0021 n0080 N75-14210 p0021 N75-13337 STEVENSON, W. H. SMITH, G. A review of initial investigations to utilize ERTS-1 data THOMPSON D N Trace gas analysis of power plant plumes via aircraft in determining the availability and distrib Classification and mapping of coal refuse, vegetative cover types, and forest types by digital processing ERTS-1 measurement - O3, NOx, and SO2 che distribution of living p0043 N75-10555 emistry - p0064, A75-13148 marine resources DO054 A75-17110 Application of ERTS-1 data to the protection and ERTS-I data for classifying native plant THOMSON F. J. Central Colorado p0004 A75-17176 management of New Jersey's coastal em Study of atmospheric effects in Skylab data [E75-10006] environment p0042 A75-17233 p0020 N75-12391 SMITH, J. L. Mapping exposed silicate rock types and exposed ferric and ferrous compounds from a space platform Microwave profiling of snowpack free-water contr STOCKHOFF, E. H. p0049 A75-17950 Remote detection of soil surface moisture SMITH, P. G. p0003 A75-17146 [E75-10008] p0036 N75-12392 The remote identification of terrain features and materials Special color analysis techniques p0064 A75-12874 STOECKELER E. G. SMITH R M Multidisciplinary analysis for highway engineering at Pennsylvania test sites:. An investigative study of An analysis of applications development systems for numoses [PB-235992/5] [E75-10103] . p0025 N75-15121 60031 N75-14263 remotely sensed, multispectral data for Observations Division of the NASA Lyndon B. Johnson THORARINSSON. S. STORTZ K Space Center [NASA-CR-140345] Environmental studies of Iceland with ERTS-1 imagery Remote sensing of western Lake Superi p0071 N75-11447 pQO48 A75-17161 p0012.A75-17105 SMITH, V. E. THORLEY, G. STRATTON, R. H. Automatic classification of eutrophication of inland lakes Future US activities based on experience gained from Analysis of pseudocolor transformations of ERTS-1 images of Southern California area [E75-10037] p0036 N75-13346 spacecraft data p0048 A75-17164 and EREP THORSTEINSSON, I. FRTS-1 p0009 N75-14253 SMOKTY, O. I. Environmental studies of Iceland with ERTS-1 imagen Influence of the atmosphere on spectral radiance and contrasts of natural formations measured from space STREES, L V. p0012 A75-17105 Computer derived coastal water classifications p0019 N75-11557 TIMOSEEV III.M. spectral signatures p0042 A75-17177 The potentialities for determining content of minor gas constituents in the atmosphere by satellites SOBTI, A Simultaneous active and passive microwave response STRETEN, N. A. A satellite view of weather systems over the North of the earth - The Skylab radscat experiment p0015 A75-17932 p0012 A75-17114 American Arctic p0063 A75-10790 TINGEY, D. L Design data collection with Skylab/EREP microwave STRONG A F Quantitative determination of stratospheric serosol elling on Lake Michigan during instrument S-[E75-10070] n0059 N75-14189 1973 observed by NOAA-2 and ERTS-1 satellites n0016 N75-10535 [E74-10808] Design data collection with Skylab/EREP microwave p0048 A75-17160 Quantitative determination of stratospheric aerosol STRUEBING, K haracteristics p0072 N75-14199 [E75-10080] Use of ERTS-1 in sea-ice studies p0045 N75-14234 [E75-10010] p0020 N75-12393 SOLLERS, S. C.

The use of remote sensing and natural indicators to STUMPF, H. G. Quantitative determination of stratospheric aerosol The use or remote sensing and national delineate floodplains - Preliminary findings p0067 A75-17142 Extensive summer upwelling on Lake Michigan during characteristics [F75-10079] p0023 N75-14198 1973 observed by NOAA-2 and ERTS-1 satell p0048 A75-17160 INNEY, L. R. The application of ERTS-1 data to the Corps of Engineers Operational use of satellite and high altitude remote SUMMERS, R. A p0015 A75-18064 sensing for the generation of input data for water demand models p0067 A75-17148 Evaluation of ERTS data utilization in develo SOLOVYEV, V. I. p0055 A75-17193 countries Geographical correlation of earth cloud cover picture obtained from unoriented satellites p0071 N75-1143 SONNERUP, B. U. O. TONELLI, A. M. SUNG. C. p0071 N75-11435 Remote sensing techniques applied to the study of Italian Design data collection with Skylab/EREP microweve instrument S-193
[E75-10070] p0059 N75-14189 volcanic areas - The results of the repetition of the airborne

p0069 A75-17223

p0073 N75-14238

I.R. survey compared to the previous data

Contribution of space platforms to a ground and airborne

remote sensing programme over active Italian volcanoes [E75-10089] p0037 N75-14188

Contribution of space platforms to a ground and eirborne remote-sensing programme over active Italian volcanoes

TORLINE, R. J.

Land use classification and ground truth correlations from simultaneously acquired aircraft and ERTS-1 MSS da p0014 A75-17188

TOWNSHEND, J. R. G.

The interpretation and use of false-colour infra-red and true colour photography of part of Argentina obtained by Skylark earth resources rockets p0067 A75-17192 A comparison of the capabilities of ERTS-1 and the Skylark earth-resources rocket for resources p0023 N75-14250 ntral Argentina TRAFGAARDH C

Measurements of meteorological parameters and air pollution spread over Stockholm. Sweden, by means of remote sensing and airborne instrumentatio

ERTS-1 contributions to the ecogenetical knowledge of the Llanos of the Orinoco (Colombia and Venezuela) p0023 N75-14245

p0018 N75-11531

TRICHEL M. C.

Summary of flight performance of the Skylab Earth Resources Experiment Package /EREP/ n0076 A75-17211

Mapping of water bodies in northern Germany from ERTS p0051 N75-14227 TRIFONOV, V. G.

Plutonic structure of the earth's crust on space p0027 A75-17125
Elements of the deep structure of the earth's crust on the space images of the East Caucasus

p0028 A75-17228

ERTS-1 data on Afar tectonics TSUCHIYA, I. TSOMBOS, P.

onment of Tokyo by p0012 A75-17102

remote sensing TULLOS, J

The role of ERTS in the establishment and of a nationwide land cover information system

[NASA-CR-141045] p0020 N75-12425 TUMANOV, O. A.

Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gases using pulsed hydrogen-fluoride laser radiation p0063 A75-11443

TUNNEL, S. H.

Correlation of missions 191, 51M and helicopter photography [NASA-CR-134289] p0058 N75-12279 p0058 N75-12279

TURNER, A. K. Land-use planning aided by computer cellular

modelling/mapping system to combine remote sensing, natural resources, social and economic data p0076 A75-17118

Classification and mapping of coal refuse, vegetative cover types, and forest types by digital processing ERTS-1 p0054 A75-17110 TURNER. C.

Geophysical monitoring for climatic cha change. The NOAA p0018 N75-11510 program

TURNER, R. E. Determination of the earth's aerosol albedo using Skylab

data [E75-10096] p0025 N75-15114 TURUNEN, T.

Geophysical variations at Sodankyla during a omagnetic storm, 17-18 December 1971 n0030 N75-13432 [REPT-9]

U

UENK, D.

Investigation into the spectral signature of agricultural crops during their state of growth p0004 A75-17189
Radar cross sections of vegetation canopies determined by monostatic and bistatic scatterometry

p0005 A75-17190

ULABY, F. Optical data processing analysis of stream patterns exhibited on ERTS-1 imagen p0056 A75-17219

Active microwave measurement of soil water content p0001 A75-16242 [NASA-CR-141685]

Simultaneous active and passive microwave response of the earth - The Skylab radscat experi

p0012 A75-17114 p0005 A75-18556 Radar response to vegetation Design data collection with Skylab/EREP microwave instrument S-193

[E75-10070] DO059 N75-14189

ULARY, S. Design data collection with Skylab/EREP microwave instrument S-193 p0059 N75-14189 [E75-10070]

ULANOWICZ, R. E. -

A survey for the use of remote sensing in the Chesapeake Bay region [NASA-CR-137476] p0043 N75-10571

UTHE. E. E.

Lidar observations of atmospheric pa p0019 N75-11559

VAN KASTEREN, H.

Radar cross sections of vegetation canopies determined by monostatic and bistatic scatterometry p0005 A75-17190

VAN KASTEREN, H. W. J.

Investigation into the spectral signature of agricultural crops during their state of growth p0004 A75-17189 p0004 A75-17189 VAN KUILENBURG, J.

Radar cross sections or vegetation.

by monostatic and bistatic scatterometry

p0005 A75-17190

VANDEHULST, H. C.

Multiple scattering in cloud layers: some results p0019 N75-11554

VANGENDEREN .I.I.

An evaluation of stereoscopic viewing of ERTS and Skylab images p0060 N75-14220

VANMELLE, M.J.

Measurements of the dielectric properties of sea water at 1.43 GHz [NASA-CR-2458] p0045 N75-13496

VANROOY, D. L.

An analysis of applications development systems for remotely sensed, multispectral data for the Earth Observations Division of the NASA Lyndon B. Johnson Snace Center

[NASA-CR-140345]

VASILYEV, O. B.

Influence of the atmosphere on spectral radiance and contrasts of natural formations measured from space DO019 N75-11557

VERBEEK, W. J. P. A. [PHL-1974-23]

Pattern recognition relevant to reconnaissance and remote sensing

b0070 N75-10573

Investigation into the spectral signature of agricultural crops during their state of growth p0004 A75-17189

VEZIROGLU, T. N. Application of remote sensing for prediction and detection

of thermal pollution [NASA-CR-139182] p0017 N75-10572

Feasibility of remote sensing for detecting thermal pollution. Part 1: Feasibility study. Part 2: Implementation

DO025 N75-15199 [NASA-CR-134453]

VIBULSRESTH, S.

ERTS applications in Thailand - A progress report p0076 A75-17123

VIEZEE, W

Lidar observations of atmospheric particulate content p0019 N75-11559

VINCENT, R. K

Skylab S-192 ratio codes of soil, mineral, and rock spectra for ratio image selection and interpretation p0003 A75-17158

Surface compositional mapping in the Wind River Range and Basin. Wyoming by multispectral techniques to ERTS-1 data p0028 A75 techniques applied p0028 A75-17224 Unsupervised mapping of geologic features and soils

p0034 A75-17225 California New theoretical models and ratio imaging techniques associated with the NASA earth resources spectral

information system [NASA-CR-140357] DO058 N75-12417 The remote identification of terrain features and materials

at a California test site: An investigative study of IPB-235991/71 p0031 N75-14261

VINOGRADOV, A. B.

The calculation of cloud shadows in modelling of the earth's surface survey from space p0014 A75-17204 VINOGRADOV, B. V.
The calculation of cloud shadows in modelling

p0014 A75-17204 earth's surface survey from space Experience in integrated geological and geographical interpretation of space photographs obtained in the USSR p0036 N75-11437

Morphographic and morphometric analysis of vegetative cover photographs in aerial photography for the mapping of vegetation [AD-785932] p0009 N75-14258

VINOGRADOVA, A. I.

Geological and geophysical studies from space and prospects for their development p0035 N75-11436 VIRCHENKO, O. V.

Automated remote measurement stem in agricultural p0064 A75-12947 meteorology

Automated remote ng system in p0007 N75-12518 grometeorology

VIRDEN L water for a thirsty p0047 A75-13578 Can spinoffs from space provide world

VORONOV, S. A.

Detection of synchronous changes in fluxes of energetic electrons and gamma-quanta during a magnetic disturbance p0064 A75-12317 disturbance

WADDELL, B. H., JR.

The use of high altitude photography and ERTS-1 imagery for wildlife habitat inventory in Kansas

p0005 A75-18066

Monitoring trophic status of inland lakes p0049 A75-18067 application of ERTS imagery WAGNER, T. W.
The application of remote sensing techniques: Technical

and methodological issues [NASA-CR-140674] D0070 N75-10564 A Skylab program for the International Hydrological Decade (IHD)

[E75-10039] DO050 N75-12408 The remote identification of terrain features and materials Pennsylvania test sites: An investigative study of

techniques IPB-235992/5) p0031 N75-14263

WALKER, J. E. Special color analysis techniques p0064 A75-12874

WARK, D. A cloud physics investigation utilizing Skylab data [E75-10021]

p0020 N75-12400 A cloud physics investigation utilizing Skylab data [E75-10099] p0025 N75-15117

BESMEX: Bering Sea marine [NASA-TM-X-62399] mammal experimen p0008 N75-13503

WATERS, J. W. Signatures of various earth surfaces measured b imbus-5 microwave spectrometer p0066 A75-1 p0066 A75-17119

WATSON, K. onnaissance from quantitative analysis thermal infrared images p0056 A75-17218 Remote sensing geophysics from Skylab

[E75-10045] p0030 N75-12413 Remote sen ing geophysics from Skylab

[E75-10051] p0036 N75-13351 Remote sensing geophysics from Skylab

p0030 N75-13352 [E75-10052] Remote sensing geophysics from Skylab [E75-10057] p0037 N75-13357

Remote se ng geophysics from [E75-10058]

p0037 N75-13358 WATSON, R. D. Prediction of the Fraunhofer

line detectivity luminescent materials p0068 A75-17221 WEBER, F. P.

Inventory of forest and rangeland resources, including forest stress [E75-10032] p0007 N75-12402

WEBSTER, W. J., JR. A radio picture of the earth p0066 A75-16191

WEECKSTEEN, G.

ERTS image contribution to geological knowledge covering different types of climates (France, Togo, Dahomey. Madagascar) p0038 N75-14240

WEEDMARK, J. Performance of a hologram matrix radar

WELBY, C. W.

Utilization of EREP data in geological evaluation regional planning, forest management, and water management in North Carolina

p0065 A75-14792

[E75-10063] p0008 N75-13363 WELCH, R. I.

A comparison of Skylab and ERTS data for agricultural crop and natural vegetation interpretation p0001 A75-12033 [AIAA PAPER 74-1219]

WELLMAN P. Australian and Soviet gravity surveys along the Australian calibration line

[BULL-161] p0031 N75-13454 WELLS, J. An overview of Texas activities in remote sensing

p0076 A75-17235 WELLS, W. T. Skylab attimeter applications and scientific results [AIAA PAPER 74-1221] p0063 A75-

p0063 A75-12035 WENDEROTH, S. Multispectral photography for earth resources

p0074 N75-15138 [NASA-CR-141283] WERTZ, D. L. Correlation of multispectral imagery with water analysis

The Ross Barnett Reservoir Remote Sensing Project p0014 A75-17165

WESTIN, F. C. Use of ERTS-1 imagery for land evaluation in Penn County, South Dakota p0055 A75-17135

Correlation of coastal water turbidity and current

Correlation of coastal water turbun, circulation with ERTS-1 and Skylab imagery p0048 A75-17181 Monitoring physical and chemical parameters of Delaware Bay waters with an ERTS-1 data collection platform [E75-10090] p0051 N75-1 p0051 N75-15108

WETZLER E.

The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 5: Inland water resource [NASA-CR-141269] p0080 p0080 N75-14209 ERSONAL AUTHOR INDEX


PERSONAL AUTHUR INDEX	
WEZERNAK, C. T.	The Coho Project: Living resources prediction feasibilit
The use of remote sensing in timnological studies p0048 A75-17163	study. Volume 2: Environmental report [PB-234058/6] p0044 N75-1068
WHEELER, R. J.	The Coho Project: Living resources prediction feasibilit study. Volume 3: System evaluation report
Water resource and hazard planning report for the Clark Fork River Valley above Missoula, Missoula County,	[PB-234059/4] p0044 N75-1068
Montana [PB-236041/0] p0051 N75-14260	WORK, E. W., JR. Utility of ERTS for monitoring the breeding habit of
WHISLER, F.	migratory waterlowl p0006 N75-1055 WRIGHT, D. J.
Application of remote sensing to state and regional problems	The Coho Project: Living resources prediction feasibilit
[NASA-CR-140805] p0078 N75-11414 WHITE, L. P.	study, volume 1 [PB-234057/8] p0044 N75-1068
Natural-resources mapping of Ethiopia from ERTS-1	The Coho Project: Living resources prediction feasibilit study. Volume 2: Environmental report
imagery p0031 N75-14235 WHITEHURST, C. A.	[PB-234058/6] p0044 N75-1068 The Coho Project: Living resources prediction feasibility
The use of color infrared imagery for the study of marsh	study. Volume 3: System evaluation report
buggy tracks p0061 N75-15127 WHITELEY, A.	[PB-234059/4] p0044 N75:1068: WUKELIC, G. E.
The role of lidar in environmental sensing p0070 A75-19274	Multidisciplinary applications of ERTS and Skylab dat in Ohio p0015 A75-1723
WIEGAND, C. L.	WUKELINC, G. E.
Land use classification and ground truth correlations from simultaneously acquired aircraft and ERTS-1 MSS data	The application of Skylab EREP data for land us classification and mapping in the Cleveland and Columbu
p0014 A75-17188 Irrigation scheduling, freeze warning and soil salinity	areas [E75-10083] p0023 N75-1420
detecting	,
[E75-10046] p0007 N75-12414 Irrigation scheduling, freeze warning and soil salinity	Y
detecting	VAVOVIEW N. A.
[E75-10050] p0050 N75-13350 Irrigation scheduling, freeze warning and soil salinity	YAKOVLEV, N. A. Geological and geophysical studies from space an
detecting [E75-10084] p0008 N75-14202	prospects for their development p0035 N75-1143 YAMAMOTO, G.
Irrigation scheduling, freeze warning and soil salinity detecting	Radiative transfer of visible radiation in turbi atmosphere p0019 N75-1155
[E75-10109] p0010 N75-15125	YARGER, H. L
WIELCHOWSKY, C. C. The remote-sensing program of the Geological Survey	Skylab study of water quality [E75-10075] p0051 N75-1419
of Alabama p0069 A75-17230 WIESNET, D. R.	YEN, J. L. Performance of a hologram matrix radar
Mapping of the 1973 Mississippi River floods by the	p0065 A75-1479
NOAA-2 satellite p0028 A75-17139 WILCOX, L. E.	YENTSCH, C. S. Ocean color, a three component system?
Fully automated compilation of gravity and geophysical maps	[NASA-CR-139145] p0044 N75-1141 YOST, E.
[AD-785838] p0032 N75-15147	In situ spectroradiometric calibration of EREP image:
WILHEIT, T. T. A radio picture of the earth p0066 A75-16191	and oceanography of Block Island Sound [E75-10042] p0072 N75-1241
WILLIAMS, C. K. Summary of flight performance of the Skylab Earth	Multispectral photography for earth resources [NASA-CR-141283] p0074 N75-1513
Resources Experiment Package /EREP/ p0076 A75-17211	YOUNES, H. A. Spectral reflectance studies on mineral deficiency in cor
WILLIAMS, D. L.	plants p0003 A75-1717
Classification and mapping of coal refuse, vegetative cover types, and forest types by digital processing ERTS-1	YOUNG, J. D. Simultaneous active and passive microwave respons
data p0054 A75-17110 WILLIAMS, R. S., JR.	of the earth - The Skylab radscat experiment p0012 A75-1711
Environmental studies of Iceland with ERTS-1 imagery p0012 A75-17105	Preliminary analysis of Skylab RADSCAT results over the ocean
WILLIAMSON, A. N., JR.	[NASA-CR-141454] p0046 N75-1479
Automated selection and location of terrain features on the basis of spectral characteristics	YUNGHANS, R. S. Application of ERTS-1 data to the protection an
[AD-785698] p0030 N75-13375 WILMARTH, V. R.	management of New Jersey's coastal environment p0042 A75-1723
Skylab earth resources program p0011 A75-13577 WILSON, L. B.	
The economic value of remote sensing of earth resources	Z
from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources:	ZACHARY, A. L.
Agriculture. Part 1: Overview [NASA-CR-141265] p0079 N75-14205	Mapping soil features from multispectral scanner data
WIMBERLY, E. T.	p0027 A75-1543 ZAFONTE, L
Surface water modeling Everglades Water Basin, Florida	Airborne survey of major air basins in Catifornia
	p0011 A75-1436
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V.	ZAITSEV, V. M.
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V. The point visibility meter - A forward scatter instrument	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base on a stereomodel of terrain
[E74-10814] p0049 N75-10540 WINSTANLEY, J, V. The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient p0070 A75-19266	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V. The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base on a stereomodel of terrain [AD-784170] p0057 N75-1145 ZAITZEFF. JB. Computer derived coastal water classifications vi
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V. The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient p0070 A75-19266 The role of lidar in environmental sensing p0070 A75-19274 WOBBER, F. J.	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base on a stereomodel of terrain [AD-784170] p0057 N75-1145 ZAITZEFF, J. B. Computer derived coastal water classifications vi spectral signatures p0042 A75-1717 ZELENKA, J. S.
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V. The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient p0070 A75-19266 The role of lidar in environmental sensing p0070 A75-19274 WOBBER, F. J. Application of ERTS-1 data to the protection and management of New Jersey's coastal environment	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base on a stereomodel of terrain [AD-784170] p0057 N75-1145 ZAITZEFF, J. B. Computer derived coastal water classifications vi spectral signatures p0042 A75-1717
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V. The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient p0070 A75-19266 The role of lidar in environmental sensing p0070 A75-19274 WOBBER, F. J. Application of ERTS-1 data to the protection and management of New Jersey's coastal environment p0042 A75-17233 Trends in the use of remote sensing for state resources	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base on a stereomodel of terrain [AD-784170] p0057 N75-1145 ZAITZEFF, J. B. Computer derived coastal water classifications vi spectral signatures p0042 A75-1717 ZELENKA, J. S. Investigation of microwave hologram techniques for application to earth resources p0067 A75-1719 ZHAGAR, IU. KH.
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V. The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient p0070 A75-19266 The role of lidar in environmental sensing p0070 A75-19274 WOBBER, F. J. Application of ERTS-1 data to the protection and management of New Jersey's coastal environment p0042 A75-17233 Trends in the use of remote sensing for state resources and environmental management p0005 A75-18068	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base on a stereomodel of terrain [AD-784170] p0057 N75-1145 ZAITZEFF, J. B. Computer derived coastal water classifications wispectral signatures p0042 A75-1717 ZELENKA, J. S. Investigation of microwave hologram techniques for application to earth resources p0067 A75-1719 ZHAGAR, IU. KH. Possibilities of calculating the earth's local gravitation field from satellite observations p0029 A75-1852
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V. The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient p0070 A75-19266 The role of lidar in environmental sensing p0070 A75-19274 WOBBER, F. J. Application of ERTS-1 data to the protection and management of New Jersey's coastal environment p0042 A75-17233 Trends in the use of remote sensing for state resources and environmental management p0005 A75-18068 WOLF, J. M. SICLOPS - A system of computer programs for rectified	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base on a stereomodel of terrain [AD-784170] p0057 N75-1145 ZAITZEFF, J. B. Computer derived coastal water classifications vi spectral signatures p0042 A75-1717 ZELENKA, J. S. Investigation of microwave hologram techniques fo application to earth resources p0067 A75-1719 ZHAGAR, IU. KM. Possibilities of calculating the earth's local gravitation field from satellite observations p0029 A75-1852 ZHAGAR, N. V. Possibilities of calculating the earth's local gravitation.
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V. The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient p0070 A75-19266 The role of lidar in environmental sensing p0070 A75-19274 WOBBER, F. J. Application of ERTS-1 data to the protection and management of New Jersey's coastal environment p0042 A75-17233 Trends in the use of remote sensing for state resources and environmental management p0005 A75-18068 WOLF, J. M. SICLOPS - A system of computer programs for rectified mapping of airborne scanner imagery p0055 A75-17156	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base on a stereomodel of terrain [AD-784170] p0057 N75-1145 ZAITZEFF, J. B. Computer derived coastal water classifications vi spectral signatures p0042 A75-1717 ZELENKA, J. S. Investigation of microwave hologram techniques for application to earth resources p0067 A75-1719 ZHAGAR, IU. KH. Possibilities of calculating the earth's local gravitation field from satellite observations p0029 A75-1852 ZHAGAR, N. V. Possibilities of calculating the earth's local gravitation field from satellite observations p0029 A75-1852
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V. The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient p0070 A75-19266 The role of lidar in environmental sensing p0070 A75-19268 WOBBER, F. J. Application of ERTS-1 data to the protection and management of New Jersey's coastal environment p0042 A75-17233 Trends in the use of remote sensing for state resources and environmental management p0095 A75-18068 WOLF, J. M. SICLOPS - A system of computer programs for rectified mapping of airborne scanner imagery p0055 A75-17156 WOODLEY, W. Rainfall estimations from geosynchronous satellite	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base on a stereomodel of terrain [AD-784170] p0057 N75-1145 ZAITZEFF, J. B. Computer derived coastal water classifications vi spectral signatures p0042 A75-1717 ZELENKA, J. S. Investigation of microwave hologram techniques for application to earth resources p0067 A75-1719 ZHAGAR, IU. KH. Possibilities of calculating the earth's local gravitation: field from satellite observations p0029 A75-1852 ZHAGAR, N. V. Possibilities of calculating the earth's local gravitation: field from satellite observations p0029 A75-1852 ZHAROV, V. Possibilities of calculating the earth's local gravitation; field from satellite observations p0029 A75-1852
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V. The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient p0070 A75-19266 The role of lidar in environmental sensing p0070 A75-19274 WOBBER, F. J. Application of ERTS-1 data to the protection and management of New Jersey's coastal environment p0042 A75-17233 Trends in the use of remote sensing for state resources and environmental management p0005 A75-18068 WOLF, J. M. SICLOPS - A system of computer programs for rectified mapping of airborne scanner imagery p0055 A75-17156 WOODLEY, W. L. Rainfall estimations from geosynchronous satellite imagery p0047 A75-13596	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base on a stereomodel of terrain (AD-784170) p0057 N75-1145 ZAITZEFF, J. B. Computer derived coastal water classifications vi spectral signatures p0042 A75-1717 ZELENKA, J. S. Investigation of microwave hologram techniques for application to earth resources p0067 A75-1719 ZHAGAR, IU. KH. Possibilities of calculating the earth's local gravitation field from satellite observations p0029 A75-1852 ZHAGAR, N. V. Possibilities of calculating the earth's local gravitation: field from satellite observations p0029 A75-1852 ZHAGOV, V. P. Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gase using pused hydrogen-fluoride laser radiation
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V. The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient p0070 A75-19266 The role of lidar in environmental sensing p0070 A75-19274 WOBBER, F. J. Application of ERTS-1 data to the protection and management of New Jersey's coastal environment p0042 A75-17233 Trends in the use of remote sensing for state resources and environmental management p0005 A75-18068 WOLF, J. M. SICLOPS - A system of computer programs for rectified mapping of airborne scanner imagery p0055 A75-17156 WOODLEY, W. L. Rainfall estimations from geosynchronous satellite imagery p0047 A75-13596 WOODMAN, R. G. Muttidisciplinary analysis for highway engineering purposes	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base on a stereomodel of terrain [AD-784170] p0057 N75-1145 ZAITZEFF, J. B. Computer derived coastal water classifications vi spectral signatures p0042 A75-1777 ZELENKA, J. S. Investigation of microwave hologram techniques fr application to earth resources p0067 A75-1719 ZHAGAR, IU. KH. Possibilities of calculating the earth's local gravitation field from satellite observations p0029 A75-1852 ZHAGAR, N. V. Possibilities of calculating the earth's local gravitation field from satellite observations p0029 A75-1852 ZHAROV, V. P. Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gase
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V. The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient p0070 A75-19266 The role of lidar in environmental sensing p0070 A75-19267 WOBBER, F. J. Application of ERTS-1 data to the protection and management of New Jersey's coastal environment p0042 A75-17233 Trends in the use of remote sensing for state resources and environmental management p0005 A75-18068 WOLF, J. M. SICLOPS - A system of computer programs for rectified mapping of airborne scanner imagery p0055 A75-17156 WOODLEY, W. L. Rainfall estimations from geosynchronous satellite imagery WOODMAN, R. G. Muttidisciplinary analysis for highway engineering	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base on a stereomodel of terrain [AD-784170] p0057 N75-1145 ZAITZEFF, J. B. Computer derived coastal water classifications vi spectral signatures p0042 A75-1717 ZELENKA, J. S. Investigation of microwave hologram techniques for application to earth resources p0067 A75-1719 ZHAGAR, IU. KH. Possibilities of calculating the earth's local gravitation: field from satellite observations p0029 A75-1852 ZHAGAR, N. V. Possibilities of calculating the earth's local gravitation: field from satellite observations p0029 A75-1852 ZHAROV, V. P. Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gase using pulsed hydrogen-fluoride laser radiation p0063 A75-1144 ZIMAN, Y. L. Space iconics
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V. The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient p0070 A75-19266 The role of lidar in environmental sensing p0070 A75-19274 WOBBER, F. J. Application of ERTS-1 data to the protection and management of New Jersey's coastal environment p0042 A75-17233 Trends in the use of remote sensing for state resources and environmental management p0005 A75-18068 WOLF, J. M. SICLOPS - A system of computer programs for rectified mapping of airborne scanner imagery p0055 A75-17156 WOODLEY, W. L. Rainfall estimations from geosynchronous satellite imagery p0047 A75-13596 WOODMAN, R. G. Muttidisciplinary analysis for highway engineering purposes [E75-10103] p0025 N75-15121 WOODS, G. Pretiminary results of fisheries investigation associated	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base on a stereomodel of terrain (AD-784170) p0057 N75-1145 ZAITZEFF, J. B. Computer derived coastal water classifications vi spectral signatures p0042 A75-1717 ZELENKA, J. S. Investigation of microwave hologram techniques for application to earth resources p0067 A75-1719 ZHAGAR, IU. KH. Possibilities of calculating the earth's local gravitation field from satellite observations p0029 A75-1852 ZHAGAR, N. V. Possibilities of calculating the earth's local gravitation: field from satellite observations p0029 A75-1852 ZHAGOV, V. P. Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gase using pulsed hydrogen-fluoride laser radiation p0063 A75-1144 ZIMAN, Y. L. Space iconics [NASA-TIF-798] Geographical correlation of TV and IR images obtaine
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V. The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient p0070 A75-19266 The role of lidar in environmental sensing p0070 A75-19274 WOBBER, F. J. Application of ERTS-1 data to the protection and management of New Jersey's coastal environment p0042 A75-17233 Trends in the use of remote sensing for state resources and environmental management WOLF, J. M. SICLOPS - A system of computer programs for rectified mapping of airborne scanner imagery p0055 A75-17156 WOODLEY, W. L. Rainfall estimations from geosynchronous satellite imagery p0047 A75-13596 WOODMAN, R. G. Muttidisciplinary analysis for highway engineering purposes [E75-10103] p0025 N75-15121 WOODS, G. Pretiminary results of fisheries investigation associated with Skylab-3 WOODWOTH, B. M.	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base on a stereormodel of terrain [AD-784170] p0057 N75-1145 ZAITZEFF, J. B. Computer derived coastal water classifications via spectral signatures p0042 A75-1717 ZELENKA, J. S. Investigation of microwave hologram techniques for application to earth resources p0067 A75-1719 ZHAGAR, IU. KH. Possibilities of calculating the earth's local gravitation field from satellite observations p0029 A75-1852 ZHAGAR, N. V. Possibilities of calculating the earth's local gravitation field from satellite observations p0029 A75-1852 ZHAROV, V. P. Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gase using pulsed hydrogen-fluoride laser radiation p0063 A75-1144 ZIMAN, Y. L. Space iconics [NASA-TT-F-798] p0057 N75-1141
[E74-10814] p0049 N75-10540 WINSTANLEY, J. V. The point visibility meter - A forward scatter instrument for the measurement of aerosol extinction coefficient p0070 A75-19266 The role of lidar in environmental sensing p0070 A75-19274 WOBBER, F. J. Application of ERTS-1 data to the protection and management of New Jersey's coastal environment p0042 A75-17233 Trends in the use of remote sensing for state resources and environmental management p0042 A75-17233 Trends in the use of remote sensing for state resources and environmental management p0055 A75-18068 WOLF, J. M. SICLOPS - A system of computer programs for rectified mapping of airborne scanner imagery p0055 A75-17156 WOODLEY, W. L. Rainfall estimations from geosynchronous satellite imagery p005MAN, R. G. Multidisciplinary analysis for highway engineering purposes [E75-10103] p0025 N75-15121 WOODS, G. Pretiminary results of fisheries investigation associated with Skytab-3 p0003 A75-17167	ZAITSEV, V. M. Use of a digital computer to construct isoline maps base on a stereomodel of terrain (AD-784170) p0057 N75-1145 ZAITZEFF, J. B. Computer derived coastal water classifications vi spectral signatures p0042 A75-1717 ZELENKA, J. S. Investigation of microwave hologram techniques for application to earth resources p0067 A75-1719 ZHAGAR, IU. KH. Possibilities of calculating the earth's local gravitation field from satellite observations p0029 A75-1852 ZHAGAR, N. V. Possibilities of calculating the earth's local gravitation: field from satellite observations p0029 A75-1852 ZHAGOV, V. P. Optical-acoustic detection of low concentrations of hydrogen fluoride, nitrous oxide, and carbon dioxide in gase using pulsed hydrogen-fluoride laser radiation p0063 A75-1144 ZIMAN, Y. L. Space iconics [NASA-TIF-798] Geographical correlation of TV and IR images obtaine

CORPORATE SOURCE INDEX

Earth Resources/A Continuing Bibliography (Issue 5)

OCTOBER 1975

Typical Corporate Source Index Listing



The title of the document is used to provide a brief description of the subject matter. The page number and the accession number are included in each entry to assist the user in locating the abstract in the abstract section. If applicable, a report number is also included as an aid in identifving the document.

AGRICULTURAL RESEARCH SERVICE, WESLACO, TEX.

Irrigation scheduling, freeze warning and soil salinity detecting [E75-10046] p0007 N75-12414

Irrigation scheduling, freeze warning and soil satinity detecting [E75-10050] p0050 N75-13350

A study of the early detection of insect infestations and density/distribution of host plants

p0008 N75-13362 [E75-10062] Irrigation scheduling, freeze warning and soil salinity

p0008 N75-14202 Irrigation scheduling, freeze warning and soil salinity [E75-10109] p0010 N75-15125

AIR FORCE AVIONICS LAB., WRIGHT-PATTERSON An application of infrared remote sensing techniques to

ecological problems [AD-786028] p0022 N75-13419 AKADEMIYA NAUK SSSR, NOVOSIBIRSK.

Solution of the radiative transfer theory problems by the onte Carlo method p0058 N75-11551 Monte Carlo method

ALABAMA UNIV., HUNTSVILLE. Determination of points of entry for potential contaminants into limestone aquifers using thermal infrared

imagery [NASA-CR-120540] p0021 N75-12546

ALASKA UNIV., FAIRBANKS.

Environmental surveys in Alaska based upon ERTS ata p0017 N75-10551 data AMERICAN UNIV., WASHINGTON, D.C.

Skylab-EREP investigations of wetlands [E74-10816] p00

p0049 N75-10542 Utilization of ERTS-1 imagery in cultivation and settlement site identification and carrying capacity estimates in Upper Volta and Niger [E75-10029] p0008 N75-13345

APPLIED PHYSICS LAB., JOHNS HOPKINS UNIV., SILVER SPRING, MD.

A radar altimeter for satellite application [AD-784373] p0 p0056 N75-10425

ARIZONA UNIV., TUCSON.

A study to explore the use of orbital remote sensing to native arid plant distribution [E74-10818] D0006 N75-10544

ARMY ENGINEER WATERWAYS EXPERIMENT STATION, VICKSBURG, MISS.

Automated selection and location of terrain features on ne basis of spectral characteristics [AD-785698] p0030 N75-13375

ARMY FOREIGN SCIENCE AND TECHNOLOGY CENTER, CHARLOTTESVILLE, VA

Use of a panoramic aerial camera [AD-784162] pc era in aerosurveying p0071 N75-11450

Use of a digital computer to construct isoline maps based

on a stereomodel of terrain [AD-784170] p0057 N75-11453 Investigation of accuracy of determination of plane coordinates by means of the Central Scientific Research Institute of Geodesy, aerial photography and cartography

airborne rangefinder [AD-785886] p0061 N75-14255

Work dealing with densification of altimetric-planometric data points employing universal instruments and radio attimeter and statoscopic indications p0061 N75-14257 [AD-785877]

Morphographic and morphometric analysis of vegetative cover photographs in aerial photography for the mapping

of vegetation [AD-785932] p0009 N75-14258 Scientific research and experimental design planning and

n0081 N75-14259 [AD-785998] The next problems in the area of the implementation of

planning for cartographical-geodetic work [AD-785882] p0032 N75-15143

Use of forest vegetation as an interpretation and large scale mapping of soils in a forest

n0010 N75-15146 Optical quantum generators in geodetic measure

ARMY MATERIAL COMMAND.TEXARKANA, TEX.
Life cycle cost study of Army Spectrometric Oil Analysis
Program (ASOAP)
[AD-786501] p0073 N75-14870

В

BATTELLE COLUMBUS LABS., OHIO.

Interaction of marine geodesy, satellite technology and [NASA-CR-137470]

nO043 N75-10671

Bistatic radar sea state monitoring [NASA-CR-137469] p0043 N75-10673

Calibration and evaluation of Skylab altimetry for geodetic determination of the geoid [E75-10064] p0030 N75-13364

The application of Skylab EREP data for land use classification and mapping in the Cleveland and Columbus

[E75-10083]

BOCHUM OBSERVATORY (WEST GERMANY).

Environmental satellites and ERTS-1 imagen p0059 N75-14217

BOEING CO., KENT, WASH.

Quantitative determination of stratospheric aerosol characteristics

p0016 N75-10535 [E74-10808] Quantitative determination of stratospheric serosol

[E75-10010] p0020 N75-12393

Quantitative determination of stratospheric aerosol characteristics p0023 N75-14198 [E75-10079]

BREVARD COUNTY PLANNING DEPT., TITUSVILLE,

Planning applications in east central Florida [E75-10015] p0021 p0021 N75-13337

BRISTOL UNIV. (ENGLAND).

Ground-truth measurements in relation to aircraft and satellite studies of agricultural land use classification in Britain p0009 p0009 N75-14251

BUREAU DE RECHERCHES GEOLOGIQUES ET MINIERES, ORLEANS (FRANCE).

ERTS image contribution to geological knowledge covering different types of climates (France, Togo, Dahomey, Madagascar) p0038 N75-14240

BUREAU OF MINERAL RESOURCES, GEOLOGY AND

GEOPHYSICS, CANBERRA (AUSTRALIA) A study of the usefulness of Skylab EREP data for earth studies in Australia

p0036 N75-12415 Australian and Soviet gravity surveys along the Australian

p0031 N75-13454 [BULL-161]

BUREAU OF NATURAL GAS, WASHINGTON, D.C. Offshore investigation: Producible shut-in leases as of January 1974, phase 2 p0036 N75-11458

C

CALIFORNIA EARTH SCIENCE CORP., SANTA MONICA.

Lineaments in basement terrane of the Peninsular Ranges, uthern California

[E75-10003] p0035 N75-11407

Fault tectonics and earthquake hazards in the Peninsular Ranges, Southern California

[E75-10005] p0035 N75-11408 Fault tectonics and earthquake hazards in the Peninsular

Ranges, Southern California p0030 N75-13338 [E75-10022]

Analysis of pseudocolor transformations of ERTS-1 nages of Southern California area

(E75-10037) p0036 N75-13346

Fault tectonics and earthquake hazards in the Peninsular Ranges, Southern California [E75-10074]

p0031 N75-14193 Fault tectonics and earthquake hazards in the Peninsular

Ranges, Southern California [E75-10108] p0039 N75-15124

CALIFORNIA UNIV., BERKELEY.

Usefulness of remote sensing techniques for the Usefulness of remote sensing techniques for the secremento-San Joaquin p0025 N75-15135

ALIFORNIA UNIV., LIVERMORE. LAWRENCE LIVERMORE LAB.

Comparison of the Southern and Northern Hemisphere general circulation characteristics as determined by satellite ozone data [UCRL-75556]

p0017 N75-10638 Feasibility study of a Gunn-diode microwave cavity spectrometer as a formaldehyde monitor [UCID-16488] p0071 N75-11311

Environmental definition program [UCID-16565]

p0022 N75-13458 Atmospheric release advisory capability: Research and

JCRL-75644-REV-2] p0025 N75-15202

CALIFORNIA UNIV., LOS ANGELES.
Proceedings of the UCLA International Conference on Radiation and Remote Probing of the Atmosphere

ASA-CR-140816] p0078 N75-11549
Lectures on the scattering of light p0019 N75-11556
Particulate sizes from polarization measurements [NASA-CR-140816]

p0019 N75-11558
CANADA CENTRE FOR REMOTE SENSING, OTTAWA

(ONTARIO).
The Canadian ERTS program p0077 N75-10550 The Canadian ERTS program p0077 N75-10550
CATHOLIC UNIV. OF AMERICA, WASHINGTON, D.C.
Earth Resources Technology Satellite-1 (ERTS-1) data
and anthropology: Use of these data in carrying capacity
estimates for sites in Upper Volta and Niger
[E75-10001] p0078 N75-11406
CENTER FOR NAVAL ANALYSES, ARLINGTON, VA.

The GLOBAL computer program and GLOBEMAP: A special purpose three-dimensional plotter subroutine [AD-785869] p0032 N75-15141 NTRE D'ETUDES PHYTOSOCIOLOGIQUES ET

ECOLOGIQUES, MONTPELLIER (FRANCE).

Qualitative study of some ERTS-1 images of the Golion project. Ecological aspects of terrestrial environments

p0023 N75-14248

CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE, TOULOUSE (FRANCE). ARNICA/ERTS-1 experiment.

Reculte perspectives pOO81 N75-14246
CHESAPEAKE RESEARCH CONSORTIUM, INC., BALTIMORE, MD.

A survey for the use of remote sensing in the Chesapeake Bay region [NASA-CR-137476]

CHICAGO UNIV.. ILL

Remote sounding of stratospheric temperatures using high resolution radiance measurements from the IRIS-D [NASA-CR-141130] p0022 N75-13452 COLORADO SCHOOL OF MINES, GOLDEN.

Geologic and mineral and water resources investigations Western Colorado, using Skylab EREP data

[74-10802] p0034 N75-10530 Index of Skylab data available at Colorado School of lines [E74-10802]

[E75-10041] p0072 N75-12410 Geologic and mineral and water resources investigations in Western Colorado, using Skylab EREP data p0058 N75-12412 [E75-10043]

Geologic and mineral and water resources investigations vestern Colorado, using Skylab EREP data [E75-10071] p0037 N75-14190 Geologic and mineral and water resources investigations in western Colorado, using Skylab EREP data [E75-10105] p0025 N75-15123 COMMITTEE ON AERONAUTICAL AND SPACE SCIENCES (U. S. SENATE). Earth resources satellites [GPO-40-560] p0078 N75-13010 COMMITTEE ON INTERIOR AND INSULAR AFFAIRS (U. S. SENATE).
National Environmental Policy Act of 1969.
Environmental indices, status of development pursuant to Sections 102(2)(B) and 204 of the act [GPO-25-314] p0077 N75-10617 COMMITTEE ON SCIENCE AND ASTRONAUTICS (U. S. HOUSE). Earth resources survey system p0078 N75-12416 [GPO-41-665] COMMONWEALTH SCIENTIFIC AND INDUSTRIAL RESEARCH ORGANIZATION, MELBOURNE (AUSTRALIA). The PUCE programme for terrain evaluation for regineering purposes. 2: Procedures for terrain engineering purposes. classification p0029 N75-10625 [PB-233847/3] CONSIGLIO NAZIONALE DELLE RICERCHE, MILAN (ITALY). Contribution of space platforms to a ground and airborne remote sensing programme over active Italian volcanoes [275-10069] p0037 N75-14188 Contribution of space platforms to a ground and airborne remote-sensing programme over active Italian volcan p0073 N75-14236 Agricultural resources investigations in northern Italy, southern France and Madagascar p0009 N75-14252 CONTROL DATA CORP., MINNEAPOUS, MINN. Correlation and registration of ERTS multispectral imagery [NASA-CR-134294] n0058 N75-12420 CORNELL UNIV., ITHACA, N.Y. Evaluation of Skylab imagery as an information service for investigating land use and natural resource
[E74-10730] p0056 p0056 N75-10526 Evaluation of Skylab imagery as an information service for investigating land use and natural resources p0020 N75-12397 [E75-10018] Evaluation of Skylab imagery as an information service for investigating land use and natural resources
[E75-10013] p0059 N75-13336 Evaluation of Skylab imagery as an information service for investigating land use and natural resources [E75-10068] p0059 N p0059 N75-14187 CORNELL UNIV., NEW YORK.

Evaluation of Skylab imagery as an information service for investigating land use and natural resources [NASA-CR-141372] p0024 N p0024 N75-15100 CORPS OF ENGINEERS, WALTHAM, MASS.
New England reservoir management
[E75-10065] p0050 N p0050 N75-13365 CORPS OF ENGINEERS, WASHINGTON, D.C. ERTS program of the US Army Corps of Engineers p0042 N75-10554 COUNCIL FOR SCIENTIFIC AND INDUSTRIAL RESEARCH, PRETORIA (SOUTH AFRICA). Research in earth resources utilization, physical ngineering, and pollution p0022 N75-13369 DARTMOUTH COLL. HANOVER, N.H. Investigation of land use of northern megalopolis using ERTS-1 imagery INASA-CR-24591 n0017 N75-10545 DEFENCE RESEARCH ESTABLISHMENT OTTAWA (ONTARIO).

An examination of whiteouts and transmission through ice fog in the visible and near infrared [DREO-TN-74-18] p0021 N75-12467 DEFENSE MAPPING AGENCY AEROSPACE CENTER. ST. LOUIS, MO. Fully automated compilation of gravity and geophysical maps [AD-785838] p0032 N75-15147 DELAWARE UNIV., NEWARK. DEPARTMENT OF THE ENVIRONMENT, OTTAWA (ONTARIO). for retransmission of water resources data [E74-10805]

DEUTSCHE FORSCHUNGS- UND VERSUCHSANSTALT FUER LUFT- UND RAUMFAHRT, OBERPFAFFENHOFEN (WEST GERMANY). Mapping of water bodies in northern Germany from ERTS p0051 N75-14227 Theoretical investigations on the determination of the vertical temperature distribution in the atmosphere assuming multispectral thermal radiation measurements at CALIF.
Image selection system
[NASA-CR-2475] ENVIRONMENTAL DATA SERVICE, WASHINGTON, [BMFT-FB-W-74-04] p0024 N75-14331 DEUTSCHES HYDROGRAPHISCHES INSTITUT, HAMBURG (WEST GERMANY) Use of ERTS-1 in sea-ice studies p0045 N75-14234 [E75-10036] E EARTH SATELLITE CORP., WASHINGTON, D.C. An evaluation of the suitability of ERTS data for the p0035 N75-10553 ANN ARBOR. purposes of petroleum exploration Developing processing techniques for Skylab data [E74-10799] p0057 N75-The ERTS cost-benefit study [PB-234065/1] p0079 N75-13373 EAST ANGLIA UNIV., NORWICH (ENGLAND).

Generalisation of ERTS data for global-scale investigations p0081 N75-14224 and methodological issues [NASA-CR-140674] ECOLE NATIONALE SUPERIEURE DES MINES, PARIS (FRANCE).

Colour composites of ERTS multispectral imagery for Study of atmospheric effects in Skylab data [E75-10006] applications in hydrology and oceanography p0045 N75-14231 and ferrous compounds from a space platform [E75-10008] p0036 ECOLE NATIONALE SUPERIEURE DU PETROLE ET DES MONTEURS, RUEIL MALMAISON (FRANCE). Recent and present sedimentation complexity of the Rhone delta and the French mediterranean coast (Golion imagery [E75-10020] project) p0045 N75-14228 ECON, INC., PRINCETON, N.J. imagery [E75-10034] The value of improved (ERS) information based on omestic distribution effects of U.S. agriculture crops p0007 N75-12423 [NASA-CR-141046] Improved (ERTS) information and its impact on U.S. [E75-10039] Developing processing techniques for Skylab data [E75-10040] p0058 N75-1 markets for agricultural commodities: A quantitiative economic investigation of production, distribution and net export effects p0078 N75-12424 [NASA-CR-141044] information system [NASA-CR-140357] The role of ERTS in the establishment and of a nationwide land cover information system
[NASA-CR-141045] Oil pollution nforcement p0020 N75-12425 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 1: Summary [NASA-CR-141268] p0079 N75-14203 [E75-10027] The economic value of remote sensing by satellite: An ERTS overview and the value of continuity of service. techniques [NASA-CR-141193] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources: Agriculture. Part 1: Overview techniques [PB-235992/5] p0079 N75-14205 [NASA-CR-141265] The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources, agriculture. Part 2: Distribution effects
[NASA-CR-141194] p0079 N75-14206 The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 3: Intensive use of living resources, agriculture. Part 3: The integrated impact of improved (ERS) information on US agricultural commodities. INASA.CR.1412621 [E75-10093] [E75-10094] [NASA-CR-141262] p0079 N75-14207 [NASA-CR-141262] p0079 N75-14207
The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 4: Forestry, wildlife and rangeland [NASA-CR-141263] p0080 N75-14208
The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 5: Inland water resources [NASA-CR-141269] [E75-10096] ESL, INC., SUNNYVALE, CALIF. of service. Volume 5: [NASA-CR-141269] JSC version ([NASA-CR-140334] (NASA-CR-141269)
The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 1: Introduction EUROPEAN SPACE RESEARCH ORGANIZATION, PARIS (FRANCE). European earth-resources satellite experiments [ESRO-SP-100] p0081 N [NASA-CR-141266] p0080 N75-14210

Monitoring physical and chemical parameters of Delaware Bay waters with an ERTS-1 data collection platform [E75-10090] p0051 N75-15108

Water survey of Canada: Application for use of ERTS-A 74-10805] p0049 N75-11409
Water survey of Canada: Application for use of ERTS-A for retransmission of water resources data
[E75-10026] p0050 N75-13342 Water survey of Canada: Application for use of ERTS-A for retransmission of water resources data [E75-10053] p0050 N75-13353

The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 6: Land use. Part 2: The role of ERTS in the establishment and updating of a nationwide land cover information system [NASA-CR-141267] p0080 N75-14211

[MASA-CR-141267]
The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 7: Nonreplenishable natural resources: Minerals, fossil fuels and geothermal energy sources [NASA-CR-141264] n0080 N75-14212

[NASA-CR-141264] p0080 N/5-14212
The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 8: Atmosphere [NASA-CR-141191] p0080 N75-14213

[MASA-CR-141191] p0080 N75-14213
The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 9: Oceans
-[NASA-CR-141270] p0080 N75-14214

CORPORATE SOURCE INDEX The economic value of remote sensing of earth resources from space: An ERTS overview and the value of continuity of service. Volume 10: Industry [NASA-CR-141192] p0080 N75-14215 ECTROMAGNETIC SYSTEMS LABS., SUNNYVALE, Catalog of earthquake photographs p0072 N75-13455 ENVIRONMENTAL RESEARCH AND TECHNOLOGY. INC., LEXINGTON, MASS.
Experimental evaluation of atmospheric effects on radiometric measurements using the EREP of Skylab
[E75-10036] p0020 N75-12406 Study to develop improved spacecraft snow survey methods using Skylab/EREP data
[E75-10081] p0073 N75-14200 NVIRONMENTAL RESEARCH INST. OF MICHIGAN, p0057 N75-10527 [E74-10799] pnuo / N75-10327
Utility of ERTS for monitoring the breeding habit of migratory waterfowl p0006 N75-10557
The application of remote sensing techniques: Technical p0070 N75-10564 p0020 N75-12391 Mapping exposed silicate rock types and exposed ferric p0036 N75-12392 Study of recreational land and open space using Skylab p0020 N75-12399 Study of recreational land and open space using Skylab A0020 N75, 12404 A Skylab program for the International Hydrological Decade (IHD) p0050 N75-12408 [275-10040] POOSS N75-12409
New theoretical models and ratio imaging techniques associated with the NASA earth resources spectral p0058 N75-12417 detection, monitoring and law Skylab water depth determination 75-10048l [E75-10048] p0050 N75-13348
The remote identification of terrain features and materials p0050 N75-13348 at a California test site: An investigative study of p0031 N75-14261 The remote identification of terrain features and materials at Pennsylvania test sites: An investigative study of p0031 N75-14263 Developing processing techniques for Skylab data [E75-10082] p0061 N75-15102 Study of recreational land and open space using Skylab p0024 N75-15110 A Skytab program for the International Hydrological Decade (IHD) p0052 N75-15111 A Skylab program for the International Hydrological Decade (IHD) p0052 N75-15112 Determination of the earth's aerosol albedo using Skylab p0025 N75-15114 Investigation of transient earth resources phenomena:
Continuation study p0073 N75-15131

Earth resources sensor data handling system: NASA

p0072 N75-12418

p0081 N75-14216

FLORIDA STATE UNIV., TALLAHASSEE.
The Coho Project: Living resources prediction feasibility study, volume 1 [PB-234057/8] p0044 N75-10681 The Coho Project: Living resources prediction feasibility study. Volume 2: Environmental report [PB-234058/6] p0 [PB-234058/6] p0044 N75-10682

The Coho Project: Living resources prediction feasibility study. Volume 3: System evaluation report [P8-234059/4] p0044-N75-10683

FREIE UNIV., BERLIN (WEST GERMANY).

Meteorological data. Volume 145 no. 1: The European climatology 1974 meteorological satellite data from the Berlin zone of reception, part 1, 1st quarter p0023 N75-13479

Meteorological data. Volume 139, no. 4: The European

climatology, 1973. Meteorological satellite-data from the Berlin zone of reception, part 4, fourth quarter

LEIDEN UNIV. OBSERVATORY (NETHERLANDS) p0023 N75-13483 for the Arcachon basin and Rhone delta p0060 N75-14223 INSTITUT GEOGRAPHIQUE NATIONAL PARIS LENINGRAD (A. A. ZHDANOV) STATE UNIV. (USSR).
Influence of the atmosphere on spectral radiance and
contrasts of natural formations measured from space G Interpretation of ERTS-1 data for investigating the French western coastal zones (Fralit program) p0019 N75-11557 GENERAL ELECTRIC CO., PHILADELPHIA, PA. ERTS 1 flight evaluation report, 23 October 1973 - 23 January 1974 p0051 N75-14230 LIBRARY OF CONGRESS, WASHINGTON, D.C.
National Environmental Policy Act of 1969.
Environmental indices, status of development pursuant to Comparison of various numerical techniques for classification of multispectral data. Application to geology p0077 N75-10546 [NASA-CR-140712] p0038 N75-14244 Sections 102(2)(B) and 204 of the act ERTS 1 flight evaluation report, 23 January - 23 April INTERNATIONAL COUNCIL OF SCIENTIFIC UNIONS, p0077 N75-10617 [GPO-25-314] p0077 N75-10547 WUBWANA UNIV. (YUGOSLAVIA). [NASA-CR-140810] Report of the meeting on drifting buoys for the first GARP Some analytical results for radiative transfer in thick atmospheres p0019 N75-11555 ERTS 1 flight evaluation report, January 23, 1973 to April 23, 1973 [NASA-CR-140809] p0077 N75-10548 global experiment [GARP-SPEC-REPT-13] p0043 N75-10674 LOCKHEED ELECTRONICS CO., HOUSTON, TEX. INTERNATIONAL INST. FOR AERIAL SURVEY AND ERTS 1 flight evaluation report, 23 April 1974-23 July JSC earth resources data analysis capabilities available EARTH SCIENCES, ENSCHEDE (NETHERLANDS). [NASA-TM-X-72029] Preliminary results of the interpretation of ERTS-1 imagery for soil surveys in the Merida region, Spain [NASA-CR-140711] p0057 N75-10570 n0077 N75-10568 Correlation of missions 191, 51M and helicopter GEOLOGICAL SURVEY, ATLANTA, GA. n0009 N75-14247 Geologic evaluation and applications of ERTS-1 image otography INTERNATIONAL TRAINING CENTRE FOR AERIAL p0034 N75-10552 [NASA-CR-134289] p0058 N75-12279 over Georgia SURVEY, DELFT (NETHERLANDS).
The ITC 8-image comparator and its possible use in ERTS Feasibility study ASCS remote sensing/compliance GEOLOGICAL SURVEY, BAY SAINT LOUIS, MISS. determination system [NASA-CR-134288] Hydrologic significance of lineaments in central and FRFP investigations n0060 N75-14222 p0007 N75-12419 IOWA UNIV. IOWA CITY. [E75-10030] p0050 N75-12401 LONDON UNIV. (ENGLAND). Experiment to evaluate feasibility of utilizing Skylab-EREP remote sensing data of tectonic analysis of the Bighorn GEOLOGICAL SURVEY, DENVER, COLO. Recognition and interpretation of spectral signatures of vegetation from aircraft and satellite imagery in Western Remote sensing geophysics from Skylab Mountains region, Wyoming-Montana p0030 N75-12413 p0008 N75-14241 [E75-10045] eensland, Australia p0030 N75-12395 LONG ISLAND UNIV., GREENVALE, N.Y. Remote sensing geophysics from Skylab [E75-10051] p0036 N75-13351 In situ spectroradiometric calibration of EREP imagery and oceanography of Block Island Sound [E75-10042] p0 Remote nsing geophysics from Skylab J p0030 N75-13352 DO072 N75-12411 [E75-10052] Multispectral photography for earth resources [NASA-CR-141283] p0074 N Application of Skylab EREP photographs to study of the DO074 N75-15138 JET PROPULSION LAB., CALIF. INST. OF TECH., modern episode of accelerated erosion in southern Arizona LOUISIANA STATE UNIV., BATON ROUGE. PASADENA. [E75-10054] p0036 N75-13354 reliminary results in the comparison of Skylab, ERTS The use of color infrared imagery for the study of marsh p0061 N75-15127 buggy tracks and RB-57 images for the detection of lineaments and fractures in Precambrian, Paleozoic, and late Tertiary rocks on and near the Colorado Plateau, North-Central Arizona Applications of Skylab EREP photographs to mapping of landforms and environmental geology in the Great Plains LOUVAIN UNIV. (BELGIUM). Preliminary evaluation of ERTS-1 imagery of Central and Midwaet p0037 N75-13360 p0060 N75-14242 [F75-10060] [E75-10055] p0037 N75-13355 Pedological interpretation of ERTS-1 imagery of the Rima-Sokoto river basin (N. Nigeria) p0061 N75-14243 Remote ser [£75-10057] nsing geophysics from Skylat The availability of local aerial photography in southern p0037 N75-13357 p0025 N75-15126 [NASA-CR-141314] Remote sensing geophysics from Skyla [E75-10058] p0 p0037 N75-13358 JOHNS HOPKINS UNIV., BALTIMORE, MD.

BESMEX: Bering Sea marine mammal experiment
[NASA-TM-X-62399] p0008 N75-13503 M GEOLOGICAL SURVEY, MIAMI, FLA. Surface water modeling Everglades Water Basin, Florida [E74-10814] MAINE STATE HIGHWAY DEPT., BANGOR. JOINT PUBLICATIONS RESEARCH SERVICE. p0049 N75-10540 Multidisciplinary analysis for highway engineering ARLINGTON, VA. GEOLOGICAL SURVEY, RESTON, VA. Status and prospects of development of space geodes p0025 N75-15121 Overall evaluation of ERTS-1 imagery for cartographic p0029 N75-10869 MAINZ UNIV. (WEST GERMANY). Automated system remote measuring Scattering and absorption from poly-dispersed aerosols [E74-10800] p0029 N75-10528 p0007 N75-12518 agrometeorology Dynamics of suspended sediment plumes in Lake p0019 N75-11553 MASSACHUSETTS UNIV., GLOUCESTER.
Ocean color, a three component system?
[NASA-CR-139145] p004 Analysis of methods for the geodetic support of complex [E74-10815] p0049 N75-10541 a geophysical surveys p0031 N75-13498 [JPRS-63621] n0044 N75-11411 Urban and regional land use analysis: CARETS and THEMATICAL APPLICATIONS GROUP, INC., census cities experiment package ELMSFORD, N.Y. [E75-10019] p0020 N75-12398 An extension of the combinatorial geometry technique Κ Urban and regional land use analysis: CARETS and census cities experiment package [E75-10056] for modeling vegetation and terrain features [AD-782883] p0006 p0006 N75-10577 o0022 N75-13356 SSERSCHMITT-BOELKOW-BLOHM G.M.B.H., KAKIOKA MAGNETIC OBSERVATORY (JAPAN). Certographic evaluation of Skylab-A S-192 scanner MUNICH (WEST GERMANY). Report of the geomagnetic and geoelectric observations, 1972 (rapid variations) Reduction of information redundancy in ERTS-1 and [E75-10078] p0031 N75-14197 [REPT-13] p0032 N75-15228 EREP data. A proposal for an experimental study p0060 N75-14225 Urban and regional land use analysis: CARETS and KANNER (LEO) ASSOCIATES, REDWOOD CITY, cities experiment package MIAMI UNIV., CORAL GABLES, FLA. p0024 N75-15109 [E75-10091] Application of remote sensing for prediction and detection Preliminary data on the UHF radiometric research of soil of thermal pollution [NASA-CR-139182] Urban and regional land use analysis: CARETS and wetness, conducted in the USSR, in 1974 census cities experiment package p0009 N75-15097 [NASA-TT-F-16151] Feasibility of remote sensing for detecting thermal pollution. Part 1: Feasibility study. Part 2: Implementation [E75-10100] p0025 N75-15118 ANSAS UNIV. CENTER FOR RESEARCH, INC., GEOLOGICAL SURVEY, TACOMA, WASH.

Evaluate ERTS imagery for mapping and detection of LAWRENCE. Detection of moisture and moisture related phenomena from Skylab NASA-CR-134453] p0025 N75-15199 changes of snowcover on land and on glaciers MILAN UNIV. (ITALY). [E74-10812] p0049 N75-10539 [E75-10035] p0007 N75-12405 Geological features on ERTS-1 images of a test area in GEOPHYSICAL OBSERVATORY, SODANKYLA p0037 N75-14237 the west-central Alps
MINNESOTA UNIV., MINNEAPOLIS. Detection of moisture and moisture related phenomena (FINLAND).

INSTITUT DE GEOGRAPHIE. AIX EN PROVENCE

Optical processing of ERTS-1 (MSS) images: Examples

Н

agnetic storm, 17-18 December 1971

at Sodankyla during a

p0030 N75-13432

from Skylab

[E75-10007]

[E75-10011]

[E75-10070]

[E75-10073]

[E75-10080]

instrument S-193

processing of ERTS imagery

nt S: 193

HUNTING SURVEYS, LTD., BOREHAM WOOD (ENGLAND)

Geophysical variations

The potential of small-scale mapping from ERTS-1 p0060 N75-14221 Natural-resources mapping of Ethiopia from ERTS-1

p0031 N75-14235

1

INFORMATICS, INC., ROCKVILLE, MD.

Soviet satellite geodesy: Basis and status of Soviet laser sodesy and long-base line interferometry p0029 N75-10524 [AD-781138/3]

[NASA-CR-141454] p0046 N75-14793 KANSAS UNIV., LAWRENCE. Skylab study of water quality [E75-10075] p0051 N75-14194

Use of feature extraction techniques for the texture and context information in ERTS imagery: Spectral and textural

Design data collection with Skylab/EREP microwave

Detection of moisture and moisture related phenomena

Design data collection with Skylab/EREP microwave

Preliminary analysis of Skylab RADSCAT results over

p0007 N75-13333

p0059 N75-13335

p0059 N75-14189

p0008 N75-14192

p0072 N75-14199

Ν

[PB-236041/0]

MISSISSIPPI STATE UNIV., STATE COLLEGE.

A study of Minnesota forests and lakes using data from Earth Resources Technology Satellites [NASA-CR-140865] p0007 N75-12421

Application of remote sensing to state and regional problems
[NASA-CR-140805] p0078 N75-11414

MONTANA STATE UNIV., BOZEMAN.

Water recourse and his

Water resource and hazard planning report for the Clark Fork River Valley above Missoula, Missoula County,

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION, AMES RESEARCH CENTER, MOFFETT FIELD, CALIF.

Systems study of an automated fire weather data [NASA-TM-X-62402] p0006 N75-12277

p0051 N75-14260

p0036 N75-11437

p0036 N75-11438

Use of TV pictures obtained from weather satellites to

Oust storms according to data of space research [NASA-TT-F-16021] p0021 N75-12463

study the geological structure of the earth

BESMEX: Bering Sea marine mammat experiment [NASA-TM-X-62399] p0008 N75-13503 NATIONAL AERONAUTICS AND SPACE Skylab: A proposal aerotriangulation with very small PHYSICS LAB. RVO-TNO, THE HAGUE (NETHERLANDS) scale photography [E75-10101] Pattern recognition relevant to reconnaissance and n0073 N75-15119 ADMINISTRATION. GODDARD SPACE FLIGHT
CENTER, GREENBELT, MD.
Third Earth Resources Technology Satellite Symposium. NATIONAL OCEANIC AND ATMOSPHERIC emote sensina 00070 N75-10573 ADMINISTRATION, WASHINGTON, D.C.

A review of initial investigations to utilize ERTS-1 data [PHI-1974-23] PURDUE UNIV., LAFAYETTE, IND. Volume 2: Summary of results [NASA-SP-356] in determining the availability and distribution of living An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado Mountains, using p0077 N75-10549 p0043 N75-10555 p0043 N75-10559 Water resources A cloud physics investigation utilizing Skylab data 75-10021] p0020 N75-12400 tomatic data processing techniques Mineral resources, geological structures, and landform p003\$ N75-10561 [E74-10803] p0057 N75-10531 An interdisciplinary analysis of multispectral satellite data A cloud physics investigation utilizing Skylab data [E75-10099] p0025 N75-1 Exploration for fossil and nuclear fuels from orbital for selected cover types in the Colorado Mountains, using DO025 N75-15117 Geologic structure in California: Three studies with ERTS-1 imagery [NASA-TM-X-70799] automatic data processing techniques IATIONAL RESEARCH COUNCIL, BANGKOK p0078 N75-12390 (THAILAND). Spectroradiometric and multispectral scanner over Lake Thailand national programme of the Earth Resources fechnology Satellite [F75-10066] p0072 N75-13366 p0077 N75-10533 [F74-10806] A demonstration of ERTS-1 analog and digital techniques An interdisciplinary analysis of multispectral satellit A demonstration of ENIS-1 analog and digital techniques applied to strip mining in Maryland and West Virginia [NASA-TM-X-70798] p0039 N75-15133 NATIONAL AERONAUTICS AND SPACE ADMINISTRATION, LANGLEY RESEARCH CENTER, LANGLEY STATION, VA. The Thailand programme of the Earth Resources for selected cover types in the Colorado Mountains, using automatic data processing techniques

[E75-10072] p0037 N75-14191 Technology Satellite [E74-10807] NAVAL RESEARCH LAB., WASHINGTON, D.C. Reformatting Skylab and MSS data over northern Terrain properties and topography from Skylab A simulation of synthetic aperture radar imaging of ocean [E75-10016] p0061 N75-15101 p0030 N75-13359 Analysis of Skylab MSS data, and calibration of thermal data over Lake Monroe, Indiana [E75-10059] [NASA-TM-X-72629] n0044 N75-12172 NEBRASKA UNIV., LINCOLN. NATIONAL AERONAUTICS AND SPACE ADMINISTRATION, LYNDON B. JOHNSON SPACE [E75-10095] Satellite pictures can help guide Sandhills irrigation [E75-10023] p0021 N75-13339 An interdisciplinary analysis of multispectral satellite data for selected cover types in the Colorado mountains, using automatic data processing techniques
[E75-10104] p0073 N75-15122 CENTER, HOUSTON, TEX.
Agriculture, forestry, range resources Use of ERTS-1 imagery to interpret wind-erosion hazard in the Sandhills of Nebraska [E75-10044] 00006 N75-10558 n0059 N75-13347 JSC earth resources data analysis capabilities available NEVADA UNIV., RENO. to EOD revision 8 [NASA-TM-X-72029] p0057 N75-10570 The ERTS-1 investigation. Volume 7: Erts-1 land-use analysis of the Houston area test site The Great Basin investigation [E75-10076] R p0037 N75-14195 The Great Basin investigation RAND CORP., SANTA MONICA, CALIF. [E75-10102] p0038 N75-15120 [E75-10009] p0021 N75-13334 The ERTS-1 investigation (ER-600), Volume 2: ERTS-1 Analysis of pseudocolor transformations of ERTS-1 images of Southern California area [E75-10037] p0036 N75-13346 NEW YORK STATE MUSEUM AND SCIENCE SERVICE, ALBANY,
Assessment of ERTS-1 imagery as a tool for regional geological analysis in New York State coastal/estuarine analysis p0051 N75-13371 READING UNIV. (ENGLAND).

A comparison of the capabilities of ERTS-1 and the The Proceedings of the Skylab Life Sciences Symposium, p0034 N75-10536 [E74-10809] Skylark earth-resources rocket for resources sur [NASA-TM-X-58154-VOL-1] NORTH CAROLINA STATE UNIV., RALEIGH. central Argentina p0023 N75-14250
RESEARCH INST. OF NATIONAL DEFENCE, Utilization of EREP data in geological evaluation regional planning, forest management, and water management in Skylab environmental and crew microbiology studies p0024 N75-14366 NATIONAL AERONAUTICS AND SPACE STOCKHOLM (SWEDEN) North Carolina Recordings of the spectral-radio-meter SIGNE during the ADMINISTRATION. MARSHALL SPACE FLIGHT CENTER, HUNTSVILLE, ALA. [E75-10063] n0008 N75-13363 ummer of 1972 NORTHERN PRAIRIE WILDLIFE RESEARCH CENTER. [FOA-2-C-2577-E1] p0072 N75-12283 OA-2-C-2577-E1]
Processing of ERTS CCT imagery of Sweden. A pilot p0060 N75-14219 MSFC integrated experiments preliminary report
[NASA-TM-X-64881] p0078 N75-10967
Application of a single laser Doppler system to the JAMESTOWN, N. DAK.

Utilization of Skylab (EREP) system for appraising changes in continental migratory bird habitat [E74-10817] p0006 RHODE ISLAND UNIV., KINGSTON. measurement of atmospheric winds [NASA-TM-X-64891] p0006 N75-10543 Laser Raman spectroscopy of solutes dissolved in water Utilization of Skylab (EREP) system for appraising from a remote platform [PB-233432/4] Computer and photogrammetric general land use study n0056 N75-10202 changes in continental migratory bird habitat of central north Alabama [NASA-TR-R-431] RICE UNIV., HOUSTON, TEX.

An analysis of applications development systems for remotely sensed, multispectral data for the Earth p0008 N75-13361 [E75-10061] p0022 N75-13372 Utilization of Skylab (EREP) system for appraising NATIONAL AERONAUTICS AND SPACE ADMINISTRATION. MISSISSIPPI TEST FACILITY, changes in continental migratory bird habitat [E75-10087] p0009 Observations Division of the NASA Lyndon B. Johnson BAY SAINT LOUIS. NORWEGIAN WATER RESOURCES AND ELECTRICITY BOARD, OSLO. p0017 N75-10560 Land use and mapping [NASA-CR-140345] n0071 N75-11447 Earth resources data analysis program, phase 2
[NASA-CR-141477] p0074 N75-15333 Evaluation of satellite remote sensing and automatic data ERTS-1 images in glaciology and hydrology: Norwegian perience p0051 N75-14233 techniques for, characterization of wetlands and coastal experience ROCKWELL INTERNATIONAL CORP., DOWNEY, Evaluation of satellite remote sensing and automatic data techniques for characterization of wetlands and coastal 0 Measurements of the dielectric properties of sea water [NASA-CR-2458] archiande n0045 N75, 13496 OHIO DEPT. OF ECONOMIC AND COMMUNITY DEVELOPMENT, COLUMBUS.

The application of Skylab EREP data for land use [E75-10028] p0050 N75-13344 NATIONAL AERONAUTICS AND SPACE ADMINISTRATION, WASHINGTON, D.C. S classification and mapping in the Cleveland and Columbus The final Skylab mission: Man at home and at work in [F75, 10083] n0023 N75-14201 [NASA-TM-X-70394] SANDIA CORP. ALBUQUERQUE IN MEX. p0078 N75-12019 OHIO STATE UNIV., COLUMBUS. Modern trends in environmental testing [SLA-73-5931] p0016 N75-10469 ERTS-B (Earth Resources Technology Satellite)
[NASA-NEWS-RELEASE-74-329] p0079 N75 Automated strip mine and reclamation mapping from RTS p0035 N75-10556 p0079 N75-13370 CIENTIFIC TRANSLATION SERVICE, SANTA Future US activities based on experience gained from RTS-1 and EREP p0009 N75-14253 Some first accuracy estimates for applications of aerial BARBARA, CALIF. ERTS-1 and EREP gradiometry [AD-786418] OREGON STATE UNIV., CORVALLIS.
The Coho Project. (2017) Orbital geology [NASA-TT-F-16000] NATIONAL CENTER FOR ATMOSPHERIC RESEARCH, p0035 N75-10887 BOULDER, COLO.

Methods of calculating infrared transfer: The earth in the lens of Salyut-3 [NASA-TT-F-16028] The Coho Project: Living resources prediction feasibility sfer: A review n0019 N75-11560 n0070 N75-10964 study, volume 1 [PB-234057/8] Space iconics [NASA-TT-F-798] n0044 N75-10681 NATIONAL ENVIRONMENTAL SATELLITE SERVICE. The Coho Project: Living resources prediction feasibility udy. Volume 2: Environmental report n0057 N75-11419 WASHINGTON, D.C.
Environmental satellite imagery, September 1974 Geometric distortions of opticommercial panoramic p0044 N75-10682 [PB-234058/6] p0057 N75-11425 n0058 N75-11571 The Coho Project: Living resources prediction feasibility Geographic correlation of television pictures obtained NATIONAL OCEANIC AND ATMOSPHERIC study. Volume 3: System evaluation report [PB-234059/4] p0044 N75-10683 OVERSEAS DEVELOPMENT ADMINISTRATION, p0071 N75-11432 ADMINISTRATION, MIAMI, FLA.

Remote sensing of ocean current boundary layer
[E74-10801] p0042 N75-Geographical correlation of TV and IR images obtained on weather satellites p0071 N75-11433 OVERSEAS DEVELOPMENT ACTION
SITRREY (ENGLAND).
A comparison of ERTS imagery with conventional aerial photography for land-resource surveys in less developed countries. Examples from the Rift Valley Lakes Basin.
Sthionia p0061 N75-14249 p0042 N75-10529 from weather satellites System for electronic transformation and geographic An assessment of the potential contributions to oceanography from Shand-held photography correlation of satellite television information Skylab visual observations and p0071 N75-11434 [E75-10017] n0044 N75-12396 Geographical correlation of earth cloud cover pictures obtained from unoriented satellites p0071 N75-11435 Remote sensing of ocean current boundary layer 75-10031] p0045 N75-14185 OXFORD UNIV. (ENGLAND). Geological and geophysical studies from space and prospects for their development p0035 N75-11436 [E75-10031] p0072 N75-11562 Infrared remote sounding Remote sensing of ocean current boundary layer 75-10067] p0045 N75-14186 Experience in integrated geological and geographical interpretation of space photographs obtained in the USSR [E75-10067] Remote sensing of ocean current boundary layer 75-10098] p0046 N75-15116

PACIFIC SOUTHWEST FOREST AND RANGE EXPERIMENT STATION, BERKELEY, CALIF.

[E75-10032]

Inventory of forest and rangeland resources, including

n0007 N75-12402

[£75-10098]

photography [E74-10804]

NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION, ROCKVILLE, MD.

Skylab a proposal aerotriangulation with very small scale

n0070 N75-10532

CORPORATE SOURCE INDEX From aerospace photographs to forecasting and calculating flows [NASA-TT-F-16006] p0059 N75-13367 Fundamentals of the space indication of earth p0074 N75-15136 [NASA-TT-F-16108] [NASA-TT-F- 10 100] SHEFFIELD UNIV. (ENGLAND). An evaluation of stereoscopic viewing of ERTS and Skylab p0060 N75-14220 SMITHSONIAN ASTROPHYSICAL OBSERVATORY. CAMBRIDGE, MASS.

Mapping of the major structures of the African rift MASS. [F75.10024] n0030 N75-13340 BOUTH DAKOTA STATE UNIV., BROOKINGS.

Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil p0006 N75-12394 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil p0007 N75-12403 Develop techniques and procedures, using multispectral systems, to identify from remotely sensed data the physical and thermal characteristics of plants and soil [E75-10077] p0008 N75-14196 STANFORD RESEARCH INST., MENLO PARK, CALIF. STANFORD UNIV., CALIF. Feasibility of using S-191 infrared spectra for geological studies from space
[E75-10088] p0038 N75-15106 Feasibility of using S-191 infrared spectra for geological studies from space [E75-10089] p0038 N75-15107 STRASBOURG UNIV. (FRANCE). ERTS-1 contributions to the ecogenetical knowledge of the Llanos of the Orinoco (Colombia and Venezuela) p0023 N75-14245 SUOMALAINEN TIEDEAKATEMIA, HELSINKI (FINLAND). lonospheric meeting [REPT-3] p0022 N75-13437 SWISS INST. OF METEOROLOGY, ZURICH. Meteorological interpretation of high resolution images for remote sensing of earth resources p0060 N75-14229 T TECHNISCHE UNIVERSITAET,
CLAUSTHAL-ZELLERFELD (WEST GERMANY):
ERTS-1 data on Afar tectonics p0031 N75-14239
TELESPAZIO, 8.P.A., ROME (ITALY). Digital land-use classification using an ERTS-1 image p0023 N75-14226 TENNESSEE UNIV., KNOXVILLE. Geographic applications of ERTS-1 imagery to landscape change [E74-10810] p0034 N75-10537 ERTS-1 imagery interpretation techniques in the Tennessee Valley [E74-10811] p0016 N75-10538 Investigation of the utility of kinetically controlled and catalytic reactions to perform environmental determinations catalytic reactions to perform environmental determinations by parallel spectrophotometric analysis [TID-26594] p0021 N75-13083 TEXAS A&M UNIV., COLLEGE STATION.

A study of a dual polarization laser backscatter system for remote identification and measurement of water pollution TEXAS INSTRUMENTS, INC., DALLAS.

Oil-on-Water secret Oil-on-Water sensor [AD-784104] p0072 N75-11454 TOHOKU UNIV., SENDAI (JAPAN).
Radiative transfer of visible radiation adiation in turbid p0019 N75-11552 atmosphere TOKYO UNIV. (JAPAN). Investigation of the environmental change pattern of [E75-10086] p0024 N75-15104

V

VARIAN ASSOCIATES, PALO ALTO, CALIF.

Airborne vapor surveillance system [AD-785585]

n0021 N75-12459 VIRGINIA INST. OF MARINE SCIENCE, GLOUCESTER POINT.

emote sensing applications in marine science programs [NASA-CR-137475] p0043 N75-10566

WASHINGTON UNIV., SEATTLE. AIDJEX bulletin number 25 [PB-235298/7]

p0045 N75-14262

WEST VIRGINIA UNIV., MORGANTOWN.

Application and sensitivity investigation of Fourier transforms for microwave radiometric inversions [NASA-CR-140754] p0070 N75-11310

[NASA-CR-140754]
WOLF RESEARCH AND DEVELOPMENT CORP.,
POCOMOKE CITY. MD.
Applicability of Skylab remote sensing for detection and

onitoring of surface mining activities [E75-10038]

75-10038] p0044 N75-12407 Applicability of Skylab remote sensing for detection and monitoring of surface mining activities
[E75-10085] p0038 N75-15103

WOODS HOLE OCEANOGRAPHIC INSTITUTION,

Cruise data report. CHAIN 115 leg 2, Southlant expedition [AD-784421] p0029 N75-10641

WORLD METEOROLOGICAL ORGANIZATION, GENEVA (SWITZERLAND).

Report of the meeting on drifting buoys for the first GARP global experiment
[GARP-SPEC-REPT-13] p0043 N75-10674

Observation and measurement of atm [WMO-368] p0018 N75-11476 A technique for measurement of sulphur dioxide and sulphate concentrations from an aircraft

p0018 N75-11498 Methods for the automatic recording of atmospheric dust concentrations and some of the results obtained

p0057 N75-11502
Aircraft measurements of the concentration and size distribution of atmospheric sulfate particles

Correlation spectroscopy techniques for the regional and global measurements of air pollution p0018 N75-11505 Geophysical monitoring for climatic change. The NOAA rogram p0018 N75-11510 Global measurements of air pollution from satellites

p0018 N75-11521 Measurement of trace stratospheric gases by inverting thermal infrared limb radiance profiles

Ground-based measurements of solar radiation by geophysical monitoring for climatic change (USA) p0018 N75-11527

Microwave rotational spectroscopy: A physical technique pools N75-11529 for specific pollutant monitoring

Measurements of meteorological parameters and air pollution spread over Stockholm, Sweden, by means of remote sensing and airborne instrumentation

Quality control in global and regional air chemistry monitoring networks: Lessons of the European Air Chemistry Network p0018 N75-11540 WYOMING UNIV. LARAMIE.

Usefulness of Skylab color photography and ERTS-1 multispectral imagery for mapping range vegetation types in southwestern Wyoming [E75-10097]

Ζ

ZENTRALSTELLE FUER GEO-PHOTOGRAMMETRIE UND FERNERKUNDUNG, MUNICH (WEST GERMANY).

Evaluation of ERTS-1 data by analogue and digital chniques p0059 N75-14218
Geologic evaluation of central Italy from ERTS-1 and p0038 N75-14238


Skylab data
ZURICH UNIV. (SWITZERLAND).

Methodological aspects and regional examples of now-cover mapping from ERTS-1 and EREP imagery of p0051 N75-14232 the Swiss Alps

CONTRACT NUMBER INDEX

Earth Resources/A Continuing Bibliography (Issue 5)

Typical Contract Number Index Listing



Listings in this index are arranged alphanumerically by contract number. Under each contract number, the accession numbers denoting documents that have been produced as a result of research done under that contract are arranged in ascending order with the AIAA accession numbers appearing first. The accession number denotes the number by which the citation is identified in the abstract section. Preceding the accession number is the page number on which the citation may be found.

AF PROJ. 8607	p0032	N75-14335
AT/2035/025	p0023	N75-14250
DA PROJ. 1G6-62708-D-05	p0006	N75-10577
DA-ARO(D)-31-124-72-G158	p0012	A75-16752
DAAD05-70-C-0197	p0011	N75-12459
	p0006	N75-12439
DAAD05-73-C-0537		
DAAK02-68-C-0089	p0005	A75-18556
DAAK02-73-C-0037	p0001 p0048	A75-16682 A75-17161
DI-14-08-0001-13519	p0048	N75-17161
DI-14-08-0001-13519	p0079	N75-13373
DI-14-08-0001-13911	p0036	N75-17407
Dt-14-31-0001-3641	p0056	N75-10202
DI-14-31-0001-3826	p0051	N75-10202
DOT-CG-31888-A	p0072	N75-11454
DOT-CG-34017-A	p0017	N75-10563
DOT-CG-34071-A	p0014	A75-17205
DSS-0GR2-0522	p0042	A75-17184
DSS-01GRKW111-2-1073	p0042	A75-17184
EPA-IAG-D179(D)	p0018	N75-11529
FH-11-7136	p0031	N75-14261
	p0031	N75-14263
F19628-72-C-0120	p0032	N75-14335
F44620-72-C-0053 MOD-AT/2035/015/SP	p0029	N75-10524
MOD-AT/2035/015/SP	p0067	A75-17192
MOD-AT/2035/025/ASA	p0067	A75-17192
NASA ORDER C-21372-C	p0023	N75-14201
NASA ORDER H-2810-B	p0050	N75-12401
NASA ORDER HT-73	p0047	A75-12873
NASA ORDER L-75902	p0018	N75-11529
NASA ORDER R-03-038-002	p0001	A75-12872
NASA ORDER S-70243-AG	p0029	N75-10528
NASA ORDER S-70243-AG-2	p0049	N75-10539 N75-10541
NASA ORDER 5-70243-AG-2	p0049 p0049	N75-10541
NASA ORDER S-70243-AG-7 NASA ORDER S-70243-G	p0049	A75-17105
NASA ORDER S-70246-AG	p0012	A75-17160
NASA ORDER S-70246-AG-1	p0041	A75-10644
NASA ORDER S-70251-AG	p0004	A75-17176
10,0	p0014	A75-17188
NASA ORDER S-70257-AG	00048	A75-17179
NASA ORDER T-4105-B	p0007	N75-12414
	p0050	N75-13350
	p0008	N75-14202
	p0010	N75-15125
NASA ORDER T-4106-B	p0007	N75-12402
NASA ORDER T-4109-B	p0008	N75-13362
NASA ORDER T-4110-B	p0070	N75-10532
	p0073	N75-15119
NASA ORDER T-4111-B	p0031	N75-14197
NASA ORDER T-4113-B	p0036	N75-13354
NASA ORDER T-4114-B	p0006	N75-10543
	n00008	N75-13381

		:
1		p0009 N75-15105
lumber	NASA ORDER T-4646-B	p0050 N75-13365
	NASA ORDER T-4647-B	p0037 N75-13355
,	NASA ORDER T-4713-8	p0042 N75-10529
		p0044 N75-12396
		p0045 N75-14185
p0043 N75-10566		p0045 N75-14186 p0046 N75-15116
PUU43 N75-1U566		' '1
1 1	'NASA ORDER T-4715-B	p0020 N75-12400
	NACA CROSS TATALS	p0025 N75-15117
	NASA ORDER T-4716-B	p0030 N75-13359
7	NASA ORDER T-5290-B	p0020 N75-12398
ACCESSION		p0022 N75-13356 p0024 N75-15109
NUMBER		p0024 N75-15109 p0025 N75-15118
11	NASA ORDER T-6555-B	p0030 N75-12413
		p0036 N75-13351
		p0030 N75-13352
ed alphanumeri-		p0037 N75-13357
er each contract		p0037 N75-13358
	NASA ORDER T-8217-B	p0003 A75-17167
s denoting docu-	NASA PROJECT 240	p0003 A75-17167
d as a result of	NASW-2481	p0009 N75-15097
.!1	NASW-2483	p0035 N75-10887
act are arranged		p0070 N75-10964
AIAA accession		p0057 N75-11419
ccession number		p0021 N75-12463
		p0059 N75-13367 p0074 N75-15136
the citation is	NASW-2558	p0074 N75-15136 p0007 N75-12423
n. Preceding the		p0007 N75-12424
umber on which		p0020 N75-12425
		p0079 N75-14206
İ		p0080 N75-14211
	NASW-2580	p0079 N75-14203
		p0079 N75-14204
		p0079 N75-14205
		p0079 N75-14207
p0032 N75-14335		p0080 N75-14208]
p0023 N75-14250		p0080 N75-14209 p0080 N75-14210
p0006 N75-10577		p0080 N75-14210
p0012 A75-16752		p0080 N75-14213
p0021 N75-12459		p0080 N75-14214
p0006 N75-10577		p0080 N75-14215
p0005 A75-18556	NAS1-10691	p0045 N75-13496
p0001 A75-16682	NAS1-10810	p0067 A75-17195
p0048 A75-17161	NAS1-12048	p0018 N75-11521
p0079 N75-13373	NAS1-12209 NAS1-12304	p0049 N75-10542
p0035 N75-11407	NAS1-12304	p0041 A75-12034 p0047 A75-16240
p0036 N75-13346	NAS2-7064	p0047 A75-16240 p0058 N75-12422
p0056 N75-10202 p0051 N75-14260	NAS2-7698	p0035 N75-11407
p0072 N75-11454		p0035 N75-11408
p0017 N75-10563		p0030 N75-13338
p0014 A75-17205		p0036 N75-13346
p0042 A75-17184		p0031 N75-14193
p0042 A75-17184	NACE 10000	p0039 N75-15124
p0018 N75-11529	NASS-10306	p0064 A75-12109
p0031 N75-14261	NASS-11074	p0011 A75-11211
p0031 N75-14263	NAS5-11431 NAS5-20021	p0011 A75-11211 p0073 N75-15131
p0032 N75-14335	NASS-20021	p0073 N75-15131 p0056 A75-17202
p0029 N75-10524 p0067 A75-17192	NASS-21635	p0056 A75-17202
p0067 A75-17192 p0067 A75-17192	NAS5-21689	p0003 A75-17146
p0007 A75-17192 p00023 N75-14201	NAS5-21726	p0034 N75-10537
p0050 N75-12401	NAS5-21748	p0030 N75-13340
p0047 A75-12873	NAS5-21749	p0017 N75-10545
p0018 N75-11529	NAS5-21752	p0047 A75-13580
p0001 A75-12872	NAS5-21756	p0021 N75-13339
p0029 N75-10528	NAS5-21764	p0059 N75-13347
p0049 N75-10539	NACE 21700	p0034 N75-10536 p0028 A75-17224
p0049 N75-10541	NASS-21783NASS-21799	p0028 A75-17224 p0010 N75-15115
p0049 N75-10540 p0012 A75-17105	NAS5-21808	p0077 N75-10546
p0012 A75-17105 p0048 A75-17160		p0077 N75-10547
D0048 A75-17180		p0077 N75-10548
p0004 A75-17176		p0077 N75-10568
p0014 A75-17188	NASS-21812	p0006 N75-10544
p0048 A75-17179	NASS-21813	p0044 N75-11411
p0007 N75-12414	NAS5-21822	p0056 A75-17219
p0050 N75-13350	NAS5-21837	p0059 N75-13335 p0041 A75-12034
p0008 N75-14202	147-03-2103/	p0041 A75-12034 p0047 A75-16240
p0010 N75-15125 p0007 N75-12402	•	p0048 A75-17181
p0007 N75-12402 p0008 N75-13362		p0051 N75-15108
p0006 N75-13362 p0070 N75-10532	NAS5-21847	p0021 N75-131337
p0073 N75-15119	NAS5-21860	p0013 A75-17130
p0031 N75-14197		p0054 A75-17130
p0036 N75-13354		*
p0006 N75-10543	NAS5-21875	p0016 N75-10538
p0008 N75-13361	NAS5-21876	p0069 A75-17230

		ОСТ	OBER 1975
5	NAS5-21970		p0078 N75-11406
5	NAS5-23133	***************************************	p0008 N75-13345 p0054 A75-17110
6			p0066_A75-17111
5 6			p0043 N75-10566 p0043 N75-10671
6,		•••••••••••••••••••••••••••••••••••••••	p0043 N75-10673
٦			p0043 N75-10571
3	NAS7-100		p0066 A75-16485 p0066 A75-17119
8			p0005 A75-17973
5 9	NAS8-30216		p0025 N75-15126
в			p0021 N75-12546 p0055 A75-17156
3	-		p0068 A75-17217
2	NAS9-10261	***************************************	p0058 N75-12417 p0001 A75-16242
3!			p0042 A75-17121
<u>:</u> :			p0028 A75-18062 p0005 A75-18556
7	NAS9-11036 NAS9-11188		p0054 A75-17116 p0074 N75-15138
۱ ۲	NAS9-11500	***************************************	p0047 A75-12873
9	NAS9-12200		p0014 A75-17157 p0057 N75-10570
3			p0058 N75-12279
5	NAS9-12776	***************************************	p0007 N75-12419 p0071 N75-11447
3	NAS9-12967		p0074 N75-15333 p0054 A75-17116
5	NAS9-13114		p0058 N75-12420
8	NAS9-13271 NAS9-13272	***************************************	p0051 N75-14194 p0020 N75-12391
3 4	NAS9-13273	***************************************	p0002 A75-17145
5			p0007 N75-12405 p0007 N75-13333
7	NA 50 12274		p0008 N75-14192 p0037 N75-14195
9	NAS9-13274		p0038 N75-15120
2	NAS9-13275		p0050 N75-12408 p0052 N75-15111
3 4			p0052 N75-15112
5	NAS9-13276 NAS9-13278		p0030 N75-13364 p0050 N75-13348
5	NAS9-13279		p0025 N75-15114
1	NAS9-13280		p0057 N75-10527 p0058 N75-12409
2	NAS9-13281		p0061 N75-15102 p0022 N75-13343
2	NAS9-13283		p0020 N75-12399
7			p0020 N75-12404 p0024 N75-15110
3	NAS9-13298		p0010 N75-15115
6	NAS9-13301		p0072 N75-13366 p0061 N75-15101
3	NAS9-13303		p0073 N75-15113 p0016 N75-10535
3 .	11200-10000		p0020 N75-12393
۱	NAS9-13305		p0023 N75-14198 p0073 N75-14200
1	NAS9-13308		p0072 N75-12411
2	NAS9-13310		p0044 N75-12407 p0038 N75-15103
5	NAS9-13312 NAS9-13313	***************************************	p0047 A75-17140 p0030 N75-12395
וו	NAS9-13317		p0003 A75-17158
5	NAS9-13321		p0036 N75-12392 p0008 N75-13363
7	NAS9-13331		p0059 N75-14189
6	NAS9-13337	,,,,,,	p0072 N75-14199 p0006 N75-12394
\$			p0007 N75-12403 p0008 N75-14196
6	NAS9-13343		p0020 N75-12406
3	NAS9-13357	,	p0038 N75-15106 p0038 N75-15107
3	NAS9-13359		p0025 N75-15121
۱ ۱	NAS9-13364		p0056 N75-10526 p0020 N75-12397
3	ı		p0059 N75-13336 p0059 N75-14187
5			p0024 N75-15100
í	NAS9-13380	,	p0057 N75-10531 p0078 N75-12390
3			p0037 N75-14191
3	NAS9-13386	,	p0073 N75-15122 p0064 A75-12036
5		,	p0003 A75-17158 p0034 N75-10530
3	············ 13334	,	p0072 N75-12410
)	!		p0058 N75-12412

CONTRACT NUMBER INDEX

	p0037 N75-14190
	p0025 N75-15123
NAS9-13642	p0046 N75-14793
NAS9-14077	p0072 N75-12418
NAS10-8333	p0054 A75-17115
NAS10-8402	p0025 N75-15199
NAS10-8498	p0017 N75-10572
NGL-05-003-404	p0002 A75-17122
NGL-06-003-200	p0002 A75-17122
NGL-15-005-112	p0027 A75-15435
1100-13-000-112	p0076 A75-17166
	p0005 A75-17166
NGL-16-001-043	p0011 A75-11211
	p0061 N75-15127
NGL-24-005-263	p0007 N75-12421
NGL-25-001-054	p0078 N75-11414
NGL-36-008-093	p0064 A75-12985
NGL-42-003-007	p0055 A75-17135
NGL-44-001-001	p0014 A75-17205
	p0017 N75-10563
NGL-50-002-127	p0066 A75-17141
NGR-05-008-029	p0011 A75-14360
NGR-14-001-194	p0022 N75-13452
NGR-30-001-040	p0011 A75-11221
NGR-39-009-218	p0012 A75-16752
NGR-49-001-056	p0070 N75-11310
NOAA-3-35400	p0063 A75-10790
NOAA-03-3-022-18	p0047 A75-13596
NOAA-04-30-2212	p0041 A75-10644
NOAA-25-22-0211	p0047 A75-13596
NOAA-043-022-28	p0044 N75-10681
	p0044 N75-10682
	p0044 N75-10683
NR PROJ. 083-004	p0029 N75-10641
NR PROJECT 387-069	p0063 A75-10075
	p0028 A75-17201
NSF C-585	p0005 A75-17973
NSF C-625	p0048 N75-14262
NSF GA-1571	p0041 A75-12719
NSF GA-27205	p0041 A75-12719
NSF GI-34809X	p0048 A75-17163
NSF GI-34809X1	p0070 N75-10564
NSF GI-36338X	p0064 A75-13148
NSF GX-33502	p0044 N75-10681
1101 QX 00000	p0044 N75-10682
	p0044 N75-10683
N00014-66-C-0241	p0029 N75-10641
N00014-68-A-0091	p0032 N75-15141
N00014-68-A-0196-0009	p0032 175-15147
N00014-69-A-0141-0008	p0063 A75-10075
N00014-03-A-0141-0008	p0003 A75-10075
N00014-69-A-0407	p0026 A75-17201
N000 14-03-A-0407	
N00017-72-G-4401	p0048 A75-17181
N00017-72-G-4401 N00123-73-2352	p0056 N75-10425
NEOD24 70 C 0224	p0055 A75-17199
N60921-70-C-0221	p0042 A75-17121
USBR-14-06-D-7358	p0002 A75-17143
USGS-14-08-001-13377	p0069 A75-17230
W-7405-ENG-48	p0071 N75-11311
W04 040	p0022 N75-13458
WRK-213	p0031 N75-14239
641-14-07-50-72	p0051 N75-13371
915-17-00-00-72	n0081 N75-14358


D-2

REPORT/ACCESSION NUMBER INDEX

Earth Resources/A Continuing Bibliography (Issue 5)

OCTOBER 1975

Typical Report /Accession Number **Index Listing**



Listings in this index are arranged alphanumerically by report number. The page number indicates the page on which the citation is located. The accession number denotes the number by which the citation is identified. An asterisk (*) indicates that the item is a NASA report. A pound sign (#) indicates that the item is available on microfiche. A plus sign (+) indicates a document that cannot be microfiched but for which one-to-one facsimile is available.

AD-781138/3	p0029	N75-10524	#
AD-782883	p0006	N75-10577	#
AD-784104	p0072	N75-11454	#
AD-784162	p0071	N75-11450	#
AD-784170	p0057	N75-11453	#
AD-784373	p0056	N75-10425	#
AD-784421	p0029	N75-10641	#
AD-785585	p0021	N75-12459	#
AD-785698	p0030	N75-13375	H
AD-785838	p0032	N75-15147	Ħ
AD-785869	p0032	N75-15141	#
AD-785877	p0061	N75-14257	#
AD-785882	p0032	N75-15143	Ħ
AD-785886	p0061	N75-14255	#
AD-785932	p0009	N75-14258	#
AD-785967	p0010	N75-15146	#
AD-785998	p0081	N75-14259	#
AD-788028	p0022	N75-13419	#
AD-786418	p0032	N75-14335	Ħ
AD-786501	p0073	N75-14870	Ħ
AD-787281	p0032	N75-15237	#
AFAL-TR-74-98	p0022	N75-13419	#
AFCRL-TR-74-0317	p0032	N75-14335	#
AFOSR-74-0962TR	p0029	N75-10524	#
AIAA PAPER 74-1215	p0075	A75-12030*	#
AIAA PAPER 74-1216	00063		#
AIAA PAPER 74-1219	p0001	A75-12033	#
AIAA PAPER 74-1220	00041	A75-12034*	#
· AIAA PAPER 74-1221	p0063	A75-12035*	ï
AIAA PAPER 74-1224	p0064	A75-12036*	#
AIAA PAPER 74-1225	p0011	A75-12037*	#
			-
AIDJEX-74-25	p0045	N75-14262	Ħ
APL-TG-1226	p0056	N75-10425	Ħ
AR-29	p0022	N75-13369	#
BMFT-FB-W-74-04	p0024	N75-14331	#
BMPR-14	p0007	N75-12402*	#
BRL-CR-159	p0006	N75-10577	Ħ
BULL-161	p0031	N75-13454	#.
CONF-740106-4	-0017	N75-10638	
CONF-740406-16	p0017	N75-10638	#.

CONF-740415-1	p0016 N	175-10469 #		
CRC-PUBL-6	p0043 N	175-10571* #		
CUNY-43	n0046 N	J75.14793* #		***************************************
•		1		
D-171	p0074 N	175-15136 * #		
DGS-209	p0032 N	175-14335 #		
DMAAC-TP-74-002	p0032 N	175-15147 #		***************************************
	•	~!		***************************************
DOC-735D4249 DOC-745D4205	p0077 N	175-10548* # 175-10546* #		
DOC-74SD4217	p0077 N	l75-10547* ∦¦		
DOC-74SD4236	p0077 N	ł75-10568* #		
DREO-TN-74-18	p0021 N	175-12467 #	E75-10040	***************************************
EO-126	-0007 8		E75-10041	***************************************
•		- (E75-10042	
EOD2876	p0058 N	175-12279 * #	E75-10044	***************************************
ERIM-101700-17-L	p0020 8	475-12391* #	E75-10045	
ERIM-101800-14-P	p0022 1	175-13343* #	E75-10047	***************************************
ERIM-101900-40-L	p0057 f	N75-10527* #	E75-10048	
ERIM-101900-42-L ERIM-101900-44-L	n0056 1	175-12409 #	E75-10050	
ERIM-102000-27-L	n0036 N	175-12392* #.	F75-10051	***************************************
ERIM-102100-11-L	DO050 N	175-13348* #	E75-10053	
ERIM-102200-14-L	p0025 N	175-15114°#	E75-10054	,,
ERIM-102300-8-L	p0050 I	N75-12408* #	E/5-10055	***************************************
ERIM-102300-12-L	p0052 /	# # N75-15112	E75-10056	***************************************
ERIM-102300-13-L	p0052	V75-15111* #i	E75-10057	
ERIM-103300-35-L	p0020 /	175-12399* #	E75-10058	***************************************
ERIM-103300-37-L ERIM-103300-40-L	DO020 F	175-12404* #!	E75-10059	
ERIM-107400-2-F	n0073 N	175-15131* #	E75-10061	
ERIM-190100-30-T	p0058 f	N75-12417* #.	E75-10062	
ERIM-193500-5-X	p0070 f	N75-10564° #	E75-10063	***************************************
ERIM-196200-5-T	p0031	N75-14263 #	E75-10064	***************************************
ERIM-196200-11-T	p0031	N75-14281 #	E75-10065	***************************************
ÉSL-SR-132	20072	J75.12418* #	575-10066	
E3E-3N-132	p0072 1	475-12-16 #·	E75-10068	
ESL-TM-456	p0058 8	N75-12422* #	E75-10069	
			E75-10070	
ESRO-SP-100	p0081 /	N75-14216 #	E75-10071	
534 10720	-0056	130 10530\$ #	E75-10072	
E74-10730			676-100/3	
E74-10800			E75-10075	
E74-10801	p0042 f	N75-10529* #	E75-10076	
E74-10802	p0034 f	N75-10530* #		
E74-10803	p0057	175-10531°#	E75-10078	
E74-10804	P0070 F	N75-10532* #		
E74-10805	p0049 F	N75-11409*#		
E74-10807	p0077 1	175-10534* #		
,E74-10808	p0016 f	N75-10535* #		
E74-10809	p0034 f	N75-10536* #		*.
E74-10810	p0034 f	N75-10537* #		
E74-10811	p0016 f	N/5-10538* #		
E74-10812	n0049 F	475-10539" #1 N75-10540* #1	E75-10087	
E74-10815	p0049	175-10541* #		***************************************
E74-10816	p0049 f	475-10542* #,		
E74-10817	p0006 f	N75-10543* #	E75-10091	
E74-10818	p0006 I	N75-10544* #		
E75-10001	p0078	1/5-11406" # 176-12200# #		
E75-10002	p0035 I	N75-11407* #		
E75-10005	p0035 I	N75-11408* #		
E75-10006	p0020 f	N75-12391*#		
E75-10007	p0007 f	N75-13333* #		
E75-10008	p0036 f	N75-12392* #	E75-10099	
E75-10009	p0021	1/5-13334" # 175-12202# #		
E75-10010	p0020 1	N75-13335* #	E75-10101	
E75-10012	p0006 f	N75-12394* #	E75-10103	
E75-10013	p0059 f	√75-13336* #	E75-10104	
E75-10014	p0030 f	N75-12395* #		
E75-10015	p0021 f	N75-13337° #		***************************************
E75-10016	p0001 /	1/5-151U1" # J75.1220E* #		
E75-10017	p0020 I	175-12390°#	E15-10109	•
E75-10019	p0020	175-12398* #	FHWA-RD-1	4-9
E75-10020			FHWA-RD-7	4-27
E75-10021			ľ	
E75-10022			FOA-2-C-25	77-E1
E75-10023			FETC UT AN	790.72
E75-10024				-780-72 -0158-74
E75-10025				-0160-74
				-

	•	OC	тов	ER 1975
ı	£75-10026		p0050	N75-13342* #
1	E75-10027		p0022	N75-13343* #
ı	E75-10028		p0050	N75-13344° #
. 1	E75-10029	***************************************	p0008	N75-13345* #
	E75-10030		p0050	N75-12401° #
ı	E75-10031	***************************************	p0045	N75-14185* #
	E75-10032 E75-10033		p0007 p0007	N75-12402* # N75-12403* #
	E75-10034	***************************************	p0020	N75-12404* #
	£75-10035	***************************************	p0007	N75-12405° #
	E75-10036		p0020	N75-12406* #
	E75-10037		p0036	N75-13346° #
	E75-10038	***************************************	p0044	N75-12407* #
	E75-10040		p0050 p0058	N75-12408* # N75-12409* #
	E75-10041	***************************************	p0072	N75-12410* #
	E75-10042 E75-10043	***************************************	p0072 p0058	N75-12411* # N75-12412* #
	E75-10044		p0059	N75-13347* #
	E75-10045		p0030	N75-12413* #
1	E75-10046 E75-10047		p0007 p0036	N75-12414* # N75-12415* #
1	E75-10048		p0050	N75-13348* #
ŀ	E75-10050		p0050	N75-13350* #
.	E75-10051 E75-10052		p0036 p0030	N75-13351* # N75-13352* #
1	E75-10053		p0050	N75-13353* #
1	E75-10054 E75-10055	,,	p0036	N75-13354* # N75-13355* #
	E75-10056		p0037 p0022	N75-13356* #
	E75-10057	***************************************	p0037	N75-13357* #
	E75-10058 E75-10059	***************************************	p0037 p0030	N75-13358* # N75-13359* #
	E75-10060	***************************************	p0037	N75-13360* #
ı	E75-10061 E75-10062		p0008	N75-13361* #
	E75-10062		p0008 p0008	N75-13362* # N75-13363* #
	E75-10064		p0030	N75-13364* #
'	E75-10065 E75-10066		p0050 p0072	N75-13365* # N75-13366* #
	£75-10067		p0045	N75-14186* #
	E75-10068		p0059	N75-14187* #
	E75-10069 E75-10070		p0037 p0059	N75-14188* # N75-14189* #
	E75-10071		p0037	N75-14190* #
	E75-10072 E75-10073		p0037 p0008	N75-14191* # N75-14192* #
	E75-10074		p0031	N75-14193* #
	E75-10075 E75-10076		p0051 p0037	N75-14194* # N75-14195* #
	E75-10077		p0008	N75-14196* #
	E75-10078		p0031	N75-14197* #
1	E75-10079 E75-10080		p0023 p0072	N75-14198* # N75-14199* #
	E75-10081		p0073	N75-14200° #
	E75-10082 E75-10083		p0061 p0023	N75-15102* # N75-14201* #
	E75-10084		p0008	N75-14202* #
	E75-10085		p0038	N75-15103* #
	E75-10086 E75-10087		p0024 p0009	N75-15104* # N75-15105* #
	E75-10088		p0038	N75-15106* #
	E75-10089 E75-10090	***************************************	p0038 p0051	N75-15107* # N75-15108* #
1	E75-10091		p0024	N75-15109* #
	E75-10092 E75-10093		p0024 p0052	N75-15110* # N75-15111* #
	E75-10093		p0052	N75-15111* #
	E75-10095		p0073	N75-15113* #
	E75-10096 E75-10097		p0025	N75-15114* # N75-15115* #
	E75-10098	***************************************	p0046	N75_15118* #
İ	E75-10099 E75-10100		p0025 p0025	N75-15117* # N75-15118* #
	E75-10100		p0073	N75-15119* #
1	E75-10102		p0038	N75-15120* #
Ì	E75-10103 E75-10104		p0025 p0073	N75-15122° #
	E75-10105		p0025	N75-15123* #
1	E75-10106 E75-10108		p0024 p0039	N75-15100° # N75-15124° #
1	£75-10109		p0010	N75-15125* #
	FHWA-RD-7 FHWA-RD-7	4-9 4-27	p0031 p0031	N75-14263 # N75-14261 #
	FOA-2-C-25	77-E1	p0072	N75-12283 #

p0081 N75-14257 # p0081 N75-14259 # p0071 N75-11450 #

REPORT/ACCESSION NUMBER INDEX

		REPORT ACCESSION	VUIVIBER INDEX		
FSTC-HT-23-0276-74	00057 N75.11453 #	NASA-CR-140674	00070 N75-10564* # I	NASA-CR-141269	00080 N75-14209* #
FSTC-HT-23-1170-73	p0032 N75-15237 #	NASA-CR-140687		NASA-CR-141270	p0080 N75-14214* #
FSTC-HT-23-1270-73	p0061 N75-14255 #	NASA-CR-140688		NASA-CR-141283	p0074 N75-15138* #
FSTC-HT-23-1296-73	pQQQ9 N75-14258 #	NASA-CR-140711		NASA-CR-141291	p0073 N75-15113* #
FSTC-HT-23-1297-73	p0010 N75-15146 #	NASA-CR-140712		NASA-CR-141292 NASA-CR-141293	
FSTC-HT-23-1558-73	p0032 N/5-15143 #	NASA-CR-140725		NASA-CR-141294	p0046 N75-15116* #
GARP-SPEC-REPT-13	p0043 N75-10674 #	NASA-CR-140726	p0045 N75-14185* #	NASA-CR-141295	p0025 N75-15117* #
	·	NASA-CR-140727	p0035 N75-11408* #	NASA-CR-141296	p0025 N75-15118* #
GPO-25-314	p0077 N75-10617 #	NASA-CR-140728		NASA-CR-141297	p0073 N75-15119* #
GPO-40-560GPO-41-665	00078 N75-13010 #	NASA-CR-140731		NASA-CR-141298 NASA-CR-141304	n0025 N75-15121* #
Gr 0-41-003	poore 1175 12416 #	NASA-CR-140732 NASA-CR-140733	20020 N75-1239/* #	NASA-CR-141305	p0073 N75-15122* #
IAF PAPER 74-116		NASA-CR-140734	p0020 N75-12399* #	NASA-CR-141306	p0025 N75-15123* #
IAF PAPER 74-139	p0011 A75-13757 #	NASA-CR-140735	p0020 N75-12400* #	NASA-CR-141314 NASA-CR-141372	p0025 N75-15126* #
IAF PAPER 74-140	00053 A75-13758 #	.NASA-CR-140736	p0030 N75-13338* #	NASA-CR-141372	
1AF PAPER 74-147		NASA-CR-140737 NASA-CR-140738	00021 N/5-13339* #	NASA-CR-141477	p0074 N75-15333* #
IAF PAPER 74-156	p0065 A75-13765	NASA-CR-140739	p0050 N75-13342* #	NASA-CR-141685	
IAF PAPER 74-165		NASA-CR-140740	p0022 N75-13343* #		
ISBN-0-642-00853-1	-0021 N75 12454 #	NASA-CR-140741		NASA-NEWS-RELEASE-74-329 .	p00/9 N/5-133/0*#
ISBN-92-63-10368-2		NASA-CR-140754 NASA-CR-140762		NASA-SP-356	p0077 N75-10549* #
		NASA-CR-140763			•
JPL-SP-43-14	p0025 N75-15126* #	NASA-CR-140771		NASA-TM-X-58118	p0051 N75-13371* #
JPRS-63231	-0020 NZE 10960 #	NASA-CR-140772	p0007 N75-12405* #	NASA-TM-X-58124 NASA-TM-X-58154-VOL-1	
JPRS-63621	n0031 N75-13498 #	NASA-CR-140773	p0020 N75-12406* #	NASA-TM-X-62399	
		NASA-CR-140774	n0044 N75-12407* #	NASA-TM-X-62402	p0006 N75-12277* #
JSC-08457	p0051 N75-13371* #	NASA-CR-140776	p0050 N75-12408* #	NASA-TM-X-64881	p0078 N75-10967* #
JSC-09275-VOL-1	p0081 N75-14358* #	NASA-CR-140777	p0058 N75-12409* #	NASA-TM-X-64891 NASA-TM-X-70394	
LEC-3949	p0057 N75-10570* #	NASA-CR-140778		NASA-TM-X-70394	
		NASA-CR-140779 NASA-CR-140780		NASA-TM-X-70798	p0039 N75-15133* #
LWL-CR-03P68B	p0021 N75-12459 #	NASA-CR-140780	p0059 N75-13347* #	NASA-TM-X-70799	p0039 N75-15132* #
M-130	-0000 NOE 10070# #	NASA-CR-140782	p0030 N75-12413* #	NASA-TM-X-72029 NASA-TM-X-72062	
M-130	DOOSS 1479-13375. #	NASA-CR-140783		NASA-TM-X-72062	
MPR-10	p0007 N75-12414* #	NASA-CR-140784	00078 N75-12415* #	NASA-TM-X-72076	p0021 N75-13337* #
MPR-11	p0050 N75-13350* #	NASA-CR-140809		NASA-TM-X-72629	p0044 N75-12172* #
MPR-12	p0008 N75-14202* #	NASA-CR-140810	p0077 N75-10547* #	NASA-TR-R-431	-0000 NTE 100708 #
MPR-13	00010 N/5-15125; #	NASA-CR-140816	p0078 N75-11549* #	NASA-1R-R-431	DUU22 N/5-133/2-#
MPR-17	p0035 -N75-11408* #	NASA-CR-140865 NASA-CR-140898	.p000/ N/5-12421* # .p0039 N75-15124* #	NASA-TT-F-798	
MPR-18	p0031 N75-14193* #	NASA-CR-140899	p0010 N75-15125* #	NASA-TT-F-16000	
MPR-19	p0039 N75-15124* #	NASA-CR-140950	p0050 N75-13348* #	NASA-TT-F-16006	
MR-12	50028 N75.15106* #	NASA-CR-140952	p0050 N75-13350* #	NASA-TT-F-16021	n0070 N75-10964* #
MR-13		NASA-CR-140953 NASA-CR-140954	DOGO N75-13351* #	NASA-TT-F-16108	
MR-14	p0038 N75-15106* #	NASA-CR-140959	p0050 N75-13353* #	NASA-TT-F-16151	p0009 N75-15097* #
MR-15	p0078 N75-12019* #	NASA-CR-140962	p0036 N75-13354* #	NOT (IDOF 34 10	-0044 NZE 10001 #
MR-15	p0038 N75-15106* #	NASA-CR-140963	p0037 N75-13355* #	NSF/IDOE-74-18 NSF/IDOE-74-19	
MR-17	p0038 N75-15107* #	NASA-CR-140964 NASA-CR-140965	n0037 N75-13350* #	NSF/IDOE-74-20	p0044 N75-10683 #
		NASA-CR-140966	p0037 N75-13358* #		
MUJWRRC-51	p0051 N75-14260 #	NASA-CR-140999	p0030 N75-13359* #	NWAG-RESEARCH-CONTRIB-266	p0032 N75-15141 #
NASA-CR-2458	n0045 N75-13496* #	NASA-CR-141000	p0037 N75-13360* #	OWRR-B-036-RI(4)	p0056 N75-10202 #
NASA-CR-2459	p0017 N75-10545* #	NASA-CR-141001 NASA-CR-141002	00008 N75-13361* #		
NASA-CR-2475	p0058 N75-12422* #	NASA-CR-141003	p0008 N75-13363* #	P-412-10	p0073 N75-14200* #
NASA-CR-120540	p0021 N75-12546* #	NASA-CR-141004	p0030 N75-13364* #	PB-233432/4	p0056 N75-10202 #
NASA-CR-134288 NASA-CR-134289	DOODS N75-12279* #	NASA-CR-141005 NASA-CR-141006	p0050 N75-13365* #	PB-233847/3	
NASA-CR-134294	p0058 N75-12420* #	NASA-CR-141006	00045 N75-14186* #	PB-234057/8	p0044 N75-10681 #
NASA-CR-134453	p0025 N75-15199* #	NASA-CR-141022	p0059 N75-14187* #	PB-234058/6	p0044 N75-10682 #
NASA-CR-137469 NASA-CR-137470	p0043 N75-10673* #	NASA-CR-141044	p0078 N75-12424* #	PB-234059/4 PB-234065/1	
NASA-CR-137475	n0043 N75-105/1 #	NASA-CR-141045	p0020 N75-12425* #	PB-235298/7	p0045 N75-14262 #
NASA-CR-137476	p0043 N75-10571* #	NASA-CR-141057	p0052 N75-15112* #	PB-235991/7	p0031 N75-14261 #
NASA-CR-139145	p0044 N75-11411* #	NASA-CR-141092	p0023 N75-14198* #	PB-235992/5	p0031 N75-14263 #
NASA-CR-139163 NASA-CR-139182	p0073 N/5-15131* #	NASA-CR-141093	p0031 N75-14197* #	PB-236041/0	p0051 N75-14260 #
NASA-CR-139162	00056 N75-10526* #	NASA-CR-141094		PHL-1974-23	p0070 N75-10573 #
NASA-CR-140334	p0072 N75-12418* #		p0037 N75-14195* # p0051 N75-14194* #		•
NASA-CR-140345	p0071 N75-11447* #	NASA-CR-141097	p0031 N75-14193* #	PR-1	p0077 N75-10534* #
NASA-CR-140357 NASA-CR-140493	DO057 N75-10527*#	NASA-CR-141098		PR-17	p0025 N75-15121* # p0008 N75-13362* #
NASA-CR-140494	p0029 N75-10528* #	NASA-CR-141099 NASA-CR-141100		PR-19	p0030 N75-13364* #
NASA-CR-140495	p0042 N75-10529* #	NASA-CR-141100			
NASA-CR-140496	p0034 N75-10530* #	NASA-CR-141102	p0037 N75-14188* #	QPR-4	p0044 N75-12407* #
NASA-CR-140497 NASA-CR-140498	p005/ N/5-10531* #	NASA-CR-141104		QPR-4	p0038 N75-15103* # p0020 N75-12391* #
NASA-CR-140499		NASA-CR-141105		QPR-6	p0020 N75-12400* #
NASA-CR-140500		NASA-CR-141107			p0020 N75-12406* #
NASA-CR-140501	p0077 N75-10534* #	NASA-CR-141108	p0008 N75-14202* #	QPR-7	p0073 N75-14200* #
NASA-CR-140502 NASA-CR-140503		NASA-CR-141130		QPR-7	p0025 N75-15117* #
NASA-CR-140504		NASA-CR-141149		QR-5	p0079 N75-13373 #
NASA-CR-140505	p0016 N75-10538* #	NASA-CR-141150			
NASA-CR-140569	p0049 N75-10539* #	NASA-CR-141152	p0038 N75-15106*#	REPT-3	p0018 N75-11476 #
NASA-CR-140570 NASA-CR-140586		NASA-CR-141153			p0022 N75-13437 # p0030 N75-13432 #
NASA-CR-140616		NASA-CR-141154 NASA-CR-141155		REPT-13	p0032 N75-15228 #
NASA-CR-140617	p0049 N75-10541* #	NASA-CR-141155		REPT-74-8	p0072 N75-12410* #
NASA-CR-140618	p0049 N75-10542* #	NASA-CR-141157	p0052 N75-15111* #	REPT-74-2001-5	
NASA-CR-140619 NASA-CR-140620		NASA-CR-141191	p0080 N75-14213* #	REPT-74-2001-6	
NASA-CR-140647		NASA-CR-141192 NASA-CR-141193	DOORD N75-14215* #	REPT-74-2002-10-VOL-3-PT-1	
NASA-CR-140650		NASA-CR-141194	p0079 N75-14206* #	REPT-74-2002-10-VOL-3-PT-2	
NASA-CR-140651		NASA-CR-141262	p0079 N75-14207* #	REPT-74-2002-10-VOL-3-PT-3	p0079 N75-14207* #
NASA-CR-140654		NASA-CR-141263	p0080 N75-14208* #	REPT-74-2002-10-VOL-4	p0080 N75-14208* #
NASA-CR-140655		NASA-CR-141264 NASA-CR-141265	00079 N75-14212*#	REPT-74-2002-10-VQL-5	p0080 N75-14209* #
NASA-CR-140656	p0020 N75-12393* #	NASA-CR-141266	p0080 N75-14210* #	REPT-74-2002-10-VOL-7	p0080 N75-14212* #
NASA-CR-140658		NASA-CR-141267	p0080 N75-14211* #	REPT-74-2002-10-VOL-8	p0080 N75-14213* #
NASA-CR-140665	p0017 N75-10563* #	NASA-CR-141268	p0079 N75-14203* #	REPT-74-2002-10-VOL-9	p0080 N75-14214* #

	-0000	N75-14215* #
REPT-74-2002-10-VOL-10	p0080	
REPT-74-2002-10-1-VOL-1	p0079	N75-14203* #
REPT-74-2002-10-2-VOL-2	D0079	N75-14204* #
REPT-128	p0050	N75-13344* #
REPT-275-025-FP2	p0074	N75-15333* #
REPT-275-025-018	p0071	N75-11447* #
1)EF1-273-023-010	p0071	1173-11777
		WAT 405008 #
RSC-53	p0017	N75-10563* #
RSL-TR-254-2	00046	N75-14793* #
100-111-204-2	pooto	
SAO-SPECIAL-REPT-361	p0030	N75-13340* #
SAO-403-004	20030	N75-13340* #
440-403-004	pooso	1075-10540 #
SAPR-2	p0078	N75-11414* #
		-
\$LA-73-5931	-0016	N75-10469 #
SLA-73-5931	poore	N75-10469 #
\$R-18	n0032	N75-14335 #
411 10	p000-	
\$13-116	p0017	N75-10545* #
TDCK-65200	-0070	N75-10573 #
TUCK-05200	puo/u	N/5-103/3 #
TI-UI-998200-01-F	p0072	N75-11454 #
., .,		
3.0.00004		War 40000 "
TID-26594	p0021	N75-13063 #
TP-19	-0029	N75-10625 #
1F-19	p0023	1475-10025 H
TR-EE/CS-74-5	p0016	N75-10538* #
TR-EE/CS-74-5	p0016	N75-10538* #
·	•	
TR-74-1	p0035	N75-11407* #
TR-74-1	p0035 p0036	N75-11407* # N75-13346* #
TR-74-1	p0035 p0036	N75-11407* #
TR-74-1	p0035 p0036 p0007	N75-11407* # N75-13346* # N75-12405* #
TR-74-1	p0035 p0036 p0007 p0007	N75-11407* # N75-13346* # N75-12405* # N75-13333* #
TR-74-1	p0035 p0036 p0007	N75-11407* # N75-13346* # N75-12405* #
TR-74-1	p0035 p0036 p0007 p0007	N75-11407* # N75-13346* # N75-12405* # N75-13333* # N75-14192* #
TR-74-1	p0035 p0036 p0007 p0007 p0008	N75-11407* # N75-13346* # N75-12405* # N75-13333* # N75-14192* #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20	p0035 p0036 p0007 p0007	N75-11407* # N75-13346* # N75-12405* # N75-13333* #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004	p0035 p0036 p0007 p0007 p0008	N75-11407* # N75-13346* # N75-12405* # N75-13333* # N75-14192* #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488	p0035 p0036 p0007 p0007 p0008 p0058	N75-11407* # N75-13346* # N75-12405* # N75-13333* # N75-14192* # N75-12279* #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004	p0035 p0036 p0007 p0007 p0008	N75-11407* # N75-13346* # N75-13333* # N75-14192* # N75-12279* # N75-11311 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488	p0035 p0036 p0007 p0007 p0008 p0058	N75-11407* # N75-13346* # N75-12405* # N75-13333* # N75-14192* # N75-12279* #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565	p0035 p0036 p0007 p0008 p0058 p0058 p0071 p0022	N75-11407° # N75-13346° # N75-12405° # N75-13333° # N75-14192° # N75-12279° # N75-11311 # N75-13458 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022	N75-11407* # N75-13346* # N75-12405* # N75-13333* # N75-14192* # N75-12279* # N75-13458 # N75-10638 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565	p0035 p0036 p0007 p0008 p0058 p0058 p0071 p0022	N75-11407* # N75-13346* # N75-12405* # N75-13333* # N75-14192* # N75-12279* # N75-13458 # N75-10638 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022	N75-11407* # N75-13346* # N75-12405* # N75-13333* # N75-14192* # N75-12279* # N75-13458 # N75-10638 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-755644-REV-2	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022	N75-11407* # N75-13346* # N75-13405* # N75-13333* # N75-14192* # N75-1279* # N75-13458 # N75-10638 # N75-15202 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022	N75-11407* # N75-13346* # N75-12405* # N75-13333* # N75-14192* # N75-12279* # N75-13458 # N75-10638 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-75644-REV-2 USAMC-ITC-02-08-73-018	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0025	N75-11407* # N75-13346* # N75-12406* # N75-13333* # N75-14192* # N75-11311 # N75-13458 # N75-15202 # N75-14870 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-755644-REV-2	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0025	N75-11407* # N75-13346* # N75-12406* # N75-13333* # N75-14192* # N75-11311 # N75-13458 # N75-15202 # N75-14870 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-75644-REV-2 USAMC-ITC-02-08-73-018	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0025	N75-11407° # N75-13346° # N75-12405° # N75-13333° # N75-14192° # N75-11311 # N75-13458 # N75-15202 # N75-14870 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75566 UCRL-75644-REV-2 USAMC-ITC-02-08-73-018 USCG-D-87-74	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0025 p0073	N75-11407* # N75-13346* # N75-12405* # N75-12405* # N75-12279* # N75-13111 # N75-13458 # N75-15202 # N75-14870 # N75-11454 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-75644-REV-2 USAMC-ITC-02-08-73-018	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0025 p0073	N75-11407* # N75-13346* # N75-12406* # N75-13333* # N75-14192* # N75-11311 # N75-13458 # N75-15202 # N75-14870 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-75564 UCRL-75644-REV-2 USAMC-ITC-02-08-73-018 USCG-D-87-74 USGS-D0-74-010	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0073 p0073	N75-11407* # N75-13346* # N75-12405* # N75-12405* # N75-12279* # N75-1311 # N75-13458 # N75-16638 # N75-15202 # N75-14870 # N75-11454 # N75-13373 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-75564 UCRL-75644-REV-2 USAMC-ITC-02-08-73-018 USCG-D-87-74 USGS-D0-74-010	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0073 p0073	N75-11407* # N75-13346* # N75-12405* # N75-12405* # N75-12279* # N75-1311 # N75-13458 # N75-16638 # N75-15202 # N75-14870 # N75-11454 # N75-13373 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75566 UCRL-75644-REV-2 USAMC-ITC-02-08-73-018 USCG-D-87-74	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0025 p0073	N75-11407* # N75-13346* # N75-12405* # N75-12405* # N75-12279* # N75-1311 # N75-13458 # N75-16638 # N75-15202 # N75-14870 # N75-11454 # N75-13373 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-75644-REV-2 USAMC-ITC-02-08-73-018 USCG-D-87-74 USGS-D0-74-010 WHO1-74-59	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0025 p0073 p0072 p0079	N75-11407* # N75-13346* # N75-12405* # N75-13333* # N75-13279* # N75-1311 # N75-13458 # N75-16638 # N75-15202 # N75-14870 # N75-13373 # N75-10641 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-75564 UCRL-75644-REV-2 USAMC-ITC-02-08-73-018 USCG-D-87-74 USGS-D0-74-010	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0025 p0073 p0072 p0079	N75-11407* # N75-13346* # N75-12405* # N75-12405* # N75-12279* # N75-1311 # N75-13458 # N75-16638 # N75-15202 # N75-14870 # N75-11454 # N75-13373 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-755644-REV-2 USAMC-ITC-02-08-73-018 USCG-D-87-74 USGS-D0-74-010 WH01-74-59 WM0-368	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0025 p0073 p0072 p0079	N75-11407* # N75-13346* # N75-12405* # N75-13333* # N75-13279* # N75-1311 # N75-13458 # N75-16638 # N75-15202 # N75-14870 # N75-13373 # N75-10641 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-755644-REV-2 USAMC-ITC-02-08-73-018 USCG-D-87-74 USGS-D0-74-010 WH01-74-59 WM0-368	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0073 p0072 p0079 p0079	N75-11407* # N75-13346* # N75-12405* # N75-13333* # N75-14192* # N75-13111 # N75-13458 # N75-16638 # N75-15202 # N75-14870 # N75-13373 # N75-10641 # N75-10641 # N75-11476 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-75644-REV-2 USAMC-ITC-02-08-73-018 USCG-D-87-74 USGS-D0-74-010 WHO1-74-59	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0073 p0072 p0079 p0079	N75-11407* # N75-13346* # N75-12405* # N75-13333* # N75-13279* # N75-1311 # N75-13458 # N75-16638 # N75-15202 # N75-14870 # N75-13373 # N75-10641 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-75564-REV-2 USAMC-ITC-02-08-73-018 USCG-D-87-74 USGS-D0-74-010 WH01-74-59 WM0-368 W74-09255	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0025 p0073 p0072 p0079 p0029 p0018	N75-11407* # N75-13346* # N75-12405* # N75-13333* # N75-14192* # N75-13279* # N75-13111 # N75-13458 # N75-15202 # N75-14870 # N75-11454 # N75-10641 # N75-11476 # N75-11476 #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-755644-REV-2 USAMC-ITC-02-08-73-018 USCG-D-87-74 USGS-D0-74-010 WH01-74-59 WM0-368 W74-09255 X-922-74-314	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0025 p0073 p0072 p0079 p0029 p0018 p0056	N75-11407* # N75-13346* # N75-12405* # N75-13333* * N75-14192* # N75-13111 # N75-13458 # N75-16638 # N75-15202 # N75-14870 # N75-10641 # N75-10641 # N75-10476 # N75-10476 # N75-11476 # N75-11312* #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-75564-REV-2 USAMC-ITC-02-08-73-018 USCG-D-87-74 USGS-D0-74-010 WH01-74-59 WM0-368 W74-09255	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0025 p0073 p0072 p0079 p0029 p0018	N75-11407* # N75-13346* # N75-12405* # N75-13333* # N75-14192* # N75-13111 # N75-13458 # N75-16638 # N75-15202 # N75-1454 # N75-10641 # N75-10641 # N75-10641 # N75-10642 # N75-11476 # N75-10420 # N75-15132* #
TR-74-1 TR-74-2 TR-239-17 TR-239-18 TR-239-20 TWP-72-004 UCID-16488 UCID-16565 UCRL-75556 UCRL-755644-REV-2 USAMC-ITC-02-08-73-018 USCG-D-87-74 USGS-D0-74-010 WH01-74-59 WM0-368 W74-09255 X-922-74-314	p0035 p0036 p0007 p0007 p0008 p0058 p0071 p0022 p0017 p0025 p0073 p0072 p0079 p0029 p0018 p0056	N75-11407* # N75-13346* # N75-12405* # N75-13333* * N75-14192* # N75-13111 # N75-13458 # N75-16638 # N75-15202 # N75-14870 # N75-10641 # N75-10641 # N75-10476 # N75-10476 # N75-11476 # N75-11312* #

1. Report No. NASA SP-7041 (05)	2. Government Accessi	on No.	3. Recipient's Catalog	No.	
4. Title and Subtitle EARTH RESOURCES			5. Report Date October 197	75	
A Continuing Bibliography	(Issue 05)	<u> </u>	6. Performing Organiz		
in continuing by a rography	(13340 0))				
7. Author(s)			8. Performing Organiza	ation Report No.	
· · · · · · · · · · · · · · · · · · ·			10. Work Unit No.		
9. Performing Organization Name and Address					
National Aeronautics and Space Administration Washington, D. C. 20546		tration	11. Contract or Grant No.		
and the second s			13. Type of Report and Period Covered		
12. Sponsoring Agency Name and Address					
		. [14. Sponsoring Agency	Code	
45.6		<u> </u>			
15. Supplementary Notes					
				•	
16. Abstract					
This bibliography lists 601 reports, articles, and other documents introduced into the NASA scientific and technical information system between January 1975 and March 1975. Emphasis is placed on the use					
of remote sensing an					
aircraft to survey a					
Subject matter is gr environmental change					
geology and mineral					
processing and distr					
and economic analysi	•	iis, Tiiseramorrea	cron una sone	,,,,	
	•				
		•			
·					
•					
17. Key Words (Suggested by Author(s))		18. Distribution Statement			
Bibliographies					
Earth Resources Program					
Remote Sensors Unclassified - Unlimited		ited			
19. Security Classif. (of this report)	20. Security Classif. (o	f this page)	21. No. of Pages	22. Price*	
Unclassified	Unclassif		150	\$4.00 HC	

PUBLIC COLLECTIONS OF NASA DOCUMENTS

DOMESTIC

NASA distributes its technical documents and bibliographic tools to eleven special libraries located in the organizations listed below. Each library is prepared to furnish the public such services as reference assistance, interlibrary loans, photocopy service, and assistance in obtaining copies of NASA documents for retention.

CALIFORNIA

University of California, Berkeley

COLORADO

University of Colorado, Boulder

DISTRICT OF COLUMBIA

Library of Congress

GEORGIA

Georgia Institute of Technology, Atlanta

ILLINOIS

The John Crerar Library, Chicago

MASSACHUSETTS

Massachusetts Institute of Technology, Cambridge

MISSOURI

Linda Hall Library, Kansas City

NEW YORK

Columbia University, New York

OKLAHOMA

University of Oklahoma, Bizzell Library

PENNSYLVANIA

Carnegie Library of Pittsburgh

WASHINGTON

University of Washington, Seattle

NASA publications (those indicated by an "*" following the accession number) are also received by the following public and free libraries:

CALIFORNIA

Los Angeles Public Library San Diego Public Library

COLORADO

Denver Public Library

CONNECTICUT

, Hartford Public Library

MARYLAND

Enoch Pratt Free Library, Baltimore

MASSACHUSETTS

Boston Public Library

MICHIGAN

Detroit Public Library

MINNESOTA

Minneapolis Public Library

MISSOURI

Kansas City Public Library

St. Louis Public Library

NEW JERSEY

Trenton Public Library

NEW YORK

Brooklyn Public Library

Buffalo and Erie County Public Library

Rochester Public Library

New York Public Library

OHIO

Akron Public Library

Cincinnati Public Library

Cleveland Public Library

Dayton Public Library

Toledo Public Library

TENNESSEE

Memphis Public Library

TEXAS

Dallas Public Library
Fort Worth Public Library

WASHINGTON
Seattle Public Library

WISCONSIN

Milwaukee Public Library

An extensive collection of NASA and NASA-sponsored documents and aerospace publications available to the public for reference purposes is maintained by the American Institute of Aeronautics and Astronautics, Technical Information Service, 750 Third Avenue, New York, New York 10017.

EUROPEAN

An extensive collection of NASA and NASA-sponsored publications is maintained by the British Library Lending Division, Boston Spa, Wetherby, Yorkshire, England. By virtue of arrangements other than with NASA, the British Library Lending Division also has available many of the non-NASA publications cited in *STAR*. European requesters may purchase facsimile copy of microfiche of NASA and NASA-sponsored documents, those identified by both the symbols "#" and "*", from: ESA - Space Documentation Service, European Space Agency, 114 Avenue Charles-de-Gaulle, 92522 Neuilly-sur-Seine, France.

OFFICIAL BUSINESS PENALTY FOR PRIVATE USE \$300

SPECIAL FOURTH CLASS MAIL Book



POSTMASTER:

If Undeliverable (Section 158 Postal Manual) Do Not Return

NASA CONTINUING BIBLIOGRAPHY SERIES

NUMBER	TITLE	FREQUENCY
NASA SP7011	AEROSPACE MEDICINE AND BIOLOGY Aviation medicine, space medicine, and space biology	Monthly
NASA SP-7037	AERONAUTICAL ENGINEERING Engineering, design, and operation of aircraft and aircraft components	Monthly
NASA SP7039	NASA PATENT ABSTRACTS BIBLIOGRAPHY NASA patents and applications for patent	Semiannually
NASA SP-7041	EARTH RESOURCES Remote sensing of earth resources by aircraft and spacecraft	Quarterly
NASA SP-7043	ENERGY Energy sources, solar energy, energy conversion, transport, and storage	Quarterly
NASA SP-7500	MANAGEMENT Program, contract, and personnel management, and management techniques	Annually

Details on the availability of these publications may be obtained from:

SCIENTIFIC AND TECHNICAL INFORMATION OFFICE

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

Washington, D.C. 20546